

LETTER TO THE EDITOR

LIST DO REDAKCJI

**SYNERGISTIC IMPACT OF EXERCISE AND GLP-1 RECEPTOR AGONIST  
TREATMENT: ADDRESSING METABOLIC SYNDROME, ABDOMINAL OBESITY,  
AND INFLAMMATION**

**SYNERGICZNY WPŁYW POŁĄCZENIA AKTYWNOŚCI FIZYCZNEJ I LECZENIA  
AGONISTAMI RECEPTORA GLP-1 NA NASILENIE ZESPOŁU  
METABOLICZNEGO, OTYŁOŚĆ BRZUSZNĄ I STAN ZAPALNY**

**Jakub Kuśmierski<sup>1(A,B,C,D,E,F)</sup>, Martyna Kuśmierska<sup>2(A,B,C,D,E,F)</sup>**

<sup>1</sup>Faculty of Medical Sciences in Zabrze, Medical University of Silesia in Katowice, Poland

<sup>2</sup>Independent Public Health Care Facility, Municipal Hospital Complex in Chorzów, Poland

Kuśmierski J, Kuśmierska M. Synergistic impact of exercise and GLP-1 receptor agonist treatment: addressing metabolic syndrome, abdominal obesity, and inflammation. Health Prob Civil. <https://doi.org/10.5114/hpc.2024.139680>

Tables: 0

Figures: 0

References: 5

Submitted: 2024 May 8

Accepted: 2024 May 14

**Address for correspondence / Adres korespondencyjny:** Jakub Kuśmierski, Faculty of Medical Sciences in Zabrze, Medical University of Silesia in Katowice, Poniatowskiego Street 15, 40-055 Katowice, Poland, e-mail: j.kusmierski@outlook.com, phone: +48 32 271 72 17

ORCID: Jakub Kuśmierski <https://orcid.org/0009-0003-2031-3071>, Martyna Kuśmierska <https://orcid.org/0009-0005-5895-2471>

Copyright: © John Paul II University in Biała Podlaska, Jakub Kuśmierski, Martyna Kuśmierska. This is an Open Access journal, all articles are distributed under the terms of the Creative Commons AttributionNonCommercialShareAlike 4.0 International (CC BY-NC-SA 4.0) License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited and states its license.

**Keywords:** GLP-1 receptor agonists, abdominal obesity, metabolic syndrome, physical activity, inflammation

**Słowa kluczowe:** agoniści GLP-1, otyłość brzuszna, zespół metaboliczny, aktywność fizyczna, zapalenie

**Dear Editor,**

We are eager to share the findings from a recent article that we had the pleasure of reading, which we believe holds profound significance. Authored by Sandsdal et al. [1], this randomized controlled clinical trial delves into the impact of a combination of exercise and GLP-

1 receptor agonist treatment on the severity of metabolic syndrome, abdominal obesity, and inflammation.

The escalating prevalence of overweight and obesity has emerged as a prominent catalyst for the surge in chronic, non-communicable illnesses worldwide. In contemporary understanding, overweight and obesity are regarded as chronic ailments. Beyond the physical health implications, individuals grappling with these conditions often face psychological and social challenges. They contend with societal biases, experience discrimination in personal and professional spheres, grapple with diminished self-esteem, and are susceptible to depression [2,3].

During the trial [1], over an 8-week period, 195 adults diagnosed with obesity but not diabetes underwent a low-calorie diet regimen of 800 kcal/day, resulting in a 12% reduction in body weight. Following this, participants were randomly assigned to one of four treatment arms for a year: placebo, moderate-to-vigorous exercise (requiring a minimum of 150 minutes per week of moderate-intensity activity, 75 minutes per week of vigorous-intensity aerobic exercise, or a proportional blend of both), the GLP-1 RA liraglutide at a dose of 3.0 mg/day, or a combined approach involving both exercise and liraglutide.

The findings of this study are highly intriguing. Following diet-induced weight loss, the severity of metabolic syndrome, as indicated by the Metabolic Syndrome Severity Z-score (MetS-Z), was reduced; a reduction that persisted in both the placebo and exercise groups after one year. Moreover, treatment with liraglutide alone and in combination with exercise led to further decreases in MetS-Z compared to the placebo group. In addition, abdominal fat percentage decreased by 2.6, 2.8, and 6.1 percentage points in the exercise, liraglutide, and

combination groups, respectively, compared to the placebo group. Notably, levels of hsCRP decreased only in the combination group when compared with the placebo group.

We find these results particularly captivating, given the multifaceted effects observed. Each of these outcomes holds significant promise in the context of reducing cardiometabolic risk. The combined reduction in metabolic syndrome severity, abdominal obesity, and inflammation suggests a comprehensive approach to addressing underlying factors contributing to cardiovascular and metabolic health [4]. It's worth highlighting that the use of glucagon-like peptide-1 receptor agonists (GLP-1 RAs) in isolation has demonstrated both safety and efficacy. Beyond facilitating weight loss, these agents have shown promise in mitigating risk factors associated with cardiovascular disease. Notably, they have been linked not only to improvements in body weight but also to favorable alterations in blood pressure and lipid profiles [5]. This holistic impact underscores the potential of such interventions to yield substantial benefits in mitigating overall cardiometabolic risk.

## References:

1. Sandsdal RM, Juhl CR, Jensen SBK, Lundgren JR, Janus C, Blond MB, et al. Combination of exercise and GLP-1 receptor agonist treatment reduces severity of metabolic syndrome, abdominal obesity, and inflammation: a randomized controlled trial. *Cardiovasc Diabetol*. 2023; 22(1): 41. <https://doi.org/10.1186/s12933-023-01765-z>
2. Purnell JQ. Definitions, classification, and epidemiology of obesity. In: Feingold KR, Anawalt B, Blackman MR, Boyce A, Chrousos G, Corpas E, et al., editors. *Endotext*. South Dartmouth (MA): MDText.com, Inc.; 2000.

3. Rubino F, Puhl RM, Cummings DE, Eckel RH, Ryan DH, Mechanick JI, et al. Joint international consensus statement for ending stigma of obesity. *Nat Med.* 2020; 26(4): 485-497. <https://doi.org/10.1038/s41591-020-0803-x>
4. Khanna D, Welch BS, Rehman A. *Pathophysiology of obesity.* Treasure Island (FL): StatPearls Publishing; 2024.
5. Ansari HUH, Qazi SU, Sajid F, Altaf Z, Ghazanfar S, Naveed N, et al. Efficacy and safety of glucagon-like peptide-1 receptor agonists on body weight and cardiometabolic parameters in individuals with obesity and without diabetes: a systematic review and meta-analysis. *Endocr Pract.* 2024; 30(2): 160-171. <https://doi.org/10.1016/j.eprac.2023.11.007>