



Ministerstwo Nauki
i Szkolnictwa Wyższego



Krajowy Naukowy
Ośrodek Wiodący
2012-2017



The 12th International Symposium

MOLECULAR BASIS OF PATHOLOGY AND THERAPY IN NEUROLOGICAL DISORDERS

November 20-21, 2014

Warsaw, Poland

ORGANIZERS

Mossakowski Medical Research Centre, Polish Academy of Sciences,
Polish Academy of Sciences, Division V Medical Sciences,
Committee of Physiological and Pharmacological Sciences
and Committee of Neurobiology

SCIENTIFIC COMMITTEE


Elzbieta Salinska-Chairperson
Leonora Buzanska
Krystyna Domanska-Janik
Jerzy Lazarewicz
Andrzej W. Lipkowski
Barbara Lukomska
Lidia Struzynska
Barbara Zablocka
Teresa Zalewska
Magdalena Zielinska

MAZOWIECKI KLASTER PEPTYDOWY

Projekt współfinansowany przez Unię Europejską, ze środków Europejskiego Funduszu Rozwoju Regionalnego
w ramach Regionalnego Programu Operacyjnego Województwa Mazowieckiego 2007-2013.



SPONSORS:



SYMBIOS LIFE SCIENCES



The growth and regeneration of axons: an *in vitro* model for studying interactions in 3D

Gerardo-Nava J.¹, Hodde D.¹, Katona I.¹, Bozkurt A.², Grehl T.³, Steinbusch H.W.M.⁴, Weis J.¹, Brook G.A.¹

¹Institute of Neuropathology, Uniklinik RWTH Aachen, Aachen 52074, Germany

²Department of Plastic, Hand and Burns Surgery, Uniklinik RWTH Aachen University, Aachen, Germany

³Department of Neurology, Ruhr-University Bochum, BG-Kliniken Bergmannsheil GmbH, Bochum 44789, Germany

⁴Department of Translational Neuroscience, Maastricht University Medical Center, Maastricht, The Netherlands

Investigations of the factors influencing mammalian axonal growth and regeneration are currently addressed using a range of *in vitro* and *in vivo* models. The *in vitro* models often employ relatively simple 2D culture systems (or the more recent development of axon diodes in micro-fluidics [5,6]) and present the advantage of ready access, manipulation and visualization of the axons and the environment in which they are growing. The *in vivo* models employ relatively expensive laboratory animals and present the advantage of studying axon growth in a more natural 3D environment of supporting cells and extracellular matrix. Our earlier *in vitro* attempts to demonstrate the support of axon regeneration by 3D bio-engineered scaffolds involved the use of explanted dorsal root ganglia (DRG) obtained from adult rats [2,3]. This approach has been useful for demonstrating the influence of the scaffold on the properties of glial cell adhesion, proliferation, migration and process extension as well as on axon regeneration and orientation of growth. However, such studies were limited sensory neurons which are easy to dissect and can be maintained in culture for extended periods of time. However, patterns of DRG axonal growth do not necessarily reflect those that may be demonstrated would be expected by motor axons: differences in axonal growth between the motor and sensory neurons (in both *in vivo* and *in vitro* models) have been demonstrated by others in response to specific cellular and molecular cues, including differences when presented with environments containing a range of growth factors [1]. Such differences highlight the value of assays capable of evaluating neuron subtype-specific responses to their environment. Here, we describe the use of the *in vitro* organotypic spinal cord slice preparation for the investigation of motor axon regeneration from early post natal (P7-P9) rat pups and demonstrate the influence of scaffold microstructure on the trajectories the axons (for a complete description see Gerardo-Nava and colleagues [4]). To date, *in vitro* investigations of such motor axon-3D scaffold interactions remain poorly characterized, largely due to the difficulties

of culturing motor neurons (MN) for extended periods *in vitro*. The organotypic spinal cord slice preparation has the advantage of studying motoneurons in a relatively normal cellular niche of the ventral horn gray matter in which their survival and axonal regeneration can be maintained for many weeks or even months.

The rat pup spinal cords were rapidly isolated in ice-cold Gey's balanced salt solution and the dura mater and superficial blood vessels carefully removed. The lumbar spinal cord was sectioned transversely (350 µm thick) with a McIlwain tissue chopper and the slices transferred to Millicell membrane inserts where previously dissected ventral roots or biomaterials (fibrin hydrogel or orientated collagen scaffold, Matricel GmbH) could be carefully apposed to the ventral surface of the slices (as described by Vyas and colleagues [7]). The preparations were then maintained in an incubator at 37°C with 5% CO₂.

Myelin clearance within the explanted ventral roots was rapid with the events following a similar pattern to that already described for Wallerian degeneration *in vivo*. By 7-14 days in culture, axons regenerated for substantial distances along the explanted nerve root, with most axons following a trajectory within the root itself. The interactions between regenerating motor axons and Schwann cells revealed different stages of maturity from simple, initial axon-Schwann cell contacts, to the ensheathment and the formation of compact myelin surrounded by basal lamina. A similar sequence of events could also be seen within the longitudinally orientated porous framework of the collagen scaffold, however, these interactions took place between migrating Schwann cells (from the residual root stumps remaining at the ventral surface of the slice) and the regenerating motor axons which formed longitudinally orientated fascicles. Although similarly extensive Schwann cell migration and axon growth was also supported by the fibrin hydrogel, the trajectory of both cells and axons followed the surface of the hydrogel rather through it.

Although the organotypic spinal cord preparation is capable of reproducing many of the events taking place in PNS Wallerian degeneration with subsequent axonal regeneration, the model has clear limitations, including the lack of any functioning vasculature, an incomplete immune system and the absence of appropriate end organ targets for the regenerating motor axons. However, despite these limitations, the model affords a number of significant advantages over assays that attempt to employ cultures of freshly isolated embryonic motor neurons (or cells lines), or studies involving laboratory animals. These include the simplicity and ease of preparation, the length of time such preparations can be maintained and the financial and ethical benefits of reducing the number of animals required to perform such studies. The model thus represents, in our opinion, an excellent platform for inves-

tigating how regenerating motor axons interact with any 3D growth-promoting environment and can be used as a cost-effective means of screening or assaying bioengineered scaffolds or the mechanism of action of drugs and bioactive agents that modify axon growth and regeneration.

References

1. Allodi I, Guzmán-Lenis MS, Hernández J, Navarro X, Udina E. In vitro comparison of motor and sensory neuron outgrowth in a 3D collagen matrix. *J Neurosci Methods* 2011; 198: 53-61.
2. Bozkurt A, Brook GA, Moellers S, Lassner F, Sellhaus B, Weis J, Woeltje M, Tank J, Beckmann T, Fuchs P, Olde Damink L, Schügner F, Heschel I, Pallua N. In-vitro assessment of axonal growth using dorsal root ganglia (DRG) explants in a novel 3D collagen matrix. *Tissue Engineering* 2007; 13: 2971-2979.
3. Bozkurt A, Deumens R, Beckmann C, Olde Damink L, Schügner F, Heschel I, Selhaus B, Weis J, Jähnen-Dechent W, Brook GA, Pallua N. In vitro cell alignment obtained with a Schwann cell enriched microstructured nerve guide with longitudinal guidance channels. *Biomaterials* 2009; 30: 169-179.
4. Gerardo-Nava J, Hodde D, Katona I, Bozkurt A, Grehl T, Steinbusch HW, Weis J, Brook GA. Spinal cord organotypic slice cultures for the study of regenerating motor axon interactions with 3D scaffolds. *Biomaterials* 2014; 35: 4288-4296.
5. Kim Y, Karthikeyan K, Chirvi S, Dave DP. Neuro-optical microfluidic platform to study injury and regeneration of single axons. *Lab Chip* 2009; 9: 2576-2581.
6. Peyrin JM, Deleglise B, Saias L, Vignes M, Gougis P, Magnifico S, Betuing S, Pietri M, Caboche J, Vanhoutte P, Viovy JL, Brugg B. Axon diodes for the reconstruction of oriented neuronal networks in microfluidic chambers. *Lab Chip* 2011; 11: 3663-3673.
7. Vyasa A, Li Z, Aspalter M, Feiner J, Hoke A, Zhou C, O'Daly A, Abdullah M, Rohde C, Brushart TM. An in vitro model of adult mammalian nerve repair. *Exp Neurol* 2010; 223: 112-118.

Use of induced pluripotent stem cells – derived neurons for disease modeling

Liszewska E.^{1*}, Rydz K.¹, Kotulska K.², Jozwiak S.², Jaworski J.^{1*}

¹International Institute of Molecular and Cell Biology, Warsaw, Poland

²The Children's Memorial Health Institute, Warsaw, Poland

*Correspondence should be addressed to eliszewska@iimcb.gov.pl, jaworski@iimcb.gov.pl

Yamanaka's breakthrough discovery of cell reprogramming, awarded with the Nobel Prize in Physiology or Medicine in 2012, showed that the fate of differentiated somatic cells is not definitively defined and can be changed [14,15]. Yamanaka's strategy relied on over expression of transcrip-

tion factors required for pluripotency such as Oct4 (octamer-binding transcription factor 4), Klf4 (Krueppel-like factor 4), Sox2 (sex determining region Y-box 2), and c-Myc [14,15] in differentiated cells (e.g., fibroblasts). Cells modified in that way change their morphology to embryonic stem cell-like one, start expressing genes, normally only active in cells of the embryonic node, in the early stages of embryonic development and can be differentiated in all cell types characteristic of the three germ layers, including those that morphology and profile of protein produced resembled neurons. Therefore they were called induced pluripotent stem cells (iPSC).

While virus-based reprogramming technology is currently one of several options [1,17], it is still the most popular one and gives researchers an easy access to human cells that were previously unavailable or difficult to achieve, such as neural stem cells or different types of neurons [3,5,7,8,13]. In addition to reprogramming followed by differentiation, recently a new technology of direct change, both *in vitro* and *in vivo*, of differentiated cells (fibroblasts, astrocytes) to neurons (transdifferentiation) emerged [12,16]. An immediate consequence of these technologies was development of a number of cellular models of the nervous system diseases. These models are used both to explore the cellular mechanisms of these diseases and for the development of new pharmacological strategies as well as nuclease based correction of genetic defects.

Thus far, induced pluripotent stem cell-derived neurons (iPSC-N) and induced neurons (IN) have been already obtained from patients with several neurological disorders including neurodegenerative diseases (e.g., amyotrophic lateral sclerosis [ALS], Parkinson's [PD], Alzheimer's [AD] and Huntington's disease [HD]), neurodevelopmental diseases (e.g., fragile X syndrome, Rett syndrome) and psychiatric disorders (e.g., schizophrenia) [3,5,7,8,13]. In majority of cases researchers' efforts focused on reproducing basic cellular features of selected diseases. Yet, in few cases new interesting observations concerning disease mechanism have been made and schizophrenia, ALS and PD can serve as good examples. Brennand *et al.* [2] using schizophrenic patients' iPSC-N revealed disturbed dendritic arbor morphology and decreased synaptic connectivity. In iPSC-N derived from ALS patients carrying mutations in *TDP-43* (*TAR DNA-binding protein 43*), Egawa *et al.* [6] discovered that changes in RNA metabolism likely underlay sensitivity of these cells to oxidative stress. Last but not least, Belmonte group discovered new feature of PD caused by *LRRK2* (*leucine-rich repeat kinase 2*) mutations, namely death of neuroprecursors, which may explain cognitive dysfunctions observed in course of PD [10]. It is worth stressing that most ambitious projects assume the use of iPSC-N to search for new drugs. Although this type of exploration requires testing hundreds of thousands of substances, pre-

viously published work focused on testing only a few substances at the same time [2,6,11]. Yet, it should be stressed that some of these studies brought interesting results, e.g. use of anacardic acid for protecting ALS-derived iPSC-N against arsenate induced cell death [6].

Among the diseases that have not been yet modeled with use of iPSC-N is Tuberous sclerosis (TSC). TSC is a developmental disease caused by mutations in genes encoding either hamartin or tuberin and manifested by tumor development, epilepsy and autism [4]. On the cellular level the major characteristic of this disease are hypertrophy and mTOR signaling hyperactivation [9]. While the hyperactivation of the mTOR pathway is considered as one of the major causes of the pathology it is not clear what molecular mechanisms downstream of this multi-task homeostatic kinase are involved in TSC development. Using overexpression of Oct4, Klf4, Sox2 and c-Myc we established several iPSC lines from patients with mutations in *TSC1*, which were differentiated to iPSC-N (FENS Abstract, Volume 6, p093.06, 2012). The TSC patient-derived iPSC-N are characterized by hypertrophy, disturbed overall morphology and increased mTOR activity (e.g., increased phosphorylation of ribosomal protein S6). Thus, we show that these iPSC-N recapitulate major cellular defects observed in TSC patients and therefore can be used for further search for molecular mechanisms contributing to TSC development.

This work has been financed by ERA-NET NEURON/06/2011 “AMRePACELL” (co-financed by NCBiR) to JJ and Homing Plus programme (HOMING PLUS/2012-5/6) of the Foundation for Polish Science, cofinanced from the European Union under the European Regional Development Fund to EL.

References

- Bayart E, Cohen-Haguenauer O. Technological overview of iPSC induction from human adult somatic cells. *Curr Gene Ther* 2013; 13: 73-92.
- Brennand KJ, Simone A, Jou J, Gelboin-Burkhardt C, Tran N, Sangar S, Li Y, Mu Y, Chen G, Yu D, McCarthy S, Sebat J, Gage FH. Modelling schizophrenia using human induced pluripotent stem cells. *Nature* 2011; 473: 221-225.
- Chailangkarn T, Acab A, Muotri AR. Modeling neurodevelopmental disorders using human neurons. *Curr Opin Neurobiol* 2012; 22: 785-790.
- Curatolo P, Bombardieri R, Jozwiak S. Tuberous sclerosis. *Lancet* 2008; 372: 657-668.
- Dolmetsch R, Geschwind DH. The human brain in a dish: the promise of iPSC-derived neurons. *Cell* 2011; 145: 831-834.
- Egawa N, Kitaoka S, Tsukita K, Naitoh M, Takahashi K, Yamamoto T, Adachi F, Kondo T, Okita K, Asaka I, Aoi T, Watanabe A, Yamada Y, Morizane A, Takahashi J, Ayaki T, Ito H, Yoshikawa K, Yamawaki S, Suzuki S, Watanabe D, Hioki H, Kaneko T, Makioka K, Okamoto K, Takuma H, Tamaoka A, Hasegawa K, Nonaka T, Hasegawa M, Kawata A, Yoshida M, Nakahata T, Takahashi R, Marchetto MC, Gage FH, Yamanaka S, Inoue H. Drug screening for ALS using patient-specific induced pluripotent stem cells. *Sci Transl Med* 2012; 4: 145ra104.
- Han SS, Williams LA, Eggan KC. Constructing and deconstructing stem cell models of neurological disease. *Neuron* 2011; 70: 626-644.
- Koch P, Kokaia Z, Lindvall O, Brustle O. Emerging concepts in neural stem cell research: autologous repair and cell-based disease modelling. *Lancet Neurol* 2009; 8: 819-829.
- Kwiatkowski DJ. Tuberous sclerosis: from tubers to mTOR. *Ann Hum Genet* 2003; 67: 87-96.
- Liu GH, Qu J, Suzuki K, Nivet E, Li M, Montserrat N, Yi F, Xu X, Ruiz S, Zhang W, Wagner U, Kim A, Ren B, Li Y, Goebel A, Kim J, Soligalla RD, Dubova I, Thompson J, Yates J, 3rd, Esteban CR, Sancho-Martinez I, Izpisua Belmonte JC. Progressive degeneration of human neural stem cells caused by pathogenic LRRK2. *Nature* 2012; 491: 603-607.
- Marchetto MC, Carron C, Acab A, Yu D, Yeo GW, Mu Y, Chen G, Gage FH, Muotri AR. A model for neural development and treatment of Rett syndrome using human induced pluripotent stem cells. *Cell* 2010; 143: 527-539.
- Niu W, Zang T, Zou Y, Fang S, Smith DK, Bachoo R, Zhang CL. In vivo reprogramming of astrocytes to neuroblasts in the adult brain. *Nat Cell Biol* 2013; 15: 1164-1175.
- Robinton DA, Daley GQ. The promise of induced pluripotent stem cells in research and therapy. *Nature* 2012; 481: 295-305.
- Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007; 131: 861-872.
- Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006; 126: 663-676.
- Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Sudhof TC, Wernig M. Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* 2010; 463: 1035-1041.
- Zhou YY, Zeng F. Integration-free methods for generating induced pluripotent stem cells. *Genomics Proteomics Bioinformatics* 2013; 11: 284-287.

Biogenesis, transport and degradation of mitochondrial proteins

Chacinska A.

International Institute of Molecular and Cell Biology, Warsaw, Poland

Mitochondria play an important role in variety of metabolic and regulatory processes and their dysfunction leads to life threatening disorders. About 1000-1500 proteins are needed to build this essential organelle. The large majority of mitochondrial proteins is synthesized on the cytosolic ribosomes, and after completion of their synthesis must

be properly targeted to mitochondria. After synthesis, precursors of mitochondrial proteins are directed to the TOM complex and subsequently to specific mitochondrial import machineries that are responsible for their sorting. Proteins destined to the intermembrane space of mitochondria follow the mitochondrial intermembrane space import and assembly (MIA) pathway. The MIA pathway provides an efficient and redox-dependent mechanism for trapping substrate proteins in the intermembrane space of mitochondria. The cytosolic steps that precede mitochondrial import have not been well understood. We identified a role for the ubiquitin-proteasome system in the biogenesis of the intermembrane space proteins. The ubiquitin-proteasome system persistently removes a fraction of intermembrane space proteins under physiological conditions, acting as a negative regulator in the biogenesis of this class of proteins. Moreover, based on unbiased proteomic approaches, we discovered that mitochondria play a role in the regulation of cellular protein homeostasis.

Mechanisms of neuroprotection by ischemic preconditioning: synaptic and mitochondrial modifications

Perez-Pinzon M.

Department of Neurology, University of Miami, Miller School of Medicine, Miami, USA

Ischemic Preconditioning (IPC) has been shown to be a robust neuroprotective approach to protect against cerebral ischemia. IPC is modelled when tissues are exposed to mild ischemic insults and after a period of recovery they become highly resistant to a subsequent, more lethal ischemic insult. My laboratory has demonstrated that the ischemic tolerance that ensues following IPC occurs at the level of mitochondria and by modifications in synaptic function. Mitochondria are key players in ischemic/reperfusion injury hours to days following cerebral ischemia and thus an important target for IPC-induced neuroprotection. Excitotoxicity following cerebral ischemia plays a major role in the pathology that ensues following cerebral ischemia. This presentation will summarize several signaling pathways involved in mitochondrial and synaptic neuroprotection against ischemia following IPC.

These studies were supported by National Institutes of Health Grants NS45676-01, NS054147-01 and NS34773.

Two faces of ER stress-from cancer to neurodegenerative diseases

Piwocka K.¹, Kusio-Kobialka M.¹, Dudka-Ruszkowska W.¹, Brewinska-Olchowik M.¹, Krysa W.², Rakowicz M.², Stepniak I.², Sulek A.²

¹Nencki Institute of Experimental Biology, Warsaw, Poland

²Institute of Psychiatry and Neurology, Warsaw, Poland

Endoplasmic reticulum (ER) is the site where proteins folding and assembly occurs. A number of stimuli, such as perturbations in calcium homeostasis or redox state, accumulation of misfolded proteins, glucose deprivation, and altered glycosylation disrupts ER homeostasis and induces ER stress. Mild chronic ER stress allows for adaptation and leads to permanent changes in cellular function and activation of pro-survival mechanisms, whereas excessive stress activates cell death execution. The balance between adaptation *versus* cell death has a tremendous physiological importance. The UPR (Unfolded Protein Response) activates three signaling branches participating in regulation between survival and apoptosis upon ER stress. Activation of the ER stress signalling can play an opposite role in cancer and neurodegenerative diseases. In cancer, activation of the UPR response plays predominantly a prosurvival and protective role, but can also correlate with induction of cell death. Previously, the cytoprotective function was mostly correlated with solid tumors. However, we showed that mild ER stress and one of the UPR branches are activated in chronic myeloid leukemia cells, and are connected with the disease progression and resistance to imatinib therapy. Attenuation of this pathway in CML cells led to inhibition of proleukemic signaling, apoptosis and increased sensitivity to imatinib. On the other hand, disturbed levels of endoplasmic reticulum stress (ER) markers were already observed in post-mortem brains, mouse models and cell cultures derived from patients with Alzheimer's (AD), Parkinson's or Huntington's disease (HD), highlighting the role of ER stress in the neurodegeneration. Moreover, targeting the UPR elements seems to be an interesting strategy to prevent loss of neurons. It was also reported that UPR proteins are activated in blood cells of AD patients, suggesting that peripheral cells may serve as a model to study the disease pathophysiology. In HD mutated huntingtin aggregates provoke ER stress and the UPR. We performed pilot studies to verify the hypothesis that level of some ER stress mediators may serve as a peripheral biomarker in HD patients, correlating to disease stage and number of CAG repetitions in carriers of CAG expansion in *HTT* locus. We assume that pathophysiological processes appear in HD prior to clinical manifestations, and this study was done to look for the biomarkers correlating to the disease

progression and indicators of the compensative mechanisms efficiency in asymptomatic subjects. Understanding the complexity of ER stress response in cancer and neurodegeneration is necessary and may help to improve current, and discover novel therapeutic strategies.

This work was supported by grants 2011/01/B/NZ3/02145 and 2013/10/E/NZ3/00673 from the National Science Centre in Poland.

Astrocytes ER Ca²⁺ signalling and its pathological relevance

Verkhatsky A.

The University of Manchester, Manchester, United Kingdom

The evolution of the central nervous system (CNS) resulted in an appearance of highly specialised neuronal networks optimised for rapid information transfer. In the course of this specialisation neuronal cells lost their metabolic independence and the ability to survive in the absence of homeostatic systems. These homeostatic systems, represented by neuroglia regulate all aspects of CNS function in physiological and pathological conditions. The neurological diseases should be therefore considered as primary gliopathologies, which determine the progression and outcome of neuropathological process. Glial function is intimately regulated by cellular calcium signalling that underlies the specific form of “glial calcium excitability”. Glial Ca²⁺ signals are triggered by activation of multiple receptors, and are primarily driven by Ca²⁺ release from the endoplasmic reticulum. In this review we summarise the role of glial calcium signalling in various forms of pathological processes including neurological and psychiatric disorders and neurodegeneration.

Who is running your gut? The gut-brain axis in brain metastasis formation by polychlorinated biphenyls

Toborek M.

Department of Biochemistry and Molecular Biology, University of Miami School of Medicine, USA

The gut microbiome is a dynamic bacterial community that interacts with the host and closely relates to human health by regulating energy metabolism and immune functions. Each host has a unique composition of gut microbi-

ota, which implies highly individual responses to environmental stressors and suggests a role for gut microbiota in future personalized health strategies. However, the influence of the microbiome on the toxicity of environmental pollutants and its role in risk assessment are largely unknown. In addition, there is an emerging interest in the role of behavioral factors in modulating toxicity of environmental pollutants. While the role of nutrition has been explored, the impact of exercise on the health effects of toxicants is not known. Because exercise can influence the outcomes of disorders known to be associated with alterations of the gut microbiome, we hypothesized that physical activity may affect the composition of the gut microbiota and thus influence the impact of environmental toxicants. In order to address this hypothesis, we investigated the effects of polychlorinated biphenyls (PCBs) and exercise on the composition and structure of the gut microbiome. In addition, we related these results to PCB-induced development of brain metastases [1,2]. Mice exercised voluntarily for 5 weeks, followed by the exposure to a mixture of environmentally relevant PCB congeners (PCB153, PCB138 and PCB180) for 48 h. The microbiome was assessed by determination of 16S rRNA. Oral exposure to PCBs disrupted the integrity of the gut barrier function and significantly altered the abundance of the gut microbiome primarily by decreasing the levels of Proteobacteria. The activity level correlated with a substantial shift in abundance, biodiversity, and microbiome composition. Importantly, exercise attenuated PCB-induced changes in the gut microbiome. Oral exposure to PCBs facilitated the development of brain metastases, the effects that was attenuated by exercise. This study provides the first evidence that oral exposure to PCBs can induce substantial changes in the gut microbiome, which may then influence their systemic toxicity. Importantly, these changes can be attenuated by behavioral factors, such as voluntary exercise.

References

1. Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. Exercise attenuates PCB-induced changes in the mouse gut microbiome. *Environ Health Perspect* 2003; 121: 725-730.
2. Sipos E, Chen L, Andras IE, Wrobel J, Zhang B, Pu H, Park MS, Eum SY, Toborek M. Proinflammatory adhesion molecules facilitate polychlorinated biphenyl-mediated enhancement of brain metastasis formation. *Toxicol Sci* 2012; 126: 362-371.

Uptake and toxicity of metal containing nanoparticles in brain cells

Dringen R.

Centre for Biomolecular Interactions Bremen, University of Bremen, PO. Box 330440, D-28334 Bremen, Germany
Centre for Environmental Research and Sustainable Technology, University of Bremen, Leobener Straße, D-28359 Bremen, Germany

Metal-containing nanoparticles (NPs) have gained considerable interest during the last decade due to their exciting properties as antibacterial agents, for consumer products as well as for technical applications and diagnostics. Due to their frequent use, the likelihood that brain cells will encounter such NPs has strongly increased. To investigate potential adverse effects of NPs on brain cells, we have studied the biocompatibility, the uptake and the metabolism of metal-containing NPs such as iron oxide NPs (IONPs), copper oxide NPs (CuONPs) and silver NPs (AgNPs) on cultured primary brain cells. Cultured astrocytes, neurons and microglial cells accumulated IONPs efficiently in a time-, concentration- and temperature-dependent manner. However, despite of an accumulation of large amounts of iron in IONP-treated astrocytes and neurons, the viability of these brain cells was not compromised. In contrast, accumulated IONPs were toxic for microglial cells, most likely due to the rapid uptake and transfer of the IONPs into the lysosomal compartment which facilitates the liberation of iron ions from the accumulated particles. The presence of fetal calf serum in the incubation medium strongly decreased the rate of IONP uptake into brain cells. While some inhibitors of endocytotic pathways lowered IONP accumulation in serum-containing medium, these inhibitors did not affect the rapid IONP accumulation in serum-free conditions. Astrocytes that had been exposed to IONPs for only 4 h remained viable for up to 7 days, hardly releasing any iron during this incubation period and showed a transient appearance of reactive oxygen species and a strong upregulation of the iron storage protein ferritin. Similarly, AgNP-treated astrocytes were not compromised in their viability and increased their content of metal-storing metallothioneins. These data demonstrate that cultured brain astrocytes deal well even with large amounts of iron or silver that are accumulated as IONPs or AgNPs. In contrast, the treatment of cultured astrocytes with CuONPs caused severe oxidative stress-mediated toxicity that was prevented by copper chelators, suggesting that rapid liberation of copper from accumulated CuONPs is involved in the observed toxicity of these NPs to astrocytes. These data demonstrate that different brain cell types differ in their vulnerability to one type of metal-

containing NPs and that NP toxicity on a given type of brain cells strongly depends on the content of applied NPs.

Impairment of glutamine/glutamate- γ -aminobutyric acid cycle in manganese toxicity in the central nervous system

Sidoryk-Wegrzynowicz M.

Cambridge Centre of Brain Repair, University of Cambridge, Cambridge, United Kingdom

Manganese (Mn) is an essential trace element that is required for maintaining the proper function and regulation of many biochemical and cellular reactions. Despite its essentiality, at excessive levels Mn is toxic to the central nervous system (CNS). The overdose accumulation of Mn in specific brain areas, such as the substantia nigra, the globus pallidus and the striatum, triggers neurotoxicity resulting in a neurological brain disorder, referred to as manganism. Mn toxicity is associated with the disruption of glutamine (Gln)/glutamate (Glu) GABA cycle (GGC). The GGC represents a complex process, since Gln efflux from astrocytes must be met by its influx in neurons. Mn toxicity is associated with the disruption of both of these critical points in the cycle. In cultured astrocytes, pre-treatment with Mn inhibits the initial net uptake of Gln in a concentration-dependent manner. Mn added directly to primary culture of astrocytes induces deregulation in the expression of SNAT3, SNAT2, ASCT2 and LAT2 transporters and significantly decreases in Gln uptake mediated by the transporting Systems N and ASC, and a decrease in Gln efflux mediated by Systems N, ASC and L. Further, Mn disrupts Glu transporting systems leading to both a reduction in Glu uptake and elevation in extracellular Glu levels. Interestingly, there appear to be common signaling targets of Mn in GGC cycling in glial cells. Namely, the PKC signaling is affected by Mn in glutamine and glutamate transporters expression and function. Additionally, Mn deregulates GS expression and activity. Those evidences could triggers depletion of glutamine synthesis/metabolism in glia cells and consequently diminish astrocytic-derived glutamine, while disruption of Glu removal/transport can mediate dyshomeostasis in neurotransmission of functioning neurons. Here we highlight the mechanistic commonalities inherent to Mn-neurotoxicity related to the astrocyte pathology and GGC impairment.

Cadmium as a pro-inflammatory factor. Chosen cadmium neurotoxicity issues

Olszowski T.

Department of Hygiene and Epidemiology, Pomeranian Medical University, Szczecin, Poland

Cadmium belongs to the environmental toxins that constitute nowadays a serious problem for public health. The aim of this paper is to summarize the existing state of knowledge on the effects of cadmium on selected inflammatory mediators and markers, such as NF- κ B, AP-1, IL-6, TNF- α , IL-1 β , IL-8, MPO, iNOS, MMPs, COX-2, PGE₂, ICAM-1, VCAM-1, PECAM-1 and CRP in humans and rodents, both *in vitro* and *in vivo*. Moreover, the results of the author's original research on the effects of cadmium on COX-1 and COX-2 gene, protein expression and enzymatic activity in THP-1 macrophages are presented. In addition, the literature review as regards the neurotoxic effects of cadmium in humans and rodents is discussed. The effects on oxidative stress, interference with calcium, interaction with other metals, the impact on neurotransmitters, apoptosis induction, estrogen-like effects and epigenetic modification may be the mechanisms underlying cadmium neurotoxicity.

New mechanisms of GPCRs function: regulation by protein-protein interaction

Georgoussi Z.

Institute of Biosciences and Applications, Laboratory of Cellular Signaling and Molecular Pharmacology, National Centre for Scientific Research “Demokritos”, 15310 Ag. Paraskevi, Athens, Greece

Evidence has demonstrated that G protein coupled receptors (GPCRs) can physically interact with a variety of accessory proteins, confirming that signal transduction of these receptors is not restricted to heterotrimeric G protein activation. Such interactions can alter the effectiveness of agonist-driven cell signalling, determine the signals generated and alter the trafficking, targeting, fine tuning and cellular localization of these receptors by providing a scaffold that links the receptors to the cytoskeletal network (Georgoussi *et al.* 2013). Opioid receptors which belong to GPCRs modulate a variety of physiological responses in the nervous system through activation of a diverse array of effector systems ranging from adenylyl cyclase, ion channels to other signalling intermediates. Opioid administration causes activation of several transcription factors, including CREB, NF- κ B, STAT3 and STAT5A/B (Tso and

Wong 2003; Mazarakou and Georgoussi 2005; Georganta *et al.* 2010). Such parallel manifestations of opioid receptor function suggest that these receptors are involved in different signaling circuits that lead to alterations in target gene expression in a pleiotropic fashion.

In the present study emphasis will be given to unconventional interacting partners of the μ - and δ -opioid receptors opioid receptors such as the regulators of G protein signalling (RGS) proteins and STAT5B. Evidence will be presented on which RGS proteins opioid receptors interact; how RGS4 and RGS2 confer selectivity to the receptors to choose a specific subset of G proteins; how activation of opioid receptors result in recruitment of RGS proteins to the plasma membrane; and how RGS proteins exert a differential modulatory effect in ERK1,2 phosphorylation, agonist-driven inhibition of adenylyl cyclase and the internalization fate of opioid receptors. Moreover evidence will be presented on how STAT5B associates with the δ -opioid receptor and forms selective pairs with selective G α and G $\beta\gamma$ subunits and RGS proteins; and how activation of the δ -opioid receptor with selective agonists promotes a multi-component signaling complex involving the STAT5B transcription factor and other signaling intermediates to mediate neuronal survival and neurite outgrowth. Understanding the mechanism of intercellular communication via G-protein-mediated signalling pathways in the nervous system is crucial for normal brain development and regulation of adult neural processes. Thus, defining the molecular determinants that control opioid receptor signaling is important to address problems related to phenomena such as pain perception, tolerance and dependence that occur upon chronic opiate administration and define whether disruption of such interactions may contribute to the pathophysiology of nervous system related disorders and to the development of novel therapeutic strategies.

This work was supported by the EU «Normolife» (LSHC-CT2006-037733) and the GSRT, Excellence II grant 3722, “NO-ALGOS”. Z.G participates in the European COST Action CM1207 (GLISTEN).

The role of opioid peptides under neuropathic pain

Mika J.

Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Despite many advances, our understanding of the involvement of endogenous opioid peptides in the develop-

ment of neuropathic pain, is not fully understood. Many recent studies show, that microglia cells play a crucial role in the maintenance of neuronal homeostasis, therefore pain may now be considered as neuro-immune disorder. The microglial production of immune factors is believed to play an important role in nociceptive transmission and opioids effectiveness. Endogenous opioidergic neurotransmission is found throughout the brain and spinal cord and appears to influence many nervous system functions, including nociception. Paradoxaly opioids can induce both, analgesia and algesia. It has recently been hypothesized that the imbalance between the pro- and anti-nociceptive activity of peptides generated from opioid prohormones is leading to the development of chronic pain. Our studies show that minocycline reduced prodynorphin mRNA levels that were previously elevated in the spinal cord and/or dorsal root ganglia following sciatic nerve injury. Moreover, the repeated administration of minocycline enhanced the analgesic effects of low-dose dynorphin (0.15 nmol) and prevented the development of flaccid paralysis following high-dose dynorphin administration (15 nmol), suggesting a neuroprotective effect. Minocycline diminished the dynorphin(15 nmol)-induced upregulation of expression of IL-1 β and IL-6 within the spinal cord. These results suggest an important role of these proinflammatory cytokines in the development of the neurotoxic effects of dynorphin. Additionally, our study of primary microglial cell culture confirmed the presence of μ - and κ -opioid receptors. Further, we provide the first evidence for the lack of δ -opioid receptors on microglial cells. Interestingly, the analgesic effect of δ -opioid receptor ligands (DPDPE, deltorphine II) in neuropathic pain is not diminished in contrast to other opioid receptor ligands (morphine, DAMGO, U50.488H, SNC80). In conclusion, our results underline the importance of neuro-glial interactions as evidenced by the involvement of IL-1 β and IL-6 and the minocycline effect in dynorphin-induced toxicity, which suggests that drugs, that alter the prodynorphin system could be used to a better control of neuropathic pain. Our results suggest, that δ -opioid receptor agonists analgesia is different from analgesia induced by μ - and κ -opioid receptors because it does not depend on injury-induced microglial activation. Therefore, δ -opioid receptor agonists appear to be the best candidates for new drugs to treat neuropathic pain.

NCN grants 2011/03/B/NZ4/00042 and 2012/06/A/NZ4/00028.

Peptides in neurological disorders

Nyberg F.

Department of Pharmaceutical Biosciences, Uppsala University, Sweden

This presentation addresses past and current research on various neuropeptides that have been implicated in neurological complications, including drug addiction, chronic pain, cognitive disorders, anxiety, and brain injury. However, considering the large and even increasing number of identified neuroactive peptides, it is necessary to limit this presentation to a few peptides or peptide systems, which have received particular attention in relation to these items. Among these are the opioid peptides, the tachykinins, calcitonin gene-related peptide (CGRP), and peptides belonging to the endocrine system. Most of these neuropeptides are not only involved in pain processing and drug addiction but are also involved in the endogenous mechanisms of regulating memory and cognitive function. Also many of these neuroactive peptides are affected and dysregulated at injury to the central nervous system (CNS). The presentation also highlights some new aspects regarding the possibility to use endocrine peptides to counteract and repair brain damages induced by addictive drugs and CNS injuries. It does not endeavor to fully cover the field but it does aim to give an idea of how various neuropeptides may be involved in the control of various neurobehavior and how peptidergic systems are affected during various diseases related to the CNS.

Peptides in angiogenesis. Development of peptidomimetic inhibitors of VEGF co-receptor NRP-1

Tymecka D.¹, Fedorczyk F.¹, Niescioruk A.¹, Grabowska K.¹, Puszko A.¹, Sosnowski P.¹, Wilenska B.¹, Witkowska E.¹, Perret G.Y.³, Starzec A.^{1,3}, Misicka A.¹

¹Faculty of Chemistry, Biological and Chemical Research Center, University of Warsaw, Warsaw, Poland

²Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

³Universite Paris 13, Li2P EA 4222, Bobigny, France

Angiogenesis, the growth of new capillary blood vessels from existing ones, is an important physiological process in embryogenesis, wound healing and reproduction. In cancer, angiogenesis permits to develop a network of intra tumour blood vessels supplying oxygen and nutrients thus enhancing tumour growth. Therefore, compounds blocking

angiogenesis could be prospective antitumour drugs, but so far only a few angiogenesis inhibitors are used in the clinic for the treatment of cancer.

One of the most important growth factors involved in angiogenesis process is vascular endothelial growth factor-165 (VEGF₁₆₅). VEGF₁₆₅ signal is transduced via VEGF receptors and significantly enhanced by association with co-receptor NRP-1. Moreover, overexpression of NRP-1 was found in different malignant cancer tissues. All these findings suggest that NRP-1 could be involved in pathological angiogenesis and inhibitors of NRP-1 have considerable potential as novel antiangiogenic and cancer therapeutics.

As a starting point for our study of searching for anti-angiogenic peptidomimetics we utilized a heptapeptide ATWLPPR (A7R). It was shown that A7R binds to NRP-1 and selectively inhibits VEGF₁₆₅ binding to NRP-1, and *in vivo* treatment with A7R resulted in decreasing breast cancer angiogenesis and growth [3]. The detailed studies have shown the importance of C-terminal arginine and that its shortest sequence preserving inhibitory activity is the C-terminal tetrapeptide LPPR [2].

This presentation reports structure-activity relationship studies of the tetrapeptide LPPR in order to understand the structural determinants for inhibition of binding VEGF₁₆₅ to NRP-1 by this tetrapeptide and to develop more resistant to enzymatic degradation peptidomimetic NRP-1 inhibitors. These compounds may serve as tools for the study of NRP-1 biology and provide a point of departure for development of more potent anti-NRP-1 drugs in the future [1].

This work was supported by NCN grant no N 204 350940. Project was carried out with the use of CePT infrastructure cofinanced by the European Union – the European Regional Development Fund within the Operational Programme “Innovative economy” for 2007-2013.

References

1. Misicka-Kesik A, Tymecka D, Fedorczyk B, Wilenska B, Sosnowski P, Witkowska E, Ladam P, Perret G, Starzec A. Polish Patent Application, P.405129.
2. Starzec A, Ladam P, Vassy R, Badache S, Buchemal N, Navaza A, Herve du Penhoat C, Perret GY. Structure-function analysis of the antiangiogenic ATWLPPR peptide inhibiting VEGF(165) binding to neuropilin-1 and molecular dynamics simulations of the ATWLPPR/neuropilin-1 complex. *Peptides* 2007; 28: 2397-2402.
3. Starzec A, Vassy R, Martin A, Lecouvey M, Di Benedetto M, Crepin M, Perret GY. Antiangiogenic and antitumor activities of peptide inhibiting the vascular endothelial growth factor binding to neuropilin-1. *Life Sci* 2006; 79: 2370-2381.

Stroke-induced neurogenesis

Kokaia Z.

Stem Cell Center, Lund University, Sweden

Stroke is caused by occlusion of a cerebral artery, which gives rise to focal ischemia with irreversible injury in a core region and partially reversible damage in the surrounding penumbra zone.

Stroke is the leading cause of disability in adult humans in developed countries. After ischemic stroke neurons are rapidly damaged and usually die but cellular loss can occur hours and days thereafter. Stroke increases stem cell proliferation in the subventricular zone (SVZ) and the generated neuroblasts migrate to the stroke-damaged area of the brain where they become mature neurons. This process may continue for several months and produce a significant number of new, functional neurons. The evidence of increased neuroblast production after stroke has also been demonstrated in human brain. A bulk of experimental evidence supports the idea that the stroke-damaged adult brain makes an attempt to repair itself by producing new neurons also in areas where neurogenesis does not normally occur, e.g., striatum and cerebral cortex. Knowledge about mechanisms regulating the different steps of neurogenesis after stroke is rapidly increasing but still incomplete. The functional consequences of stroke-induced neurogenesis and the level of integration of the new neurons into existing neural circuitries are poorly understood. In order to have a substantial impact on the recovery after stroke, this potential mechanism for self-repair needs to be markedly enhanced, primarily by increasing the survival and differentiation of the generated neuroblasts. Moreover, for efficient repair, optimization of neurogenesis most likely needs to be combined with promotion of other endogenous neuroregenerative responses, e.g., protection and sprouting of remaining mature neurons, and transplantation of stem cell-derived neurons and glia cells.

Injury-induced neurogenesis in the adult striatum

**Magnusson J.P.¹, Göritz C.¹, Tatarishvili J.²,
Smith E.M.K.³, Kokaia Z.², Lindvall O.², Frisén J.¹**

¹Department of Cell and Molecular Biology, Karolinska Institute, SE-171 77, Stockholm, Sweden

²Wallenberg Neuroscience Center, Lund Strategic Research Center for Stem Cell Biology and Cell Therapy, Lund University Hospital, SE-221 84 Lund, Sweden

³Division of Translational Cancer Research, Lund University, SE-223 63 Lund, Sweden

The adult brain is not good at repairing itself. After brain injury, functional recovery is generally restricted to behavioral compensation and rewiring of spared circuits and there is very limited, if any, replacement of dead neurons. In the adult mammalian brain, neuron-producing stem cells do exist in two spatially restricted regions, the dentate gyrus of the hippocampus and the subventricular zone lining the lateral ventricles. These stem cells are important for normal brain function and their primary role does not seem to be to repair injuries; however, the discovery of these cells has garnered much speculation about whether the brain's inherent capacity to make new neurons can be harnessed to improve functional recovery after injury. It is likely that a detailed understanding of the mechanisms that govern adult neurogenesis will be crucial if this prospect is to become reality. The striatum has recently emerged as a brain region that sustains neurogenesis into adulthood in humans; in addition, this region is known to support injury-induced neurogenesis in animal models of stroke. In my talk, I will discuss recent findings regarding the extent, origin and molecular mechanisms governing stroke-induced neurogenesis in the striatum – results that highlight a previously unknown level of plasticity in the adult brain.

Poster session I

[A1]

Intervertebral disc regeneration with the use of autologous population of chondrocytes and its analysis in terms of morphology and activities

Abaaslam F.¹, Wluka A.², Głowacki M.¹, Czernicki Z.¹, Lipkowski A.W.²

¹Department of Neurosurgery, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Department of Neuropeptides, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Introduction: Lumbosacral spine pains affect nearly 80% of the population. In the last two decades various biological methods aimed at treatment of the intervertebral disc diseases have been developed. Out of all methods, the cell therapy based on the transplantation of autologous chondrocyte cells seems promising. This procedure stimulates the matrix production, increases the population of chondrocytes and equates the disturbed balance within the disc.

The aim of the study was to establish the best method of obtaining chondrocytes for autologous cell transplantations in cases of intervertebral disc diseases. This project takes into account the duration of the disease, the place of collecting the material from the disc, the quantity of the collected material and the degree of destruction of the intervertebral disc, in order to derive the optimal, autologous material to be administered to the intervertebral space during operation.

Material and methods: The population of the study consisted of 47 patients whose age ranged from 18 to 40. Those patients were diagnosed with a single-level lumbar degenerative disc disorder on the basis of clinical and MRI examinations. The research topic of the present study dealt with the chondrocyte cells culture derived from the intervertebral disc by means of a single-level discectomy at L4-S1 levels. The analysis also included the target material removed from the intervertebral disc.

Results: The study revealed that the transplantation of cultured cells produced a good radiological and clinical effect. Tests in confocal microscope showed good integration of transplanted cells with the matrix and their correct activity. It was observed that the relation between the production of typical components of the matrix-collagen I, II, III, IV, proteoglycans, aggrecans and anomalous for the matrix-collagen type IV in the initial material was appropriate.

Conclusions: The conducted examinations show that the transplantation of chondrocyte cells derived from the culture to the previously operated intervertebral space suppresses degenerative processes of the disc and influences its recreation. Tests in the light and confocal microscope indicate a good integration of transplanted cells with the matrix and their correct productive activity as the production of typical components of the matrix-collagen I, II, III, IX, proteoglycans and aggrecans and anomalous for the matrix-collagen type I. The examination of the matrix-collagen type D could be the subject for future research. Such research might verify the role of this collagen in a healthy nucleus pulposus, and its functions in both processes of degeneration and regeneration of the intervertebral disc. To conclude, lack of inflammatory processes and other clinical complications after the transplantation of cells derived from the autologous chondrocytes culture show that this procedure is safe and beneficial for patients.

[A2]

Adhesion to the endothelial wall – the crucial step of cell homing into the tissue after their systemic transplantation

Andrzejewska A.¹, Nowakowski A.¹, Janowski M.^{1,2,3}, Koniusz S.¹, Walczak P.³, Grygorowicz T.⁴, Jablonska A.^{1,2}, Lukomska B.¹

¹NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Cellular Imaging Section and Vascular Biology Program, Institute for Cell Engineering, the Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

³Russel H. Morgan Vascular Biology Program, Division of MR Research, The Johns Hopkins University School of Medicine, Baltimore, USA

⁴Laboratory of Advanced Microscopy Techniques, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Application of mesenchymal stromal cells (MSCs) is a new and promising approach for the treatment of neurological disorders. The success of clinical application of MSCs relies upon the efficient recruitment of these cells into appropriate tissues. Systemic infusion of MSCs is preferred as a minimally invasive method of transplantation however cell migration from the vasculature into injured brain is inefficient. Leukocytes, circulating in the blood are known for their ability to migrate through the endothelial wall into surrounding tissues and this is a multi-step process that includes rolling and adhesion to the endothe-

lial cells. Initial capture of the leukocytes on the endothelium is promoted by the adhesion proteins i.e. selectins and integrins present on the leukocyte cell surface, interacting with their specific ligands expressed on activated endothelial cells. We hypothesize that the overexpression of $\alpha 4\beta 1$ (VLA-4) integrin in MSCs will enhance their migration towards ischemic brain regions after systemic transplantation.

Material and methods: Human bone marrow mesenchymal stromal cells (hBM-MSCs) (Lonza) were employed. Overexpression of $\alpha 4$ subunit of VLA-4 in hBM-MSCs was obtained by mRNA-ITGA4 based transfection. The naïve (non-transfected) and modified (mRNA-ITGA4 transfected) hBM-MSCs were labelled with iron nanoparticles coupled with Rhodamine-B (Molday, BioPAL) and used in microfluidic assays with microchannels coated with TNF- α activated rat brain endothelial cell line (RBE-4). The experiments with microfluidic device are based on the use of special microchannels, where laminar flow mimics physiological blood flow *in vivo*. Microchannels are covered with endothelial cells in the way that seeded cells resemble physiological blood vessel wall by their shape and diameter. The hBM-MSCs are forced to pass through microchannels with the flow and their interactions and adhesion rate to endothelial cells layer are assessed. The systemic transplantation of 10^5 naïve hBM-MSCs labelled with iron nanoparticles into internal carotid artery of ouabain-induced stroke rat was observed using MRI. The displaying was performed during the cells infusion and the day after surgery.

Results and conclusions: During the *in vitro* observation naïve hBM-MSCs interacted with rat endothelial cells. After pressure-driven perfusion of hBM-MSCs through the microfluidic channels even naïve cells were found to roll and be captured at the surface of endothelial cell layer. The effect of $\alpha 4\beta 1$ overexpression in MSCs on endothelial adhesion is currently being investigated. The model of microfluidic assay is a useful tool that can be used to assess the adherent properties of engineered hBM-MSCs before their intravenous or intra-arterial transplantation. The *in vivo* MRI research revealed a huge loss of transplanted hBM-MSCs the day after surgery in comparison to their amount observed directly after transplantation. This may be caused by their weak adhesiveness. We hope that our further study will show both *in vitro* and *in vivo* enhanced adherent properties of engineered hBM-MSCs.

Supported by a NCR&D grant No 101 in ERA-NET NEURON project: "MEMS-IRBI".

[A3]

Hypoxic ventilatory response after central dopamine receptor blockade in rat model of Parkinson's disease

Andrzejewski K.¹, Budzinska K.¹, Kaczynska K.²

¹Department of Respiratory Research, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Laboratory of Respiratory Reflexes, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Parkinson's disease (PD) patients apart from motor dysfunctions exhibit also respiratory disturbances such as: dyspnea, restrictive pulmonary function, respiratory dysrhythmias, upper airway dysfunction. Our study was aimed to investigate the effect of impairment of nigrostriatal dopaminergic system on resting breathing and hypoxic ventilatory response in experimental model of PD. Further, the involvement of central dopamine D₂ receptors in the hypoxic ventilatory response was examined. In our experiments we used neurotoxin 6-hydroxydopamine (6-OHDA) model with unilateral, double injection into the right medial forebrain bundle (MFB).

The experiments were performed on adult Wistar rats microinjected with 6-OHDA or vehicle according to the stereotaxic coordinates. Ventilatory parameters: tidal volume, minute ventilation, and frequency of breathing were measured in conscious rats with the use of whole body plethysmograph apparatus before and two weeks following the lesion. Changes in body weight and behavioral cylinder test were evaluated at the same time points and compared with the results obtained in sham-operated animals. Effects of centrally acting dopamine D₂ receptor antagonist, haloperidol (1 mg/kg *i.p.*), on ventilatory responses during resting breathing and exposure to hypoxia (8%) were studied before and after 6-OHDA injection.

Two weeks after 6-OHDA treatment the cylinder test displayed limb use asymmetry. Body weight showed lesser increase in comparison to the sham-operated animals. No effect of nigrostriatal pathway degeneration to the resting normoxic respiration was observed. Injection of 6-OHDA caused increase in tidal volume and decrease in respiratory rate during respiratory response to hypoxia with no alteration in minute ventilation. Haloperidol treatment in intact animals evoked significant increase in V_T stimulation in response to hypoxia. Fourteen days after lesion the effect of haloperidol injection was not observed.

In conclusion, stimulatory action of haloperidol via D₂ central dopamine receptor on tidal volume in reaction to hypoxia suggests inhibitory involvement of central dopamine to hypoxic ventilatory response. Augmentation of tid-

al volume response to hypoxia present after lesion in the MFB area, resulted from a deficit of dopamine connected with degeneration of dopaminergic neurons evoked by 6-OHDA treatment.

[A4]

The influence of dopaminergic system lesion and repeated administration of pramipexole and imipramine on BDNF and trkB mRNAs expression

Berghauzen-Maciejewska K., Wardsa J., Glowacka U., Kuter K., Mroz B., Ossowska K.

Department of Neuropsychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Motor dysfunctions in Parkinson’s disease (PD) are frequently preceded by comorbid depression. Emerging pieces of evidence suggest that BDNF alterations are associated with neurodegenerative and neuropsychiatric disorders. Postmortem studies in depressed patients showed reduced expression of mRNA and protein levels of BDNF in hippocampus (HIP) and prefrontal cortex, as well as in the serum and plasma of patients (suicide) with depression. The above effects were reversed by administration of different classes of antidepressant drugs. Similar results were observed in animal models of stress and depression.

Clinical studies indicate that dopamine D3/D2 receptor agonist – pramipexole (PRA) is the most effective compound in the treatment of depression in PD. The action of classical tricyclic antidepressant drug – imipramine (IMI) in PD patients is controversial. This drug causes a number of side-effects, including sedation and may even worsen the condition of patients.

The aim of the present study was to search for the mechanisms involved in the “depressive-like” behavior in an animal model of depression in the preclinical phase of PD.

Male Wistar rats were bilaterally injected with 6-OHDA (15 µg/2.5 µl) into the ventral striatum (vSTR). IMI was injected *i.p.* (10 mg/kg) once a day and PRA *s.c.* (1 mg/kg) twice a day for two weeks. On the 15th day after the surgery (24 h after the last drug injection) the forced swimming test (FST) and motility were examined. After decapitation the BDNF and trkB mRNAs were investigated using *in situ* hybridization.

The lesion of vSTR induced by 6-OHDA was moderate (ca. 50% of STR dopamine level), increased immobility in FST, but did not influence motility of rats suggesting

“depressive-like” changes without motor disturbances. PRA, but not IMI, shortened the immobility in FST in lesioned rats. 6-OHDA administration reduced the BDNF and/or trkB mRNAs in HIP, amygdala and habenula, while IMI reversed the lesion effect in the dentate gyrus of HIP. PRA consistently reduced the BDNF and trkB mRNAs in mesolimbic and nigrostriatal systems in lesioned animals.

These results seem to indicate that the presymptomatic dopaminergic lesion may induce “depressive-like” symptoms which are reversed by PRA but not by IMI. The PRA-induced reduction of BDNF-trkB signaling pathway may be related to its clinical efficacy.

[A5]

The N-terminal portion of mitofusin 2 inhibits proliferation of cells

Boratynska-Jasinska A.^{1*}, Kawalec M.^{1*}, Beresewicz M.¹, Kabzinska D.², Zablocka B.¹

¹Molecular Biology Unit, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Neuromuscular Unit, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

*These authors contributed equally to this work.

Mitofusin 2 (Mfn2) is a nuclear encoded protein involved in mitochondrial fusion. It has two C-terminal transmembrane domains that anchor Mfn2 in the mitochondrial outer membrane, while both the NH₂- and COOH-terminal ends are exposed to cytoplasm.

Proper mitochondrial function is maintained by mitochondrial dynamics that is the balance between mitochondrial fusion and fission. Disruption of this balance, resulting in excessive mitochondrial fission has been suggested to have a role in apoptosis activation by facilitation of Bax insertion and activation in mitochondria. It has been known for a long time that mutations in MFN2 gene are associated with Charcot-Marie-Tooth disease type 2A (CMT2A). On the other hand, a recent cancer studies have shown that Mfn2 may be responsible for inhibition of cell proliferation. Over-expression of Mfn2 protein has an anti-proliferative effect in hepatoma cells and exerts pro-apoptotic effect in hepatocellular carcinoma cells. It has also been shown that the expression of Mfn2 gene is reduced in actively proliferating cancer cells. On this basis, we decided to study the influence of Mfn2 mutations on mouse embryonic fibroblasts (MEFs) viability. We tested here: (i) proliferation rate of MEFs: wild type MEF (MEFwt) and Mfn2-null MEF (MEF^{Mfn2^{-/-}}); (ii) the effect of over-expres-

sion of wild type human Mfn2 gene and (iii) the effect of over-expression of mutated (R400X and R250W) Mfn2. These mutations were detected in Polish patients diagnosed with CMT2A. In this study we observed that depletion of Mfn2 gene in MEF^{Mfn2-/-} increased proliferation rate, significantly reducing their doubling time in comparison to MEFwt. We found that over-expression of wild type Mfn2 gene in these knock-out cells decreased their growth rate to that observed in wild type MEF. Interestingly, the expression of R400X Mfn2 mutant in MEF^{Mfn-2/-2} resulted in further elongation of doubling time of MEF^{Mfn-2/-2}. Additionally, by Western Blot, we showed that the mutation R400X generates truncated, 45 kDa, N-terminal part of mitofusin 2. Such a protein does not possess transmembrane domains located downstream of HR1 region and thus should not influence fusion of mitochondria. Taken together, these findings indicate that Mfn2 has a significant anti-proliferative effect in MEF cells and this function is mainly exerted by N-terminal part of the protein without engagement of its mitochondrial fusion activity.

This project was supported by NSC grant NN402474640.

[A6]

Neuroprotective effects of metabotropic glutamate receptors group II (mGluR2/3) agonists preconditioning in animal model of birth asphyxia: Investigations into molecular mechanisms

Bratek E., Ziembowicz A., Salinska E.

Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Hypoxic-ischemic encephalopathy is an abnormal neurobehavioral state in which the predominant pathogenic mechanism is impaired cerebral blood flow that may result in neonatal death or be manifested later as cerebral palsy or mental deficiency. Recent investigations have not provided us with promising neuroprotective compounds to reduce perinatal hypoxic-ischemic (HI) brain injury and there is a growing interest in engaging endogenous neuroprotective mechanisms. mGluR2/3 activation before or after ischemic insult results in neuroprotection and the exact mechanism of this effect is not clear. The aim of proposed study is to investigate whether mGluR2/3 activation before or after hypoxia-ischemia reduces brain damage and if induced ischemic tolerance acts through activation

of antioxidant enzymes and decrease of oxidative stress. We used an animal model of hypoxia-ischemia (H-I) on 7-day old rat pups. Animals were anesthetized and the left common carotid artery was isolated and double – ligated and then cut between the ligatures. After completion of the surgical procedure the pups were subjected to hypoxia (7.2-7.4% oxygen in nitrogen for 75 min at 35°C). Control pups were sham-operated (anaesthetized and left c.c.a. dissected, but not ligated). Animals were injected intraperitoneal with specific mGluR2 (LY 379268) and mGluR3 (NAAG) agonists 24 and 1 hour before H-I (5 mg/kg of body weight). Weight deficit of the ischemic brain hemisphere, radical oxygen species (ROS) content, activity of antioxidant enzymes (superoxide dismutase – SOD, glutathione peroxidase – GPx, catalase – CAT) and level of reduced glutathione (GSH) were measured.

Our results show a neuroprotective effect of mGluR2/3 agonists. Both agonists decreased tissue weight loss in ischemic hemisphere independently on the time of application (from 40% in H-I to 10-20% in treated). H-I resulted in increased activity of SOD in the injured hemisphere, which was further potentiated by mGluR2/3 agonists application 1 h before H-I. The activity of glutathione peroxidase (GPX) was also increased by H-I and its activity was potentiated by both agonists. The decrease by H-I GSH level observed in ischemic hemisphere was not observed after agonists application, the effect was better expressed when agonists were administered 1 h before H-I. However, CAT activity was positively modulated only by mGluR2 agonist – LY379268. Our results show that both mGluR2/3 antagonists reduce ROS level in the injured hemisphere (HI).

Conclusions: The results show that activation of mGluR2 and mGluR3 before H-I insult triggered neuroprotective mechanisms, which probably partly engage defense against oxidative stress by activation of antioxidant enzymes. The effect is more distinct when agonists are applied in a short time before H-I.

This work was made under MMRC-KNOW 2013-2017 project.

[A7]

Photic modulation of infra slow oscillatory activity in the intergeniculate leaflet of the rat model of absence epilepsy

Chrobok L., Palus K., Orłowska-Feuer P., Lewandowski M.H.

Department of Neurophysiology and Chronobiology, Institute of Zoology, Jagiellonian University, Cracow, Poland

The intergeniculate leaflet of the thalamus (IGL) is a small retinorecipient brain nucleus participating in the regulations of circadian functions. The IGL consists of at least two GABAergic neuronal populations expressing either neuropeptide Y (NPY) or enkephalin (ENK). NPY-positive cells project to the main rhythm generator – the suprachiasmatic nucleus (SCN), whereas ENK-positive cells send their axons to the contralateral IGL. ENK-positive neurons were shown to receive direct retinal input and generate action potentials in an infra slow oscillatory (ISO) manner with a period of multiseconds during *in vivo* recordings. This characteristic pattern of activity consists of two phases: intraburst, with the high firing rate, and extraburst, in which the neuron is quiet. Interestingly, ISO depends on the contralateral retina activity, as it is completely abolished by its deactivation and is only present under photopic conditions.

Absence epilepsy (AE) is characterised by the presence of spike-wave discharges (SWDs) in EEG. WAG/Rij rats, used in the present study, are well established model of AE. This neurological disease is often connected with retinal pathologies and sleep-wake cycle abnormalities. The occurrence of SWDs is modulated by the circadian timing system. The aim of the present study was to verify the presence of ISO activity in the IGL of WAG/Rij rats and whether it is modulated by light.

The experiments were performed on urethan-anaesthetised male WAG/Rij rats by using *in vivo* single-unit extracellular recordings with simultaneous ECoG measurements. ECoG electrode was placed above the frontal cortical area, whereas the recording electrode was placed in the IGL. Experimental protocol consisted of two stages. First, the baseline ISO activity was recorded in the IGL, during REM- and NREM-like phases of urethan sleep. Next, the light was turned off and the activity of the tested neuron was recorded for at least the duration of both phases.

Obtained results confirmed the presence of infra slow oscillatory activity in the IGL of WAG/Rij rats. The period of oscillations was in the range of multiseconds. The characteristic of oscillations was comparable to ISO activity described previously in the IGL of Wistar rats. Two phases

of oscillations: intra- and extra-bursts, were clearly distinguishable. In contrast to Wistar rats, all of the recorded ISO neurons remained active during the dark conditions. The features of oscillations, such as the period and mean intra- and extra-burst activity were strongly influenced by light. Namely, the period was shortened and activity was decreased. The relationship between the ISO activity and phases of urethane sleep was observed.

We hypothesise that this response to light of ISO neurons in the IGL may be due to both: abnormalities at the level of the retina and disinhibition of the IGL network itself. Our parallel *in vitro* patch clamp studies indicate that the IGL's GABAergic synaptic transmission in WAG/Rij rats is down-regulated. Based on the acquired results, further studies on the IGL of WAG/Rij can serve as a novel insight into absence epilepsy.

Supported by the Ministry of Science and Higher Education project “Diamentowy grant III” K/PMI/000190, 2014-2017.

[A8]

Neuroprotective potential of KU55933, an inhibitor of ATM kinase, against hydrogen peroxide- and doxorubicin-induced cell death in human neuroblastoma SH-SY5Y cells

Chwastek J., Jantas D., Lason W.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

The experimental data collected during the recent years point to additional functions of ATM (ataxia-telangiectasia mutated) kinase apart its involvement in DNA repair system. Since the engagement of this protein in regulation of cellular response to oxidative stress, insulin signaling and mitochondrial homeostasis has been suggested, it started to be the subject of research in diabetes, cancer and neurodegenerative diseases. In order to widen the knowledge about the role of ATM kinase in neurodegeneration, in the present study we tested the effect of pharmacological inhibition of ATM kinase by specific inhibitor KU55933 on cell death of human neuroblastoma SH-SY5Y cells, induced by oxidative stress (hydrogen peroxide) and pro-apoptotic factors – staurosporine and doxorubicin (Dox). The data showed that KU55933 (0.01-20 μ M) being not toxic when given alone to undifferentiated (UN-) and retinoic acid (RA-)-differentiated SH-SY5Y

cells, partially attenuated the cell death induced by hydrogen peroxide in both cell types (about 19-25%), as was confirmed by MTT reduction assay. The protection evoked by KU55933 against oxidative stress-evoked cell damage was relatively higher in RA-SH-SY5Y, comparing to UN-SH-SY5Y, when given the effective concentrations (0.1-10 μ M and 0.1-1 μ M for RA- and UN-SH-SY5Y cells, respectively). Moreover, KU55933 (0.1-10 μ M) almost completely blocked the cell damage induced by Dox in RA-SH-SY5Y, being only slightly protective in UN-SH-SY5Y cells (1 μ M with 10% protection). In contrast, KU55933 (0.01-10 μ M) did not affect cell damage evoked by staurosporine in both cell types. Further studies showed, that KU55933 completely blocked the Dox-induced caspase-3 activity in both, UN- and RA-SH-SY5Y cells, whereas did not change the hydrogen peroxide-evoked increase in this enzyme activity. The higher neuroprotective effectiveness of KU5593 in RA- vs. UN-SH-SY5Y was not related to changes in ATM mRNA abundance since we did not observe any significant differences in *Atm* gene expression between both cell types as was confirmed by quantitative PCR method. Finally, we demonstrated that KU5593 at higher concentration (20 μ M) significantly increased the cell death induced by hydrogen peroxide, doxorubicin and staurosporine but only in UN-SH-SY5Y cells. Altogether our data point to neuroprotective potential of ATM kinase inhibitor, KU55933 against hydrogen peroxide- and doxorubicin-evoked injury in SH-SY5Y cells with engagement of caspase-3 inhibition only in the latter one. Moreover, the results showed better usefulness of retinoic acid-differentiated SH-SY5Y cells over undifferentiated one for further study of the mechanisms of ATM kinase inhibition-mediated neuroprotection.

The study was supported by statutory funds of the Institute of Pharmacology, Polish Academy of Sciences.

[A9]

Alteration of sphingosine kinase 1 by amyloid beta (1-42) and its implication in cell death

Cieslik M., Czapski G.A., Strosznajder J.B.

Department of Cellular Signaling, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Sphingosine kinase (SphK) is a key enzyme involved in maintaining sphingolipid rheostat in the brain. Deregulation of sphingolipid metabolism has been associated with accumulation of amyloid beta (A β) peptides and could play a crucial role in pathogenesis of Alzheimer's disease

(AD). Sphingolipids are main constituents of lipid rafts and important regulators of the activity of secretase β (BACE) and A β liberation from its precursor protein.

The aim of this study was to investigate the role of extracellular A β_{1-42} peptides in regulation of SphK expression and activity. Moreover, the question arose what kind of molecular events, regulated by SphK and A β_{1-42} , are responsible for cells survival or death.

In this study rat pheochromocytoma (PC12) cells were subjected for 24 and 96 h to A β_{1-42} oligomers (1 μ M) or to SphK inhibitor (SKI II 10 μ M) and other pharmacological compounds. The experiments were performed using biochemical, spectrophotometric and qRT-PCR methods.

Our data indicated that A β_{1-42} induced time-dependent alteration of SphK1 expression and activity. Exposure to A β_{1-42} for 24 h enhanced SphK activity, but after 96 h the activity was significantly reduced. Similar changes were observed for the mRNA level of Sphk1 gene, but the expression of Sphk2 gene was not changed. Both A β_{1-42} and SKI II evoked time-dependent cell death. The level of free radicals was negatively correlated with cell viability. These two compounds evoked up-regulation of the expression of antioxidative enzymes, catalase and superoxide dismutase. The significant increase in the level of apoptosis inducing factor (AIF) that could be involved in antioxidative defence in mitochondria was also observed. Moreover A β_{1-42} and SKI II evoked reduction of glycogen synthase kinase-3 β (Gsk-3 β) immunoreactivity and altered the balance between pro- and anti-apoptotic proteins. Exposure of cells to A β_{1-42} or SKI II resulted in the increase in gene expression for pro-apoptotic BAX protein and in decrease of anti-apoptotic Bcl-xL and Bcl-2 proteins. Several pharmacologically active compounds were then investigated for their cytoprotective effects against A β_{1-42} -evoked cytotoxicity, but only α -pifithrin (20 μ M), an inhibitor of p53 protein, protected cells against death.

Our data indicated that inhibition of SphK, by the alteration of sphingolipid rheostat, may be the trigger of apoptotic signaling evoked by Alzheimer's A β_{1-42} peptides.

Our study was supported by a Grant of NSC 2013/09/B/NZ3/01350.

|A10|

The effects of maternal immune activation on the behavior of rat offspring: a focus on changes in postsynaptic proteins – Shank

Cieslik M.¹, Kudlik M.¹, Boguszewski P.², Adamczyk A.¹

¹Department of Cellular Signaling, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Department of Neurophysiology, Laboratory of Limbic System, Nencki Institute of Experimental Biology, Polish Academy of Sciences, Warsaw, Poland

Autism spectrum disorders (ASDs) are neurodevelopmental diseases characterized by two core symptoms, impaired social interactions and communication, and ritualistic or repetitive behaviors. Recent human genetic studies indicate that genes identified as altered in individuals with ASDs often encode synaptic proteins. Shank family proteins (Shank1, Shank2, and Shank3), that belong to synaptic scaffolding proteins, are involved in the regulation of morphological and functional maturation of dendritic spines and synapse formation. These proteins interact with NMDA receptor and metabotropic glutamate receptor complexes in the postsynaptic density (PSD). Previous data indicated that mutation in *SHANK1* contributes to the symptoms of autism. In *Shank1(-/-)* mice impaired communication characterized by low levels of ultrasonic vocalizations, as compared to wildtype *Shank1(+/+)* littermate controls, was observed.

Both epidemiological and biochemical evidence suggests that occurrence of ASD may be linked to immune perturbations that took place during fetal development. Therefore, “maternal immune activation” (MIA) model, where the research is carried out in offspring from female rodents subjected to an immune stimulus during early pregnancy, is commonly used as animal models of ASD. The present experiments were performed using male offspring of Wistar rats treated intraperitoneally with saline or lipopolysaccharide (LPS, 100 µg/kg) on gestational day 9.5, to mimic a bacterial infection. Autism-associated behaviors and the function of postsynaptic proteins Shank were examined in the offspring of the treated animals. Behavioral tests were conducted to assess social communication, motions and anxiety, play behaviors, as well as learning and memory. Data analysis was carried out exclusively on male rats. Gene expression and protein level of Shank was analyzed using quantitative Real-Time RT-PCR and Western blot methods, respectively. We observed that male offspring from LPS-treated animals showed impaired communication (expressed by ultrasonic vocalization on postnatal day 9-13, PND 9-13). Moreover, in the spon-

taneous alternation (T-maze) test, adult male offspring (PND 40-50) from LPS-treated mothers showed impaired learning and memory function. The motions and anxiety responses (Open space test, PND 40-50) and play behaviors (Tickling, PND 40-50) were not altered. When the experimental animals have reached adulthood, the changes in the expression of *SHANK1*, *SHANK2* and *SHANK3* was investigated. The data showed the down-regulation of *SHANK1* and *SHANK2* expression in the cerebral cortex, without changes in *SHANK3* expression in all analyzed brain structures. Summarizing, our findings indicate that offspring from mothers subjected to immune stimulation during gestation show an impaired communication and dysregulation of synaptic proteins belonging to Shank family. The lower expression of Shank proteins may lead to profound alterations in synaptic function and social behavior.

Supported from IMDIK statutory theme 17.

|A11|

Effect of prenatal stress on spleen cytokine production in female offspring

Curzytek K., Duda W., Kubera M.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Numerous studies have shown that stressful events occurring in early life have a powerful influence on the development of the central nervous system. Prenatal stress also modifies the immune response of the offspring. It is known that chronic activation of immune-inflammatory pathway play an important role in the pathophysiology of depression. The spleen, as a crucial component of the immune system, can produce a lot of agents i.e. cytokines, chemokines and neurotrophic factors. Proinflammatory cytokines and chemokines produced by the spleen can enhance depressive-like behaviours via influencing synaptic transmission and neuronal plasticity. The aim of present study was to evaluate the disrupting effect of prenatal stress on spleen releasing cytokines and other factors in female offspring.

Pregnant Sprague-Dawley rats were subjected daily to three stress sessions (immobilization) from day 14 of pregnancy until delivery. At 4 months of age, adult female offspring of control and stressed rat dams were subjected to behavioural testing (forced swim test). After other behavioural tests (i.e. locomotor activity, elevated plus

maze test), animals were killed by rapid decapitation and the brain tissue and organs were dissected. Spleen derived cells were cultured with concavalin A (Con A, 2.5 µg/ml) for 24 hours, and supernatants were collected. The level of spleen cytokines was measured by RayBio Rat Cytokine Antibody Array – the protein array system. Our studies showed that prenatal stress caused increased ability of splenocytes to produce Fas-ligand, fractalkine, granulocyte-macrophage colony-stimulating factor (GM-CSF), monocyte chemoattractant protein-1 (MCP-1), tumor necrosis factor (TNF)- α and decreased to produce interleukin (IL)-1 α , IL-10, IL-6 and tissue inhibitor of metalloproteinase-1 (TIMP-1). Our studies revealed that prenatal stress significantly changed secretory spleen functions, that can be crucial in disturbance of homeostasis in central nervous system.

The study was supported by Grant POIG.01.01.02-12-004/09 (DeMeTer).

[A12]

Cyclin dependent kinase 5 and its modulatory role in transcription of inflammation-related genes in the brain. Relevance to pathomechanism of Alzheimer's disease

Czapski G., Wilkaniec A., Gassowska M., Strosznajder J.B., Adamczyk A.

Department of Cellular Signalling, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Epidemiological and genetic studies demonstrated an important role of inflammatory reaction in the pathomechanism of Alzheimer's disease (AD), but the precise course of immune system activation and characterization of regulatory processes have not been fully elucidated. In this study we focused on the role of Cyclin dependent kinase 5 (Cdk5) in inflammatory signaling. Cdk5 is serine/threonine kinase that is involved in regulation of neuronal development and function. Moreover, it has been implicated in pathomechanism of AD and other neurodegenerative disorders. Overactivation of Cdk5 has deleterious effects on neuronal cells, leading to hyperphosphorylation of MAP tau, which is responsible for microtubule destabilization and impairment of axonal transport. Additionally, inhibition of this enzyme was proposed to be a prospective strategy preventing neurodegeneration.

The aim of this study was to analyze the role of Cdk5 in neuroinflammatory signaling in animal model of AD. Adult male mice C57BL6 received single intracerebroventricular injection of A β ₁₋₄₂ oligomers (1 nmol). Lipopolysaccharide (LPS) was administered intraperitoneally (ip.) to evoke activation of inflammatory reaction. Roscovitine (50 mg/kg b.w.) was applied *i.p.*, as an inhibitor of Cdk5. Alterations of gene transcription were analyzed using real-time RT-PCR. The level and phosphorylation of proteins was examined by Western blotting (WB) and by using multiplexed cytometric bead array (CBA).

Our results demonstrated that LPS-induced neuroinflammatory processes evoked rapid increase in phosphorylation of tau protein at Ser396 in hippocampus. In parallel, we observed enhanced phosphorylation of Glycogen synthase kinase-3 β (Gsk-3 β) at Ser9, indicating inhibition of this tau-kinase. However, the rise of phosphorylation of Cdk5 and acceleration of Calpain-dependent p25 formation suggested that overactivation of Cdk5 may be responsible for hyperphosphorylation of tau protein during neuroinflammation. By using an animal model of AD we have found that A β peptides evoked activation of Cdk5 via p25 formation and by enhancement of Cdk5 phosphorylation at Tyr15. Moreover, administration of A β induced rapid increase in expression of proinflammatory genes for Tnfa and Il1b, Il6, and concomitant rise in mRNA level for anti-inflammatory Il10. Additionally, an increase in the level of mRNA for Nos2 was observed, suggesting activation of nitrosative/oxidative processes in hippocampus. By using WB and CBA methods we have found that Roscovitine significantly affected A β -induced inflammatory signaling in brain. Pre-treatment with Roscovitine reduced the level of Il1b in the mouse hippocampus 3 h after administration of A β .

Our results demonstrated for the first time that inhibition of Cdk5 significantly affects A β -evoked expression of inflammation-related genes in mouse hippocampus. These data indicate the critical role of Cdk5 in regulation of acute neuroinflammatory signaling in the brain.

Supported by a grant from The National Science Centre 2011/03/B/NZ3/04549.

|A13|**Involvement of huntingtin associated protein 1 in regulation of SOCE in medium spiny neurons from transgenic YAC128 mice model of Huntington’s disease****Czeredys M.¹, Maciag F.^{1,2}, Kuznicki J.¹**¹International Institute of Molecular and Cell Biology in Warsaw, Laboratory of Neurodegeneration, Warsaw, Poland²Warsaw University of Technology, Faculty of Chemistry, Warsaw, Poland

Huntington’s disease (HD) is a hereditary neurodegenerative disease caused by the expansion of a polyglutamine stretch in the huntingtin (HTT) protein, and characterized by dysregulated calcium homeostasis (Gaicomello *et al.* 2013). One of the mechanisms that regulate calcium homeostasis is store-operated Ca²⁺ entry (SOCE). It is a process in which the depletion of Ca²⁺ stores in the endoplasmic reticulum (ER) induces Ca²⁺ influx from the extracellular space. The Ca²⁺ sensor’s stromal interaction molecules 1 and 2 (STIM1 and STIM2) detect changes in Ca²⁺ concentration in the ER. In response to Ca²⁺ depletion, they rearrange into punctate structures by interacting with ORAI calcium channels in plasma membrane (PM). These structures force ER membranes toward PM and activate members of the Orai family (Orai1, Orai2, and Orai3). This results in Ca²⁺ entry by these channels (Frischauf *et al.*, 2008). The mechanism by which mutated HTT affects SOCE is unknown. Therefore, we assessed alterations of Ca²⁺ signalosome in the striatum of a transgenic YAC128 mouse, a model of HD. We found that in medium spiny neurons (MSN) from these mice, the activity of SOCE was enhanced about 25%, which confirmed previous results of Wu (2011). This could be explained by increased expression of huntingtin-associated protein 1 (Hap1) mRNA and Hap1 protein in the striatum of YAC128 mice (Czeredys *et al.* 2013) and may explain data of Wu *et al.* (2011) and Tang *et al.* (2003, 2004). In this work we are investigating the involvement of Hap1 in the regulation of SOCE in neurons from YAC128 mice. Using western blot we found that Hap1 protein is specific for MSN neurons, and was not found in primary cultures of astrocytes. We prepared constructs in lentiviral system for Hap1 overexpression in HEK293 cells. We are now analyzing if the increased level of Hap1 affects SOCE in HD neurons.

|A14|**Ceramide and sphingosine-1-phosphate the critical regulator of the PI3-K/Akt/GSK-3 β signaling pathway and cell fate****Czubowicz K., Strosznajder R.P.**

Laboratory of Preclinical Research and Environmental Agents, Department of Neurosurgery, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Bioactive sphingolipids: ceramides, sphingosine and sphingosine-1-phosphate (S1P) are crucial regulators of cell survival and death. S1P promotes cell proliferation and viability acting as primary and secondary messenger and exerts its effect by binding to five specific G protein coupled-receptors (S1PR1-5). Its precursors ceramide/sphingosine activate the pro-apoptotic pathways. The relationship between these sphingolipids and pro-survival pathway regulated by PI3-kinase/Akt is till now not fully elucidated. The PI3-K/Akt mediates a wide range of cellular functions, including cell proliferation, survival and cell metabolism. Alteration of ceramide/S1P biostat and signaling is implicated in brain ischemia, cancer, and also neurodegenerative diseases including Alzheimer’s disease.

The aim of this study was to investigate the role of S1P in molecular mechanisms of neuronal cells death in the model of oxidative stress evoked by cell-permeable C2-ceramide. The experiments were carried out on human neuroblastoma SH-SY5Y cells, using spectrometric and immunochemical methods. Our study indicated that C2-ceramide (25 μ M) inhibited the pro-survival PI3-K/Akt pathway followed by activation of glycogen synthase kinase-beta (GSK3 β). Moreover, after ceramide treatment we observed the lower phosphorylation of pro-apoptotic Bad protein. Conversely, exogenously added S1P (1 μ M) increased the viability of neuronal cells through S1PR (1, 3) receptors-dependent mechanism as was demonstrated by us using specific receptors antagonists (W123 and VPC23019). In addition S1P and the agonist of S1P receptor(s) (SEW2871 and pFTY720) increased the activity of PI3-K/Akt. Furthermore, it was revealed that S1P enhanced the phosphorylation of Bad protein. Summarizing, our data indicate that modulation of S1P receptor(s) signaling may offer therapeutic strategy in treatment of neurodegenerative diseases characterized by accumulation of ceramides.

Supported by statutory budget of MRC theme no. 1.

[A15]

Neuroprotective potential of neuropeptide Y receptor agonists in ischemic *in vitro* model

Domin H.¹, Jantas D.², Smialowska M.¹

¹Department of Neurobiology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Much of evidence indicates that neuropeptide Y (NPY) and Y receptors agonists may have neuroprotective potential against excitotoxic damage, however little is known about their effect in ischemic neurodegenerations. Our previous studies demonstrated that Y2 receptor agonist, NPY 13-36, given intracerebroventricularly 30 min after the onset of ischemia, significantly diminished the infarct volume in the rat transient middle cerebral artery occlusion (MCAO) model. The present study was aimed at investigating the role of Y1, Y2 and Y5 receptors in neuroprotection in primary neuronal cortical cultures, exposed to oxygen-glucose deprivation (OGD) for 3h as an *in vitro* ischemic injury paradigm. Therefore we evaluated the effects of YR agonists (0.01-1 μ M) such as Y2R agonist (NPY 13-36), Y5R agonist [cPP¹⁻⁷, NPY¹⁹⁻²³, Ala³¹, AiB³², Gln³⁴]-hpp or Y1R agonist [Leu³¹, Pro³⁴]-NPY on ischemia-induced LDH release and MTT reduction in the cortical cultures 24 h after the termination of OGD. The YR agonists were applied twice: just before the start of OGD and immediately after the end of OGD, or once: 30 min or 1 h after the termination of OGD. It was found that double application of the Y2R or the Y5R agonists significantly attenuated neurotoxicity. Of particular importance is the finding that those Y2R or Y5R agonists, given once 30 min after OGD also significantly decreased ischemic injury. The neuroprotective effects of Y2R or Y5R agonists were inhibited by corresponding antagonists to those receptors. No protection was found after the Y1 receptor agonist. We also observed a significant induction of calpains after OGD, evidenced by an increase in spectrin alpha II 145 kDa cleavage product, and that effect was diminished by double application of Y2R agonist, NPY 13-36, but not by Y5R agonist. In conclusion, our results indicate that particularly Y2 receptors may play an important role in the attenuation of damage after ischemia.

The study was supported by grant No. N N401 091037 from MS&HE and by funds for statutory activity of the Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland.

[A16]

Genetic variants and oxidative stress in peripheral lymphocytes of the patients with Alzheimer's disease

Dorszewska J.¹, Oczkowska A.¹, Prendecki M.¹, Florczak-Wyspianska J.², Dezor M.¹, Karlikowska M.^{1,3}, Postrach I.¹, Kozubski W.²

¹Laboratory of Neurobiology, Department of Neurology, Poznan University of Medical Sciences, Poznan, Poland

²Chair and Department of Neurology, Poznan University of Medical Sciences, Poznan, Poland

³University College of London, London, United Kingdom

Alzheimer's disease (AD) leads to generation of β -amyloid ($A\beta$) and oxidative stress. Oxidative stress and apolipoprotein E (ApoE) are associated with DNA damage, which leads to apoptosis induction e.g. in cells expressing wild-type p53. The *TP53* gene is found to be the most frequently mutated gene in human tumor cells. Post-mortem studies have indicated that patients diagnosed with a brain tumor also had unreported signs of AD. In AD, apoptosis can also be induced by the immune system elements, e.g. tumor necrosis factor alpha (TNF-alpha). 8-Oxoguanine DNA glycosylase 1 (OGG1) is a main DNA repair enzyme that excises of 8-oxo-2'-deoxyguanosine (8-oxo2dG) from DNA.

The purpose of this study was to analyze the mutation status in highly conservative exon 7 of *TP53* gene and *APOE* genotypes and the extent of oxidative DNA damage (8-oxo2dG), and expression of p53, OGG1 and TNF- α protein levels in peripheral lymphocytes of AD patients and controls.

The studies were conducted on 31 patients with AD, including 18 women and 13 men aging 52-85 years. The control group included 31 individuals, 18 women and 13 men aging 40-83 years. The *TP53* exon 7 was analyzed by DNA sequencing. The *APOE* genotyping analysis was carried out with the use of Real-Time PCR and DNA sequencing. The levels of 8-oxo2dG were determined using HPLC/EC/UV (high-pressure liquid chromatography system with electrochemical and UV detection) technique. The p53, OGG1 and TNF- α protein levels were analyzed by Western blotting.

Our studies on the sequencing of *TP53* exon 7 revealed the presence of two various mutations only in AD patients, a missense mutation C748A (3.4%) present in an *APOE* ϵ 3/ ϵ 4 alleles carrier and a silent mutation C708T (20.1%) in AD patients with various *APOE* genotypes, ϵ 2/ ϵ 3, ϵ 3/ ϵ 3, ϵ 3/ ϵ 4. Only a person (control subject and AD patient) with one or two alleles of *APOE* ϵ 2 had few times higher level of 8-oxo-2dG compared to non-carriers of this allele. Moreover, in AD patients with *APOE* ϵ 2/ ϵ 3 genotype and a missense

mutation of *TP53* gene (15 times) and a silent mutation of *TP53* gene (2 times) the level of 8-oxo2dG was higher than in patients without analyzed *TP53* gene mutations. However, in the controls with *APOE* $\epsilon 3/\epsilon 4$ genotype the level of 8-oxo2dG and OGG1 was nearly twice higher compared to subjects with the *APOE* $\epsilon 3/\epsilon 3$ genotype. In contrast, in AD patients with *APOE* $\epsilon 3/\epsilon 4$ genotype 8-oxo2dG level was 1.5 times lower than in patients with *APOE* $\epsilon 3/\epsilon 3$ genotype, although the level of OGG1 in both AD groups was decreased compared to the appropriate controls. In the controls and AD, p53 and TNF- α protein levels were similar, regardless of *APOE* genotype.

It appears that especially *APOE* $\epsilon 2$ and $\epsilon 4$ together with *TP53* gene mutations may be associated with the development of oxidative stress in AD.

[A17]

Cytotoxic effects of zinc on cholinergic SN56 neuroblastoma and C6 astroglia cells

Dys A., Ronowska A., Gul-Hinc S., Klimaszewska-Lata J., Szutowicz A., Bielarczyk H.

Department of Laboratory Medicine, Medical University of Gdansk, Gdansk, Poland

Zinc excess in the synaptic cleft may be one of early pathologic signals triggering chronic neurodegenerative events. The aim of this work was to find relationships between Zn accumulation and integrity of cholinergic and astroglial cells. Exposition of cAMP/RA-differentiated (DC) and nondifferentiated (NC) cholinergic SN56 neuroblastoma cells and astroglial C6 cells to Zn yielded its concentration dependent accumulation. Zn accumulation caused concentration-dependent death of both, neuronal and astroglial cells. After 24 h exposition of SN56 cells to 0.15 mmol/L Zn, their mortality was equal to 35% and 50% at cation levels equal to 4.0 and 5.5 nmol/mg protein for NC and DC, respectively. In these conditions, the mortality of astroglial cells was close to 1-2% only, at intracellular Zn levels of 1.6 and 2.1 nmol/mg protein for NC and DC, respectively. Higher, about 0.25 mmol/L Zn concentration was required to evoke mortality of astroglial cells, similar to those seen in neuronal cells. In such conditions Zn levels in astroglia were about 6.4 and 27.0 nmol/mg protein, respectively. The lower accumulation of Zn in astroglial cells C6 than in neuronal cells SN56 may be due to the less density of voltage-gated calcium channels and higher density of ZnT1. In neuronal cells, Zn caused a decrease of mitochondrial Ca²⁺ content and increased level of Ca²⁺

in the cytoplasmic compartment, which may be one of the mechanisms of Zn-dependent increase in mortality of cholinergic neurons. In this study we examined the effects of accumulation of Zn in cholinergic neurons and adjacent astrocytes on activity of enzymes involved in energy metabolism, including PDHC, aconitase and IDH. One day culture of neuronal cells SN56 with 0.15 mmol/L Zn caused about 60% suppression of PDHC and aconitase activities. In these conditions we observed no changes in activity of these enzymes in astroglial cells. In C6, such as 0.22 mmol/L Zn concentration caused 60% and 50% inhibition of aconitase and IDH activities, respectively. The high susceptibility of cholinergic neurons and a relative high resistance of astrocytes to neurodegeneration, induced by cytotoxic concentration of zinc, may be caused by the higher inhibition of neuronal aconitase and PDHC activities. The inhibition of these enzymes reduces the synthesis of acetyl-CoA, and the flow rate of the tricarboxylic acid cycle, which leads to a decrease in ATP synthesis and cell damage. These results indicate, that zinc overload that leads to the disturbance of energy metabolism may be one of the causes of early damage of neuronal cells yielding to the development of neurodegenerative diseases.

Supported by MWiSW St57 and MN 01-0108/08 MUG projects.

[A18]

Neuroprotective effects of memantine, hyperbaric oxygen or hypobaric hypoxia do not add up when applied together in 7-day-old rat hypoxia-ischemia model

Gamdzik M., Ziembowicz A., Salinska E.

Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Despite major advances in monitoring technology and knowledge of fetal and neonatal pathologies, perinatal asphyxia or, more appropriately, hypoxic-ischemic encephalopathy (HIE), remains a serious condition that causes significant mortality and long-term morbidity. In our previous studies we proved that hyperbaric oxygen (HBO) and hypobaric hypoxia postconditioning (HH) reduce brain damage in experimental model of neonatal hypoxia-ischemia by over 60%. The aim of present study was to evaluate the effect of NMDA receptor (NMDAR) antagonist

memantine, and memantine combined with HBO or HH on neonatal hypoxic-ischemic (HI) brain injury.

We used an established 7-day postnatal rat model of hypoxia-ischemia. Briefly, the ipsilateral common carotid artery of a rat pup is exposed and ligated. The contralateral artery is left intact. The animal is then subjected to a period of hypoxia at normal atmospheric pressure (7.7% O₂ for 75 min at 35.5°C). HBO (2.5 ATA) or HH (0.5 ATA) were applied 1 or 6 h after H-I for 60 min. Memantine in dose of 20 mg/kg of body weight was applied 15 min before HBO or HH. These treatments were repeated for 3 following days.

Memantine applied 1 or 6 h after H-I reduced ipsilateral hemisphere weight loss from 39.2% to 19.3% and 26.8%, respectively. Memantine combined with HBO decreased brain weight loss to 25% and 28.2%, when applied 1 and 6 h after H-I, respectively. Similarly, memantine combined with HH decreased weight loss to 21.9% and 28.8%.

Memantine applied 1 or 6 h after H-I reduced number of TUNEL-positive cells in ipsilateral hemispheres by 62 and 58%, respectively. Memantine combined with HBO or HH decreased number of positive cells by 21% and 10% (HBO), and 39% and 36% (HH), respectively. Memantine combined with HBO or HH caused protection, but the effect was less significant than effect of memantine used alone.

Our results show that HBO and HH do not increase protection caused by memantine alone. It is possible that HBO and HH, which are means of manipulation of oxygen concentration, modulate NMDAR activity and alter balance between synaptic and extrasynaptic NMDAR activity when applied in combination with memantine. There is some data showing that high dose of memantine can block not only damaging extrasynaptic activity, but also normal synaptic transmission, which interferes with pro-survival protective pathways dependent on synaptic activity, causing detrimental effects leading to cell death, which was visualized by TUNEL staining. This effect could be strengthened or triggered by HBO and HH, which would explain why memantine alone was more neuroprotective.

This project was funded by the National Science Centre based on decision no. DEC-2012/07/N/NZ4/02072.

[A19]

Pre-chiasmatic subarachnoid hemorrhage leads to tight junction disassemblance and enhanced neurodegeneration in experimental rat brain

Gendosz D., Liskiewicz A., Jedrzejowska-Szypulka H., Lewin-Kowalik J.

Medical University of Silesia, School of Medicine, Chair and Department of Physiology, Katowice, Poland

Aims: Our aim was to develop model of arterial bleeding into the pre-chiasmatic cistern resembling subarachnoid hemorrhage (SAH) from aneurysm rupture in the anterior part of the circle of Willis and to explore different aspects of its effect upon adjacent as well as distant areas of rat brain and extracellular.

Methods: Pre-chiasmatic SAH (pSAH) was produced by injection of 200 µL of fresh autologous arterial blood into pre-chiasmatic cistern in rat brain. 24 hrs following the surgery, animals were perfused transcardially and whole brains were collected. Brains were coronally cryosectioned and stained with Fluoro Jade C (FJC) and immunostained to visualize cleaved caspase 3, GFAP, zonulin-1, occludin-1 and claudin-5.

Results: We found numerous FJC positive neurons around the site of blood application as well in more distant areas and its number was especially high in neocortex and hippocampus. Distribution of caspase 3 positive cells was similar to that for FJC. To examine the presence of astrogliosis we carried out GFAP staining. We found numerous astrocytes with morphological alterations characteristic for their activation: thick processes and enlarged cell bodies. Another observations acquired from immunostaining reactions against proteins building tight junctions of BBB suggest their degradation after SAH comparing to controls.

Conclusions: Obtained results indicate that administration of arterial blood directly to pre-chiasmatic cistern leads to neuronal damage and stimulation of astrogliosis not only in the vicinity of injection site but also in more distant regions like somatosensory cortex and hippocampus. Moreover obtained results suggest also SAH crucial role in tight junction proteins degradation.

[A20]

The influence of kissorphan (KSO) on the expression of sensitization to hyperlocomotor effect of morphine and ethanol in mice

Gibula-Bruzda E.¹, Kotlinska J.H.¹, Trzcinska R.², Silberring J.^{2,3}

¹Department of Pharmacology and Pharmacodynamics, Medical University, Lublin, Poland

²Centre of Polymer and Carbon Materials, Polish Academy of Sciences, Zabrze, Poland

³Faculty of Materials Science and Ceramics, AGH University of Science and Technology, Cracow, Poland

Drug reward and phenomenon of sensitization play a crucial role in the development of drug dependence. It has been proposed that sensitization exhibits an important function in the etiology and maintenance of drug seeking behavior as well as in relapse to this behavior after a period of drug abstinence (Robinson and Berridge 1993; Stewart and Badiani 1993). The endogenous brain opioid system is believed to play an important role in mediating reward effects of many psychostimulants. Therefore the aim of the present study was to investigate whether a novel derivate of kisspeptin-10 with affinity to NPF receptors – kissorphan (KSO) (Tyr-Asn-Trp-Asn-Ser-Phe-NH₂) affects the acute hyperlocomotor effect of morphine (10 mg/kg, s.c.) and ethanol (2.4 g/kg, 20% w/v, i.p.), and influences the expression of morphine- and ethanol-induced sensitization to their hyperlocomotor effects. Sensitization to the hyperlocomotor effect of ethanol and morphine was induced according to the previously established methods (Kotlinska *et al.* 2007).

Our study indicated that acute administration of KSO (1, 3 and 10 nmol/300 µl, intravenously (*i.v.*)) inhibited the expression of morphine- and ethanol-induced sensitization at all used doses. Furthermore, KSO inhibited the acute hyperlocomotor effect of morphine at doses of 3 and 10 nmol/300 µl and acute hyperlocomotor effect of ethanol at the dose of 10 nmol/300 µl. KSO, given alone, did not change the locomotor activity of mice. In conclusion, our experiments indicated that KSO attenuated the acute morphine and ethanol hyperlocomotor effect and the expression of sensitization to the hyperlocomotor effect of morphine and ethanol. Therefore we anticipate that KSO may be involved in both of these effects and may reveal anti-opioid activity.

[A21]

Role of astrocytes in nigrostriatal system degeneration. HPLC analysis of dopamine and its metabolites after glia and neurons dysfunction

Glowacka U., Olech L., Kuter K.

Department of Neuropsychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Objectives: Our aim was to investigate if prolonged energetic dysfunction of astrocytes would induce dysfunction of dopaminergic neurons in substantia nigra (SN) and striatum (STR), or influence dopaminergic neurotransmission changed by selective anti-dopaminergic toxin injection.

Background: The degeneration of dopaminergic neurons in SN is underlying cause of movement disorders observed in Parkinson’s disease (PD). The role of astrocytes supporting those neurons in both energetic and trophic way is still unrecognized. It is probable that prolonged dysfunction of astrocytes could increase dopaminergic neurons vulnerability.

Material and methods: We have prepared PD rat model of selective nigrostriatal dopaminergic system degeneration by intracerebral, stereotaxic injection of 3 µg/3 µl 6-hydroxydopamine (6-OHDA) into medial forebrain bundle (MFB). Astrocytes dysfunction was induced by 7-days constant infusion of fluorocitrate (FC, 0.5 µl/hr) into SNc using osmotic ALZET minipumps. Control animals received solvent (0.2% ascorbic acid) into MFB and blank cannulas into SNc. Animals were sacrificed 1 or 4 weeks later. HPLC-EC analysis of DA, its metabolites and turnover rates in SN and STR was performed.

Results: Selective dopaminergic neurons lesion caused decrease in DA, DOPAC and 3-MT levels in STR and SN already after one week, and after 4 weeks this decrease further progressed. Increased turnover rates suggested compensatory mechanisms in STR. Induced metabolic dysfunction of astrocytes in SN caused decrease in DA level in STR, only after one week. No change in DA was observed in SN, but its metabolites DOPAC and HVA increased significantly after 7 days, and returned to almost control levels after 4 weeks. Double (glial and neuronal) dysfunction enhanced DA and metabolites decreases in STR after one week, but no further aggravation with time was detected. No enhancement of deficits was visible in SN, probably due to increased DA metabolism and turnover.

Conclusions: We conclude that astrocytes dysfunction seriously influences DA metabolism and accelerates dopa-

minergic transmission breakdown after neuronal degeneration.

The study was supported by the NCN grant nr 2012/05/B/NZ4/02599.

[A22]

Influence of caffeine and selective antagonists of adenosine A₁ and A_{2A} receptors on DA and 5-HT release induced by MDMA in the mouse striatum

Gorska A.M., Golembiowska K.

Department of Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

MDMA (3,4-methylenedioxymethamphetamine) used recreationally is the main component of ecstasy tablets which also contain another psychostimulants such as caffeine (CAF) to gain stronger effect. Amphetamines cause an increase in motor activity, anxiety, aggression. Their chronic use can lead to mental disorders like schizophrenia or depression and may influence progression of neurodegenerative diseases.

The pharmacological action of amphetamines is directed primarily at neurotransmitter systems: catecholamines – dopamine (DA), norepinephrine (NA) and serotonin (5-HT). Amphetamines cause increase in monoamines level in the synaptic cleft by blocking their reuptake and inhibit their inactivation through monoamine oxidase A and B. CAF is a weak stimulant without abuse properties. It abolishes mental fatigue and improves thinking. Pharmacological mechanism of CAF is based on blockade of adenosine A₁ and A_{2A} receptors which may affect neurotransmitter release. Our earlier data have shown that CAF (10 mg/kg) given together with MDMA (20 or 40 mg/kg) potentiated stimulatory effect of MDMA on extracellular level of DA and 5-HT in the mouse striatum.

In the present study we aimed to understand, if potentiating effect of CAF on DA and 5-HT release produced by MDMA occurs via blockade of adenosine A₁ and A_{2A} receptors.

All experiments were conducted on wild-type male mice from the breeding in the Institute of Pharmacology, PAS, weighing 20-35 g. Animals were anaesthetized with ketamine (7.5 mg/kg) and xylazine (1 mg/kg) and a vertical microdialysis probes were implanted into the striatum (AP: +1.0, L: +1.8, V: -3.8). The following drugs were injected intraperitoneally: MDMA (20 or 40 mg/kg), CAF (10 mg/kg),

KW-6002 (1.25 or 2.5 mg/kg), DPCPX (1.25 or 2.5 mg/kg). The extracellular concentration of DA, 5-HT and extraneuronal DA metabolite, 3-methoxytyramine (3-MT) was quantified by HPLC with coulochemical detection.

Both adenosine A₁ and A_{2A} receptor antagonists, similarly to CAF, enhanced DA and 5-HT release induced by MDMA. However, adenosine A₁ receptor antagonist only weakly influenced DA and 5-HT release induced by MDMA. To the contrary, adenosine A_{2A} receptor antagonist markedly increased effect of MDMA on DA and 5-HT release. It can be speculated that effect of CAF on MDMA-induced release of DA and 5-HT in the mouse striatum is mediated mainly by adenosine A_{2A} receptors.

Supported by grant from National Centre of Science (NCN) no 2013/11/N/NZ7/00740.

[A23]

Involment of calcium channels and receptors in neuronal soce

Gruszczynska-Biegala J.¹, Sladowska M.^{1,2}, Kuznicki J.¹

¹International Institute of Molecular and Cell Biology, Warsaw, Poland

²Undergraduate student from the Warsaw University of Life Sciences – SGGW, Warsaw, Poland

Changes in the intracellular calcium (Ca²⁺) concentration are considered to be one of the crucial signalling pathways in cells. Increased concentration of Ca²⁺ in cytoplasm is caused by influx from extra- or intracellular sources of Ca²⁺. In non-excitabile cells, such as lymphocytes, store-operated calcium channels (SOCCs) are the primary pathway for the Ca²⁺ influx. The extracellular Ca²⁺ influx through the tightly regulated SOCCs in the plasma membrane (PM) is the basis of the store-operated Ca²⁺ entry (SOCE). The process relies on refilling the cytoplasm and the endoplasmic reticulum (ER) with Ca²⁺ after their IP₃-dependent release to the cytoplasm. The key proteins involved in this process are sensors of Ca²⁺ located in ER (STIM1 and STIM2) and the Ca²⁺ channel-forming protein (ORAI1) present in the PM. Interaction between these proteins leads to formation of complexes, visible under the fluorescent microscopy as puncta, and results in entry of Ca²⁺ into the cytoplasm. Ca²⁺ is then transported to ER by the activity of the Ca²⁺-ATPase of SERCA pump, which refills ER stores. In contrast to non-excitabile cells, Ca²⁺ homeostasis in neurons is maintained primarily by voltage-gated calcium channels (VGCCs) as well as ion exchangers or

various receptors (e.g. metabotropic and ionotropic glutamate receptors). Recent studies demonstrate that SOCE also occurs in neurons. Our earlier data indicate that both STIMs are involved in calcium homeostasis in neurons, but each plays a distinct role in SOCE (Gruszczynska-Biegala *et al.*, PLoS ONE 2011) and can form complexes with endogenous ORAI1 (Gruszczynska-Biegala and Kuznicki, J Neurochem 2013). Recently, SOCE has become a subject of intense research in regard to disturbances in Ca²⁺ homeostasis observed in the epilepsy and neurodegenerative diseases (Steinbeck *et al.*, Exp Neurol 2011).

The aim of this study is to determine if, in addition to ORAI1, there are neuronal specific channels or receptors, which interact with STIM proteins during SOCE. The potential PM partners of STIMs, VGCC, glutamate receptors and Na⁺-Ca²⁺ exchanger, were selectively inactivated by chemicals in order to assess their role in SOCE in cortical neurons prepared from 19-day-old embryonic Wistar rat brains. After 16 days of culture, single-cell Ca²⁺ levels in neurons were recorded using the ratiometric Ca²⁺ indicator dye Fura-2AM in the presence of different inhibitors such as nimodipine (VGCC), MK-801 (NMDAR), MCGG (mGluR), bepridil (exchanger). In some experiments neurons were treated with thapsigargin to deplete the ER stores following the Ca²⁺ re-addition to assess SOCE. The difference between Ca²⁺ response after treatment of neurons with inhibitors and response without them showed whether and how much Ca²⁺ influx is sensitive to the blockers of potential STIMs partners. The discovery of the new proteins interacting with STIM proteins will allow a better understanding of the mechanisms of SOCE.

This work was supported by funds from a National Science Centre (2011/01/D/NZ3/02051, JGB).

[A24]

Influence of P2X7R antagonist on glia activation in the early time point of EAE

Grygorowicz T.¹, Lenart J.², Lenkiewicz A.¹, Struzynska L.¹

¹Laboratory of Pathoneurochemistry, Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Laboratory of Pharmacochemistry, Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Multiple sclerosis (MS) is an autoimmune, neurodegenerative disease which is the most frequently the reason of

disabilities of young adults, and is a very serious problem for contemporary medicine.

Progressively developing plaques of demyelination with oligodendrocyte and axonal death are observed during the course of MS. Within the central nervous system astroglia and microglia are important players in inflammation. Responding to the inflammatory signal, they become activated, change their morphology, release cytokines and up-regulate marker-proteins such as GFAP and Iba1, respectively. Both of these cell populations express purinergic receptor P2X7 which is likely involved in mechanisms connected with proinflammatory cytokine (IL-1b) release and cell death.

Experimental autoimmune encephalomyelitis (EAE) is a commonly used model of MS. EAE is evoked by injection of vaccine containing complete Freund's adjuvant, homogenate of guinea pigs' spinal cord and inactivated *Mycobacterium tuberculosis*. Animals, female Lewis rats, develop neurological symptoms similar to those observed in humans during MS.

The aim of this study was to investigate whether blocking of P2X7R by its selective antagonist BBG (Brilliant Blue G) influences the expression of astroglia and microglia specific proteins during the course of EAE.

In the inductive phase of EAE (4th day post immunization, 4 dpi) we observed an early activation of both astroglia and microglia, which was connected with overexpression of GFAP and Iba1, respectively. Overexpression of markers was also present at the peak of the disease (10 dpi). To elucidate the involvement of P2X7R in the mechanisms of glia activation during EAE, we examined brain tissue of EAE rats treated with BBG (50 mg/kg of body weight) starting from day 0 pi to day 6 pi. The immunofluorescent staining of brain specimens at 4 and 10 dpi, revealed decreased astroglia activation.

The activation of astroglia measured by fluorescence intensity showed that the immunoreactivity of GFAP in EAE group was 7-fold higher than that in EAE + BBG group at 4 dpi. At the peak of the disease the immune reactivity was 2-fold higher in EAE group than in EAE + BBG group. These results were confirmed by Western Blot analysis and qPCR.

Microglia activation in EAE group started at 4 dpi and proceed to the peak of the disease. In EAE + BBG group activated microglial cells were not observed before the peak of the disease. Intensity of the immunoreactivity in EAE group was 2-fold higher than in EAE + BBG group at 4 dpi. These results were also confirmed by Western Blot analysis and qPCR.

We may conclude that administration of BBG during the course of EAE results in the decreased reactivity of glial cells pointing out that P2X7R is actively involved in the mechanisms of cell activation during the disease.

Supported by grant no: 2012/05/N/NZ4/02191.

|A25|

Paracrine effect on energy metabolism of cholinergic SN56 cells in cytotoxic conditions

Gul-Hinc S., Ronowska A., Bielarczyk H., Szutowicz A.

Department of Laboratory Medicine, Medical University of Gdansk, Gdansk, Poland

Microglial cells, through the proinflammatory mediators are key elements initiating inflammatory reactions in the brain. They are linked with pathomechanism of Alzheimer's and other neurodegenerative diseases. Prolonged study of microglial cells have shown their neuroprotective, as well as neurotoxic effects on cholinergic neurons. The aim of this work was to investigate modifying effects of microglia on cholinergic neuronal SN56 cells subjected to common neurotoxic signals. One of this factors is zinc. Chronic exposure of differentiated (DC) SN56 cells to 0.150 mM Zn caused inhibition of pyruvate dehydrogenase (PDH) and aconitase activity (50%), decrease of acetyl-CoA level (30%), as well as loss of cell viability (60%). This results are consistent with our earlier findings (Ronowska *et al.* 2007). To investigate the paracrine effect of microglial cells on cholinergic SN56 we used conditioned medium of microglial cells. We reveal that conditioned medium of microglial cells has no effect on the survival of SN56 cholinergic cells, as well as activities of studied enzymes. Despite of that, conditioned medium of microglial cells partially prevented Zn-induced loss of DC viability to 20%, as well as PDH and aconitase activity, and acetyl-CoA level. These data demonstrate that neuroprotective effects of conditioned medium of microglial cells in Zn-challenged cholinergic neurons were linked with alterations in their acetyl-CoA metabolism. To clarify the molecular mechanism of this paracrine interactions more research is needed.

Supported by projects: GUMed ST-57, MN-15.

|A26|

Lowered expression of gliosis-associated markers in aging rats fed a high-fat diet

Gzielo-Jurek K.¹, Setkowicz Z.¹, Kielbinski M.¹, Ploszaj J.¹, Janeczko K.¹, Gazdzinski S.²

¹Department of Neuroanatomy, Jagiellonian University, Cracow, Poland

²Military Institute of Aviation Medicine, Warsaw, Poland

Introduction: Obesity is strongly associated with insulin resistance and type 2 diabetes, hypertension, dyslipidemia. Together with abdominal obesity, these symptoms are recognized as a condition called metabolic syndrome (MS). MS is recognized as one of the main causes of avoidable mortality and morbidity in developed countries. In Poland it affects up to 25% of the population. In addition to the cardiovascular risks associated with MS, it is being increasingly recognized as a potential source of neurological disease, especially in aging patients. Obesity or MS are associated with loss of brain-derived neurotrophic factor (BDNF) signaling, oxidative stress, blood-brain barrier dysfunction and chronic inflammation. Here, we used rats fed a high-fat diet (HFD), consisting of 40% fat, 40% sugar and 10% fiber. The animals were maintained on either HFD or normal chow for a year, to simulate obesity-associated effects in the aging brain. We then studied glial activation by immunohistochemical (IHC) staining for astrocytic marker GFAP and microglial protein Iba-1.

Material and methods: Male Wistar rats were assigned randomly to HFD or normal diet (ND) groups at age of 45-50 days, then placed in single cages with constant access to the type of food assigned. The experimental animals were then maintained on their respective diets for a year, followed by sacrifice and perfusion fixation with 4% formalin in PBS. Brains were processed into 30 μ m slices with a vibratome, and IHC for GFAP and Iba-1 were performed. The resulting slides were then photographed under a 4x objective. Panoramic images of whole stained hippocampi were assembled with Microsoft ICE. Next, an unsupervised local thresholding algorithm based on the Bernsen method, implemented in ImageJ, was used to measure the immunoreactive area fraction in the hippocampal formation. The resulting data were compared by Student's *t*-test.

Results: Somewhat surprisingly, we have found a significant ($p < 0.005$) reduction in immunopositive area fraction for both markers in animals from the HFD group. This is in accordance with the results of behavioral and magnetic resonance studies in the same population of animals, performed recently by our research group. In these studies, animals fed a HFD for 12 months exhibited better

spatial learning in the radial arm maze paradigm in comparison to controls. Similarly, the levels of *N*-acetylo-aspartate, loss of which is generally associated with aging-related changes and dementia, was higher in the HFD group. Further morphometric studies of astrocytes and microglial cells are underway, aimed at elucidating the nature of the effects of HFD in aging rats.

This work was supported by NCN 2011/03/B/NZ4/03771.

[A27]

Neuroprotective effects of undecylenic acid (UDA) against hydrogen peroxide-, staurosporine- and doxorubicin-induced cell damage in human neuroblastoma SH-SY5Y cells

Jantas D.¹, Piotrowski M.², Lason W.¹

¹Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Jerzy Haber Institute of Catalysis and Surface Chemistry, Polish Academy of Sciences, Cracow, Poland

Undecylenic acid (UDA), a naturally occurring an 11-carbon unsaturated fatty acid has been used from several years as an economical antifungal agent and nutritional supplement. Recently, an additional clinical utility of UDA has been proposed as a promising component of nanocarriers for the oral delivery of proteins and peptides. Moreover, the potential usefulness of UDA as neuroprotective drug has been suggested, based on the ability of this agent to inhibit μ -calpain activity. In order to provide more evidence on neuroprotective potential of UDA, we tested this compound against oxidative stress (hydrogen peroxide)- and pro-apoptotic factors (staurosporine and doxorubicin)-evoked cell damage in human neuroblastoma SH-SY5Y cells. The data showed that UDA (1-40 μ M) decreased the hydrogen peroxide-induced cell damage as confirmed by MTT reduction and LDH release assays. Moreover, the protection mediated by UDA in that model was connected with attenuation of hydrogen peroxide-mediated necrotic (PI staining) and apoptotic (caspase-3 activation) changes. Using Western blot analysis we demonstrated that UDA attenuated the hydrogen peroxide-stimulated increase in 145 kDa and 120 kDa of α , β spectrin II products, specifically cleaved by calpains and caspases, respectively. Next, we showed that UDA partially protected SH-SY5Y cells against the staurosporine- and doxorubicin-evoked cell death, however this effect was not connected with its

influence on caspase-3 activity. Further studies at the St model of cell injury revealed that UDA decreased the St-induced mitochondria-cytosolic translocation of AIF (Apoptosis Inducing Factor) which is responsible for caspase-3 independent execution of apoptotic cell death. Finally, we demonstrated that inhibitor of PI3-K/Akt (LY294002) but not MAPK/ERK1/2 (U0126) pathway blocked the protection mediated by UDA in all tested models of neuronal cell injury. The data showed neuroprotective effects of UDA in various models of SH-SY5Y cell injury with the involvement of activation of pro-survival PI3-K/Akt pathway in that effect. Moreover, both caspase-3-dependent and -independent mechanisms could be involved in the mechanism of UDA-mediated protection in SH-SY5Y cells.

[A28]

Altered arginine metabolism in cells transfected with human wild type amyloid precursor protein (APP)

Jesko H.¹, Wilkaniec A.¹, Cieslik M.¹, Hilgier W.², Gassowska M.¹, Strosznajder J.B.¹, Lukiw W.J.³, Adamczyk A.¹

¹Department of Cellular Signalling, Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Department of Neurotoxicology, Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

³Neuroscience Center of Excellence and Departments of Neurology and Ophthalmology, Louisiana State University Health Sciences Center, New Orleans, USA

Amyloid-beta ($A\beta$) peptide, the product of $A\beta$ precursor protein (APP) proteolysis plays a central role in Alzheimer's disease (AD). Recent evidence suggests that alteration of urea cycle enzymes might be also involved in AD pathology. Our previous results showed that $A\beta$ is associated with the induction of oxidative/nitrosative stress and stimulation of arginine metabolism by Nitric Oxide (NO) synthase (NOS). Although urea cycle is associated with the citrulline-NO pathway, the impact of $A\beta$ on arginine degradation by its enzymes is unknown. Therefore, in the present study we investigated the expression of genes involved in the urea cycle in rat pheochromocytoma (PC12) cells overexpressing human wild type APP (*APPwt*). We compared gene expression levels of neuronal, endothelial and inducible *NOS* isoforms (*NNOS*, *ENOS*, *INOS*), urea cycle enzymes and corresponding amino acids in control PC12 and *APPwt* cells. Expression of each *NOS* isoform and urea cycle enzymes: Arginase-1 (*ARG1*) and Arginase-2 (*ARG2*),

Ornithine carbamoyltransferase (*OTC*), Argininosuccinate synthase (*ASS*), Argininosuccinate Lyase (*ASL*) and Ornithine Decarboxylase 1 (*ODC1*) was measured using quantitative Real-Time RT-PCR. Determination of amino acids arginine, ornithine and citrulline were carried out by high performance liquid chromatography (HPLC). Moreover, we investigated the changes in miRNAs responsible for modulation of gene expression of urea cycle in AD brains. Our data demonstrated that the overexpression of human wild-type *APP* in PC12 cells resulted in elevation of intracellular and secreted A β . The expression of urea cycle genes is disturbed both in *APPwt*-transfected PC12 cells and in AD brains. The expression of *ARG1* and *ARG2* enzymes which are responsible for the lysis of arginine to ornithine and urea was significantly reduced in *APPwt* cells by about three- and two-fold, respectively. *ODC1*, which catalyzes the decarboxylation of ornithine to form putrescine was reduced in *APPwt* cells by 36%. The expression of *ASS* that catalyzes the condensation of citrulline and aspartate to form argininosuccinate was also down-regulated in *APPwt* cells by 24% without changes in *ASL* gene expression. Our results indicated that arginine, the substrate for both NO synthesis and urea cycle, was unchanged in *APPwt* cells comparing to control. However, citrulline and ornithine were found to be significantly elevated in *APPwt* cells by 22% and 14%, respectively. We then investigated how the expression of *NOS* isoforms changed in each cell lines. Real-time PCR analysis revealed significant elevation of *NNOS* and *ENOS* mRNA by about six-fold and 50%, respectively in *APPwt* cells comparing to control, whereas *iNOS* was not detectable. In addition, we have found elevated level of miRNA-9 and -128a in AD brains that might modulate the expression of *ASS* and *NOS*, respectively. These results indicated that A β peptide is involved in the urea cycle modification in PC12 cells. We conclude that this peptide might have important role in AD pathomechanism via alteration of the arginine metabolism by *NOS* and urea cycle enzymes.

Supported from IMDIK statutory theme 17.

[A29]

TLR2 and TLR4 antagonists act like analgesics and potentiate opioids effectiveness under neuropathic pain

Jurga A.M., Rojewska E., Kolosowska N., Popiolek-Barczyk K., Makuch W., Przewlocka B., Mika J.

Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Neuropathic pain is one of a current major clinical problems since its pathogenesis is still not known. It may appear as a consequence of nervous system damage or as a co-symptom of many diseases. There is rising amount of hypotheses indicating toll like receptors (TLRs) to play a role in neuropathic pain development and that TLR-dependent glial activation reduce opioids analgesia. The goal of our experiments was to evaluate the contribution of TLR2 and TLR4 in modulation and maintenance of neuropathic pain and their ability to amplify the morphine and buprenorphine effectiveness in neuropathic pain model. Experiments were performed on Wistar male rats and consisted of biochemical analysis (Western blot, qPCR) of the lumbar ipsilateral spinal cord tissue and behavioral tests (von Frey's for tactile allodynia, cold plate for thermal hyperalgesia). Neuropathic pain was developed using Bennett's model (chronic constriction injury (CCI) to the sciatic nerve). TLR antagonists were administered once per day for 7 days after CCI: LPS-RS (TLR4&TLR2 antagonist) and LPS-RS Ultrapure (TLR4 specific antagonist) – allodynia and hyperalgesia were measured 1-3 hours after drug administration on day 2 and 7 after CCI. Opioids (morphine or buprenorphine) were injected once on day 7, after CCI and 60 min after morning antagonists administration. All experiments were carried out according to the recommendations of International Association for the Study of Pain (Zimmermann *et al.* 1983) and the NIH Guide for Care and Use of Laboratory Animals and were approved by the Local Bioethics Committee (Cracow, Poland). Biochemical analysis shows on day 7 after CCI the strong up-regulation of TLR2 and TLR4 mRNA and protein levels as measured on the ipsilateral dorsal (L4-L6) spinal cord level as compare to the naïve rats. Repeated intrathecal administration of LPS-RS and LPS-RS Ultrapure attenuated allodynia and hyperalgesia as measured by von Frey and cold plate tests. Additionally, we examined if chronic administration of TLR receptor antagonists influence opioids analgesia on 7th day. Our preliminary data suggest that both TLR2 and TLR4 indeed play significant role in neuropathic

pain development, since their levels are elevated in neuropathic pain conditions.

Grant OPUS 2011/03/B/NZ4/00042.

|A30|

Contribution of CCK_A receptors to cardiovascular and respiratory effects of cholecystokinin in anesthetized rats

Kaczynska K., Szereda-Przestaszewska M.

Laboratory of Respiratory Reflexes, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Cholecystokinin (CCK) is a neurotransmitter, neuro-modulator and a gastrointestinal hormone present in the central and peripheral nerve terminals and in endocrine cells. Within the brain CCK is engaged in feeding/satiety, pain/analgesia, panic/anxiety and neurological disorders.

In the nodose ganglia, abounded with cell bodies of primary afferent fibers transmitting sensory information from the cardiorespiratory organs, mRNA for CCK_A and CCK_B receptors is expressed. Nucleus of the solitary tract, a key brainstem area for the integration of autonomic functions, receiving projections from afferents of the nodose ganglia, displays immunoreactivity for CCK and its receptors.

Therefore, we investigated a share of vagal input at infra- and supra-nodosal level and the contribution of CCK_A and CCK_B receptors to the cardiorespiratory responses produced by an intravenous injection of cholecystokinin in anesthetized rats.

Cholecystokinin administered intravenously at a dose of 50 µg/kg induced short-lived decline in tidal volume and respiratory rate resulting in depression of minute ventilation. Mid cervical vagotomy had no effect on CCK-evoked ventilatory changes, whereas supranodosal denervation abolished slowing down of the breathing. Cardiovascular response to the CCK challenge was characterized by a transient decrease, followed by an augmentation in the mean blood pressure (MAP) in the intact animals. Following vagotomy performed at both levels, the declining phase of MAP was abrogated. Blood pressure changes were associated with decreased heart rate present in all neural states. All cardiovascular and respiratory effects were antagonized by pretreatment with devazepide – CCK_A receptors' antagonist, whereas CI988 – an antagonist of CCK_B receptors, was ineffective.

In conclusion, our results indicate that CCK modulates slowing down of respiratory rhythm via CCK_A receptors located in the nodose ganglia, and depresses tidal volume via the central CCK_A dependent mechanism. CCK-evoked declining in blood pressure may be due to the activation of vagal afferents, whereas pressor responses seem to be possibly mediated by the activation of CCK_A receptors in the vasculature and/or central nervous system. Bradycardia was probably induced by the direct action of CCK on heart pacemaker cells.

|A31|

Involvement of 5-HT_{1A} receptors in mirtazapine and risperidone co-treatment effect on NA, DA and 5-HT extracellular level in the rat frontal cortex

Kaminska K.¹, Golembiowska K.¹, Rogoz Z.^{1,2}

¹Department of Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²The Podhale State Higher Vocational School, Nowy Targ, Poland

Major depressive disorder (MDD) is a chronic mental illness with a life time prevalence of 5-12% in adult men and 9-26% of adult women. Although initial antidepressant (AD) therapy significantly reduces symptoms of depression in many patients, only 50-60% of persons with MDD respond to the treatment. Moreover, ca. 30-40% of MDD patients never achieve symptom resolution by means of a standard AD therapy. The problem of AD-resistant depression has been the subject of a number of thorough studies, with no apparent therapeutic success, though. Hence, there is a strong need for an alternative antidepressant treatment. Several clinical reports have postulated a beneficial effect of an additional low dose risperidone to ongoing treatment with ADs. Like other atypical antipsychotic drugs, risperidone is known to produce minimal extrapyramidal side-effects compared to classical antipsychotics. It has been proposed that risperidone in lower doses acts mainly by blocking 5-HT_{2A} serotonin receptors, while at higher doses it mostly blocks D₂ dopamine receptors. Moreover, it has been shown that mirtazapine, a noradrenergic and specific serotonergic AD is an agonist of 5-HT_{1A} receptor and an antagonist of the central α₂-auto- and hetero-adrenoreceptors. It also blocks 5-HT₂ and 5-HT₃ receptors and displays a very low affinity for DA receptors and a high affinity for histamine H₁ ones.

The aim of our study was to understand the mechanism of clinical efficacy of combination an antidepressant and an antipsychotic in drug-resistant depression.

We studied the effect of an AD (mirtazapine) and risperidone (an atypical antipsychotic) given separately or jointly, on the extracellular levels of dopamine (DA), serotonin (5-HT) and noradrenaline (NA) in rat frontal cortex. The animals were given single intraperitoneal injections of risperidone (1 mg/kg) and mirtazapine (10 and 20 mg/kg). The release of monoamines was investigated using a microdialysis in freely moving animals, and assayed by HPLC with coulochemical detection.

Mirtazapine dose-dependently increased the cortical extracellular levels of DA, 5-HT and NA. Similarly, risperidone increased the cortical extracellular levels of the monoamines studied. The combination of mirtazapine (20 mg/kg), or both doses of mirtazapine (10 and 20 mg/kg), with risperidone produced significantly stronger release of DA and NA, respectively (but not of 5-HT release) compared to the effect of administration of a single drug. The increase in the DA (but not NA) release induced by mirtazapine plus risperidone was partly blocked by the selective 5-HT_{1A} antagonist WAY 100635 (0.2 mg/kg).

The obtained results suggest that the ability of risperidone to augment the therapeutic efficacy of mirtazapine in treatment-resistant depression may be due, at least in part, to the potentiation of mirtazapine-induced increases in cortical DA and NE. Furthermore, among other mechanisms, 5-HT_{1A} receptors may play some role in this action.

This study was financially supported by statutory funds of the Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland.

The most prevalent of these is the focal cortical dysplasia (FCD), a cryptogenic malformation of brain development resulting in loss of cortical lamination, formation of cortical and subcortical ectopias, disrupted white matter and other abnormalities. Animal models aimed at simulation of these phenomena rely on controlled physical or pharmacological disruption of neurodevelopment. The onset of this disruption, i.e. the day of gestation when the stimulus is applied, determines the pattern of changes and their severity.

Here, we used prenatal gamma irradiation at embryonic days E13, E15, E17 or E19 in comparison with untreated controls, to elucidate the effects of irradiation-induced dysplasia on the volume of different brain structures. We have previously shown that the presence of dysplastic changes interacts with epileptogenic factors affecting seizure susceptibility and the expression of glial and neuronal markers. We hypothesize that after pilocarpine injection – a widely studied model of epileptic seizures – different patterns of dysplasia would differentially affect the resulting changes in brain volume.

Animals irradiated with a dose of 1 Gy at E13 ($N = 11$), E15 ($N = 9$), E17 ($N = 9$) or E19 ($N = 8$), as well as normal controls ($N = 10$) were given pilocarpine (250 mg/kg) at P60. After 72 h the animals were sacrificed, perfused transcardially with 10% formalin and the brains were removed. T2-weighted images taken with a 9.4 T magnetic resonance scanner were then obtained. Using a computer-assisted Cavalieri method, the volumes of the entire brain, hippocampus and brainstem structures (midbrain, pons and medulla oblongata) were estimated. Volumes were compared between groups to show differentially affected regions. Also, within each group, comparisons were made with age-matched untreated animals: irradiated at E13 ($N = 12$), E15 ($N = 12$), E17 ($N = 12$) or E19 ($N = 12$) and naïve controls ($N = 14$). ANOVA with HSD corrected for unequal N or Student's t -tests were used.

In both pilocarpine-treated and untreated animals, brain volume was reduced in a time-dependent manner, with animals irradiated earlier showing a more profound loss in brain volume. In animals after pilocarpine injection, hippocampal volume normalized to whole brain volume follows an opposite trend, with greatest effects in animals irradiated at E17 or E19. In addition, pair-wise comparison of naïve vs. pilocarpine treated rats revealed an attenuated effect of seizures on brain and hippocampal volume in animals irradiated at E19. This is consistent with the notion that dysplasia modifies the effects of pro-convulsive stimuli, with effects depending on the onset of disturbed neurogenesis.

Supported by National Center for Science (NCN) grants DEC-2012/05/N/NZ2/00641 and 2011/01/B/NZ4/00586.

[A32]

Brain swelling after pilocarpine-induced epilepsy: effect of cerebral malformation in rats

Kielbinski M.¹, Gzielo-Jurek K.¹, Setkowicz Z.¹, Banasik T.², Ploszaj J.¹, Weglarz W.², Janeczko K.¹

¹Department of Neuroanatomy, Jagiellonian University, Cracow

²Laboratory of In Vivo Imaging, Institute of Nuclear Physics, Polish Academy of Sciences, Cracow

Brain edema and gliosis accompany neuronal loss as the main effects of epileptic seizures, both in patients and in animal models, such as pilocarpine administration.

Cerebral malformations in humans are strongly associated with intractable epilepsy, especially in pediatric patients.

|A33|

Neurochemical changes linked to antipsychotic-like phenotype of mGlu4 knockout mice

Kleczek N.^{1,2}, Slawinska A.¹, Marciniak M.¹, Wieronska J.¹, Pilc A.¹

¹Department of Neurobiology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Department of Neurophysiology and Chronobiology, Institute of Zoology, Jagiellonian University, Cracow, Poland

Introduction: The metabotropic glutamate 4 (mGlu4) receptor has been proposed as an important target in the search of novel antipsychotic drugs. Several orthosteric and allosteric activators of that receptor were shown to possess antipsychotic effects in animal models of schizophrenia. In the present study we investigated the role of mGlu4 receptor in the pathophysiology of schizophrenia. We used mGlu4 knockout mice to see, if the lack of the receptor results in any behavioral changes in animal models of psychosis. Moreover, we investigated a possible neurochemical changes in the selected proteins known as the markers of psychosis, e.g. 5-HT1A receptor, GAD65 and GAD67 enzymes. The level of proteins was investigated in the stratum, hippocampus and prefrontal cortex.

Material and methods: Adult C57BL/6J mice with mGlu4 receptor knockout (mGlu4 KO) and wild type (WT) mice were used in this study. Mice with mGlu4 receptor knockout were generated in Institute of Pharmacology, Polish Academy of Sciences in Krakow. Animals were maintained under standard laboratory condition on 12-hour light/12-hour dark cycle at a room temperature of 19-22°C with free access to food and water. To investigate phenotype of mGlu4 KO we used behavioral models of selected symptoms of schizophrenia, such as DOI-induced head twitches and one-trial recognition task. Animals were treated with GABAB antagonist CGP 55845 (10 mg/kg, *i.p.*) or 5-HT1A antagonist, WAY 100635 (0.2 mg/kg, *s.c.*), 30 and 45 min before test, respectively. The relative mRNA expression of 5-HT1a receptor, GAD-65 and GAD-67 enzymes was measured with the use of quantitative Real-Time Reverse Transcriptase PCR.

Results: Knockout mice had a reduced response to psychostimulant in DOI-head twitches reflected by decreased number of episodes, as well as less sensitivity to MK-801-induced disruption in one-trial recognition task. The behavioral effect observed in mGlu4 knockouts in DOI-induced head twitches was reversed after administration of CGP 55845 and WAY 100635. In the mRNA studies the decreased level of 5-HT1A mRNA in the frontal cor-

tex was observed, but not in the hippocampus or striatum. In contrast, the increases in GAD-65 and GAD-67 mRNA levels in frontal cortex were noticed. The level of GAD65 mRNA was also increased in the hippocampus, but neither of GADs were changed in the stratum.

Conclusions: The present study shows an antipsychotic-like phenotype of mGlu4 mutant mice which can be reversed by GABAB antagonist (CGP55845) or 5-HT1A antagonist (WAY100635). Furthermore, the changes in the mRNA level of selected proteins known to be important in schizophrenia are relevant to the phenotype observed in the behavioral studies and stays in line with human studies.

|A34|

Sildenafil, an inhibitor of phosphodiesterase 5 (PDE5), affects asymmetric behavior and monoamine metabolism in unilaterally 6-OHDA-lesioned rats treated chronically with L-DOPA

Kmiec I., Lenda T., Kaminska K., Czarnecka A., Lorenc-Koci E.

Department of Neuro-Psychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

The dopamine replacement therapy with L-DOPA still remains the most widely used treatment for Parkinson's disease (PD). However, over time, the effectiveness of the drug decreases, while motor complications, such as dyskinesia, develop. Therefore, the search for a new therapeutic approach that can provide benefit over L-DOPA therapy is still ongoing. Recently, attention of researchers has focused on the nitric oxide (NO) – soluble guanylyl cyclase (sGC) – cyclic GMP signaling pathways in the striatum as a new target for the treatment of PD.

The aim of the present study was to examine the effects of chronic *i.p.* treatment with the phosphodiesterase 5 (PDE5) inhibitor, sildenafil in combination with the widely used antiparkinsonian drug, L-DOPA on asymmetric behavior and monoamine metabolism in the striatum (STR) and substantia nigra (SN) of unilaterally 6-OHDA-lesioned rats. The experiments were performed on male Wistar rats injected unilaterally with 6-OHDA (8 µg/4 µl) into the medial forebrain bundle. Two weeks later, rats exhibiting more than 100 contralateral rotations per 1 h in the apomorphine (0.25 mg/kg) test, indicative on an extensive loss of the nigrostriatal neurons, were treated chronically

with sildenafil (2 mg/kg) and L-DOPA (12.5 mg/kg), alone or in combination, once daily for successive 14 days. Rotational behavior was also recorded immediately after the first and penultimate doses of the examined compounds. Rats were killed 1 h after the last doses of these drugs, their STR and SN were dissected on an ice-chilled plate and stored at -80°C until further procedures were applied. The tissue concentrations of dopamine (DA) and its metabolites were determined using an HPLC method.

Our results showed that sildenafil did not evoke contralateral rotations after acute or chronic treatment. In contrast, L-DOPA produced distinct contralateral rotations already visible 10 min after its first dose and lasting almost 100 min. Moreover, after chronic L-DOPA treatment their number was higher than after its acute injection. However, after combined acute administration of sildenafil and L-DOPA, the number of contralateral rotations was markedly higher while after chronic treatment lower than in the group receiving L-DOPA alone. Regarding DA concentration, chronic sildenafil administration affected the tissue DA level neither in the ipsi- nor contralateral STR and SN. Chronic treatment with L-DOPA alone significantly increased the tissue DA content only in the ipsi- and contralateral SN when compared to the corresponding side of the 6-OHDA-lesioned groups receiving sildenafil. However, chronic combined treatment with these compounds produced markedly higher increases in the tissue DA contents in the ipsilateral STR and SN than in the group of rats receiving L-DOPA alone.

The obtained results suggest that administration of a low dose of the PDE5 inhibitor, sildenafil jointly with L-DOPA modulates rotational behavior and tissue DA concentrations in the nigrostriatal system of unilaterally 6-OHDA-lesioned rats.

This work was supported by a grant from the National Science Centre of Poland, based on decision No. UMO-2011/01/B/NZ4/01581.

[A35]

The evidence for potential of human bone marrow mesenchymal stem cells to cross the blood-barrier after their intra-arterial transplantation

Koniusz S.¹, Andrzejewska A.¹, Janowski M.^{1,2}, Lukomska B.¹

¹NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Russel H. Morgan Department of Radiology and Radiological Science, Division of MR Research, The Johns Hopkins University School of Medicine, Baltimore, USA

Introduction: Mesenchymal stem cells (MSC) have recently emerged as promising candidates for cell-based therapy in neurological disorders. Based on differentiation potential of MSC initial studies focused on the regenerative capacity of these cells. However, over time it became clear that MSC mediated their positive effect rather through the production of trophic factors. The mechanisms of action used by MSC i.e. immunomodulation, cell protection, pre-reparation and stimulation of angiogenesis have been elucidated *in vitro* but the data from pre-clinical studies have been unclear. Furthermore, the potential for minimally invasive delivery of MSC therapy in term of clinical transplantation is of particular interest. Critical questions include MSC homing to ischemic tissues from peripheral blood and the efficiency of this process. The aim of the study was to analyze the presence, distribution and quantity of human bone marrow mesenchymal stem cells (hBM-MSCs) after their intra-arterial delivery in focal cerebral ischemic rats.

Material and methods: The experiments were performed in adult male Wistar rats. To induce rat brain injury ouabain – Na/K ATPase pump inhibitor have been chosen which brings metabolic and structural cell changes that mimic ischemic deep-brain, stroke-like lesion. For the surgical procedure rats were anesthetized and immobilized in a stereotactic apparatus, then a small burr hole was drilled in the cranium over the right hemisphere, the needle was lowered into the right striatum and $1\ \mu\text{l}/50\ \text{nmol}$ ouabain was injected. Two days after the ischemic insult, 5×10^5 hBM-MSC (Lonza) tagged with iron nanoparticles conjugated with rhodamine (Molday, BioPAL) were transplanted into the right internal carotid artery. After 1, 3, 7, 14, 21 and 28 days post-transplantation rats were sacrificed, the brains were removed and quickly frozen. Then immunocytochemical analysis of hBM-MSC present in the brain, cervical lymph nodes and spleen was performed using anti-human (CD44, STEM121 and Ku80) antibodies.

Results: The preliminary results of ongoing studies revealed the presence of transplanted cells in the rat brain. hBM-MSC injected intra-arterially were visible in the right hemisphere in the area between cortex and stratum of ouabain-injured rat brain near the ischemic lesion. The fluorescent signal of Molday particles and positive staining for human antigens were observed until 28 days of observation after hBM-MSC transplantation.

Conclusions: Intra-arterial delivery of hBM-MSC promotes their migration to ischemic sites of the brain where they may release paracrine factors for the local or systemic effect. The further studies of hBM-MSC presence in the lymphatic tissues as well as neurotrophic, neuroprotective and immunomodulating function of transplanted cells in focal cerebral ischemic disorders are in progress.

This work was supported by KNOW 06 project: "The role of bone marrow mesenchymal stem cells in CNS repair of brain ischemia disorders".

[A36]

Influence of dopamine agonists on the harmaline-induced tremor measured by force plate actimeters

Kosmowska B., Glowacka U., Wardas J., Ossowska K.

Department of Neuro-Psychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Harmaline is a β -carboline derivative which is commonly used for modelling one of the most frequent movement disorder, essential tremor (ET), in animals and screening potential tremorolytic drugs. Oscillation frequency of the harmaline-induced kinetic/postural tremor is 10-12 Hz in rats. It is generally accepted, that the primary cause of tremor induced by this compound is an abnormal synchronous activation of the olivo-cerebellar glutamatergic climbing fibres. This activation leads to increased glutamate release in the cerebellum and induction of complex spike activity of Purkinje cells of the cerebellar cortex. However, mechanisms underlying harmaline-induced tremor seem to involve also other neurotransmitter systems.

The aim of the present study was to examine the influence of the non-selective agonists of dopamine receptors: apomorphine (D_1 - D_5) and pramipexole (D_2 , D_3) on the harmaline-induced tremor in rats. Propranolol (the first-line treatment of ET) was used as a reference compound. Since tremor is difficult to quantify and distinguish from other

motor abnormalities in animals, this behaviour was measured objectively by fully automated Force Plate Actimeters. To quantify the intensity of the tremor, the following parameters were measured: AP1 (power in the 0-8 Hz band), AP2 (power in the 9-15 Hz band) and tremor index (difference in power between the AP2 and AP1). In order to examine the locomotor activity, the total distance travelled by an animal was also measured. In some groups of animals the focused stereotypy or AP2/distance ratio was analysed additionally.

The generalised tremor induced by harmaline administered at a dose of 15 mg/kg *i.p.* was related to an increase in the AP2 parameter and the tremor index. Propranolol injected at a dose of 20 mg/kg *i.p.* reversed both the above effects of harmaline. Apomorphine administered at doses of 0.5 and 1 mg/kg *s.c.* further enhanced AP2 and at the lower dose also the tremor index elevated by harmaline. The increase in the AP2 was stronger than an enhancement of locomotor activity induced by apomorphine in the harmaline-treated animals. Pramipexole (0.1, 0.3 and 1.0 mg/kg *s.c.*) had no effect on tremor index, but at dose of 0.1 mg/kg significantly reduced AP2 parameter.

The present study suggests that apomorphine enhances the tremor induced by harmaline, and this effect is at least partly independent of hyperactivity induced by this dopamine agonist. In addition, the results after pramipexole administration revealed the contribution of D_2/D_3 receptors in regulation of tremor in harmaline model of ET.

The study was supported by Statutory Funds of the Department of Neuro-Psychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland.

[A37]

Effects of different stressors on amygdalar monoamines

Krotewicz M.¹, Bajer M.¹, Krotewicz M.K.²

¹Department of Neurobiology, Faculty of Biology and Environment Protection, University of Lodz, Lodz, Poland

²Department of General and Colorectal Surgery, Medical University of Lodz, Lodz, Poland

The amygdala has attracted interest as concerns its implication in emotion – behaviour interactions. The amygdala complex receives innervation from the noradrenergic, serotonergic and dopaminergic pathways originating from the brain stem. The monoaminergic systems participate in the regulation of emotional behaviour. On the other hand, it was shown that different stressors have an impact

on monoaminergic neurotransmission. In order to compare the neurochemical effects of different stressors we exposed rats to the most popular anxiety tests. The anxiety tests are widely used to study behaviour of rats in the "stress" – inducing situation. In these different ethological procedures, animals are exposed to unknown environment differing in size, shape, lighting, time of exposure etc. The most popular anxiety tests for the study of exploration and emotionality in rodents have become the open field (OF) and the elevated plus-maze (EPM). The influence of various pharmacological agents on behaviour and physiological functions have been investigated in these tests. It should also be noted that the anxiety tests are used interchangeably in the investigation of anxiety mechanisms. In the present study we undertook a simultaneous examination of noradrenergic, serotonergic and dopaminergic systems activity in the amygdala of rats exposed to one of the three anxiety tests: open field (OF), elevated plus maze (EPM) and transitions test (TT). The experiments were performed at the same time on adult male Wistar rats, housed in groups under controlled conditions. The rats were individually subjected to one of the anxiety tests (OF, EPM or TT) and sacrificed by decapitation immediately after a test procedure. The concentration of noradrenaline, dopamine, serotonin and their metabolites were determined in the amygdala using high-performance liquid chromatography with electrochemical detection. The results of our study showed that stressogenic procedure of the used anxiety tests exerted influence on indices of the monoaminergic systems activity in the amygdala. Especially, the exposure of rats to the used tests (OF, EPM and TT) was accompanied by increase in serotonin turnover in the amygdala. The obtained data revealed that the exposition of rats on the stressogenic situation deeply increase serotonin metabolism in the amygdala region. It is likely that the elevation of serotonin turnover in this emotional brain region sensitize the serotonergic system on influence of serotonergic ligands. The presented results confirm our previous observation that the transition test (TT) is a good tool to study behavioural effects of anxiolytic agents acting on serotonergic system.

[A38]

Opioid receptors expression and PGP 9.5-positive epidermal nerve fiber density in psoriasis – relationship with itch

Kupczyk P.¹, Reich A.², Wysokinska E.³, Gajda M.⁴, Holysz M.⁵, Hwang T.⁶, Kobuszewska A.², Nowakowska B.^{1,7}, Drukala J.⁸, Szepietowski J.C.²

¹Laboratory of Immunogenetic and Tissues Immunology, Department of Clinical Immunology, Ludwik Hirszfild Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, Wroclaw, Poland

²Department of Dermatology, Venereology and Allergology, Wroclaw Medical University, Wroclaw, Poland

³Laboratory of Molecular Immunobiology of Cancer, Department of Experimental Oncology, Ludwik Hirszfild Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, Wroclaw, Poland

⁴Department of Histology, Jagiellonian University Medical College, Cracow, Poland

⁵Department of Biochemistry and Molecular Biology, Medical University of Poznan, Poznan, Poland

⁶Department of Pharmacology, Medical University of Minnesota, MN, USA

⁷Laboratory of Cellular and Tissues Engineering, Department of Cell Biology, Division of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Cracow, Poland

⁸Laboratory of Tissues Immunology, Medical Center of Ludwik Hirszfild Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, Wroclaw, Poland

Opioid receptors (OR): *OPRM1/MOR*, *OPRK1/KOR*, *OPRD1/DOR*, *OPRL1/ORL* are highly expressed in central nervous system (CNS) and to lesser degree outside the CNS in peripheral tissues like lungs, heart, vessel walls, immune cells and in the skin. OR in the skin participate in transmission of molecular signals from CNS and control keratinocyte differentiation, proliferation, and apoptosis. OR are also suggested as potential targets for peripheral pain and itch therapy.

However, molecular studies on itch in psoriasis gave inconclusive results. The most widely discussed theory mentioned downregulation of skin nervous system and imbalance of different classes of neurobiological pathways. OR theory suggested that *OPRM1/MOR* system is responsible for promotion, while *OPRK1/KOR* system for suppressing itch signals in the skin and CNS. To prove this suggestion, 20 psoriasis patients with and without itch underwent punch biopsies from non-lesional and lesional skin. The results of study of OR expression and epidermal nerve density were compared with the clinical parameters. Itch intensity have been assessed using VAS (Visual Analogue Scale), while severity of psoriasis with

PASI (Psoriasis Area and Severity Index). Control group consisted of skin samples obtained from 20 healthy volunteers. A multiplex real-time PCR with UPL probes were used to determine gene expression of *OPRM1/ACTB* and *OPRK1/ACTB* by relative quantification analysis. Protein expression evaluation was performed by indirect immunofluorescence on skin frozen sections using anti-MOR and anti-KOR antibodies labelled with secondary fluorochrome conjugated antibodies. Fluorescence intensity analysis was done using ImageJ.

OPRM1/MOR expression was similar in all groups, while *OPRK1* gene show decreased expression in lesional psoriatic skin compared to controls. A lower expression of *OPRK1* gene was noted in patients with itch compared to those without itch. KOR expression was not significantly different in lesional and non-lesional skin of patients without itch compared to control group. Lesional skin with itch showed significantly lower KOR expression compared to non-lesional one or control group. Epidermal innervations analysis show significantly higher PGP 9.5-positive nerve density in non-lesional and lesional skin compared to control and exhibited positive correlation with VAS and reverse with PASI.

Based on our results, we believe that imbalance of *OPRK1/KOR* and constant *OPRM1/MOR* expression with downregulation of cutaneous nervous system may be relevant neurobiological and molecular pathogenic factors in psoriatic itch.

compensatory processes could depend on increased energetic demands for surviving cells and intracellular adaptations in mitochondrial function and structure. Our aim was to investigate if functional compensation of motor deficit after selective degeneration of dopaminergic neurons would influence activity of mitochondrial Cx I and IV and cause adaptation in their SCx assembly.

Material and methods: We have prepared PD rat model of selective nigrostriatal dopaminergic system degeneration by intracerebral, stereotaxic injection of 3 µg/3 µl 6-hydroxydopamine into medial forebrain bundle. Animals showed decreased locomotor activity 3 days after the operation and this disability was functionally compensated after 4 weeks. Using solubilized crude mitochondrial membrane fraction from striatum we performed in-gel enzymatic activity measurements of complex I and IV, after blue-native electrophoresis.

Results: Cx I amount of SCxs and individual Cx I₁, as well as performance of SCxs, (but not I₁) decreased at both time-points after operation, to the same degree. At the same time specific activities were mostly not changed. There were no significant changes in Cx IV amounts but performance of Cx IV within SCxs was decreased – mostly after 4 weeks (I₁III₂IV₃, I₁III₂IV₂). Also, specific activity of SCxs was decreased only 4 weeks after toxin injection, correlating with functional compensation in behavior. No changes were detected in Cx IV individual or dimeric form.

Conclusions: The decrease in specific activity of Cx IV in form of SCxs indicated its role in compensatory mechanisms.

The study was supported by the Statutory Funds of the Institute of Pharmacology, PAS, Cracow, Poland and Technische Universität Darmstadt, Germany, DAAD scholarship to KK.

[A39]

Activity and supercomplexes assembly of mitochondrial complex I and IV in striatum during process of functional compensation for deficits in dopaminergic system in rat

Kuter K.¹, Ossowska K.¹, Dencher N.²

¹Department of Neuropsychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Physical Biochemistry, Department of Chemistry, Technische Universität Darmstadt, Darmstadt, Germany

Introduction: At the early stages of Parkinson’s disease (PD) motor degeneration of dopaminergic system is masked by potent compensatory mechanisms. Mitochondrial oxidative phosphorylation complexes (Cx) I, III and IV form higher order structures – supercomplexes (SCxs) that improve their function by channeling substrates, stabilizing protein assemblies and reducing ROS formation. Com-

[A40]

Gender differences in the neurochemical response of trigeminal ganglion neurons to peripheral inflammation in mice

Kuzawinska O., Lis K., Cudna A., Balkowiec-Iskra E.

Department of Experimental and Clinical Pharmacology, Medical University of Warsaw, Warsaw, Poland

The trigeminal system, with first-order neurons in the trigeminal ganglion (TG), provides sensory innervation to craniofacial tissues, many of which are associated with

chronic pain conditions, such as trigeminal neuralgias, migraines and temporomandibular joint (TMJ) disorders. Our previous studies showed that inflammatory reaction in the area of TG nociceptive endings (induced by tooth injury) affects neurochemical properties of TG ganglia, causing increase in concentration of both the neurotrophin brain-derived neurotrophic factor (BDNF) and well-established modulator of TG pain reaction – calcitonin gene-related peptide (CGRP). Moreover, our *in vitro* data showed that among three proinflammatory cytokines examined (IL-1, IL-6, TNF), only TNF was capable of influencing TG ganglia properties. In the present study, we have employed CFA-induced TG inflammation model to begin investigating the underlying molecular mechanisms of the inflammation-induced changes in trigeminal neurons properties.

A model of oro-facial inflammation was obtained by local injection of 30 µl of CFA to the mice left whisker pad. Controls were injected with the same volume of saline. Seven and 14 days after CFA injection mice were euthanized by spinal cord dislocation. Both TG ganglia were removed for quantitative PCR analysis. To study the effect of peripheral inflammatory reaction evoked by CFA administration on TG function we measured the expression of BDNF, CGRP and three proinflammatory cytokines – IL-1β, IL-6 and TNF, separately in the left and the right trigeminal ganglia, both in male and female mice, 7 and 14 days after induction of inflammation.

Our results indicate that response of TG ganglia to peripheral inflammatory reaction is gender-dependent, what may explain differences in frequency and severity of trigeminal nerve-associated disorders observed between women and men.

TG neurons in female mice showed increased expression of CGRP, this neuropeptide may act as the main mediator of trigeminal signaling during migraine.

This project has been supported by the Polish National Science Center, based on the Decision No. DEC-2012/05/B/NZ4/02385.

Research subject implemented with CePT infrastructure financed by the European Union – the European Regional Development Fund within the Operational Programme "Innovative economy for 2007-2013".

[A41]

Maraviroc modulates neuropathic pain by the CCR5 and CCR2 chemokine receptors

Kwiatkowski K., Rojewska E., Makuch W., Mika J.

Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Neuropathic pain develops as a result of pathological lesions within the nervous system. Recently were identified pronociceptive properties of chemokines and chemokine receptors. It was documented that the expression of CCR5 receptor is increased as a result of nerve injury, and it seems to be crucial for hyperalgesia and allodynia development. CCR5 is mostly expressed by microglial cells. What is more, previous research demonstrated that CCR5 and CCR2 chemokine receptors can link up in heterodimers. In order to understand the potential role of CCR5 and CCR2 in neuropathic pain, the aim of our studies was to examine the influence of selective antagonist of CCR5 receptor (maraviroc) on hyperalgesia and allodynia under neuropathy. Additionally we measured expression of CCR5, CCR2 and glial cell markers in neuropathic pain model. All experiments were carried out according to IASP rules (Zimmermann 1983). According to Yaksh and Rudy (1976) Wistar rats were implanted with intrathecal (i.th.) catheters. After seven days we performed chronic constriction injury (CCI) of the sciatic nerve according to Bennett and Xie (1988). On day 3 and 7 after CCI two behavioral tests were conducted to measure hyperalgesia (cold plate test) and allodynia (von Frey test). qRT-PCR was used to measure CCR5, CCR2, C1q and GFAP mRNA levels in spinal cord and dorsal root ganglia (DRG) in vehicle- and maraviroc-treated CCI-exposed rats. We provided evidence that chronic (once a day) i.th. administration of maraviroc attenuated the neuropathic pain symptoms on day 3 and 7 in CCI-exposed rats. qRT-PCR analysis indicate that maraviroc decreased activation of C1q-positive cells in spinal cord and DRG as measured on day 7 after CCI. Additionally we validated at the same time point that expression of CCR5 and CCR2 receptors is increased in spinal cord as well as in DRG after sciatic nerve injury. Interestingly, those changes are diminished by maraviroc chronic administration. Our results suggest that modulation of CCR5 and CCR2 by maraviroc might be a potential novel therapeutic approach for neuropathic pain treatment.

Supported by NCN2011/03/B/NZ4/00042 and statutory funds.

[A42]

Chronic treatment with a nitric oxide donor, molsidomine and L-DOPA, alone or in combination, differently affect cGMP concentration in the striatum of unilaterally 6-OHDA-lesioned rats

Lenda T., Kmiec I., Kaminska K., Czarnecka A., Konieczny J., Lorenc-Koci E.

Department of Neuropsychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

It is well-known that dopamine (DA) modulates the synthesis of second messengers: cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP). DA acting via striatal DA D1 receptor stimulation, activates adenylate cyclase (AC) and initiates cAMP synthesis while via DA D2 receptor stimulation inhibits this reaction. Moreover, DA-mediated stimulation of DA D1 receptors localized on the striatal nNOS-containing interneurons initiates NO synthesis while stimulation DA D2 heteroreceptors evokes opposite effects. Hence, DA-induced initiation of NO synthesis and then NO-mediated activation of soluble guanylyl cyclase (sGC) lead to the increased cGMP production. The cGMP formation is tightly regulated not only by enzymes contributing to its synthesis but also by an enzyme involved in its degradation. In the DA-deafferented rat striatum 6-8 weeks after lesion, the cGMP level was significantly decreased while cAMP increased. Chronic treatment with L-DOPA differently affected the cAMP and cGMP levels in the striatum (STR) of dyskinetic and non-dyskinetic rats. In dyskinetic rats, the nucleotide levels dramatically decreased in the ipsi- and contralateral STR. In non-dyskinetic rats, cAMP increased while cGMP decreased only in the contralateral STR.

The aim of the present study was to examine the effects of chronic treatment with the nitric oxide donor molsidomine and L-DOPA, alone and in combination, on the cGMP concentration in the STR of unilaterally 6-OHDA-lesioned rats.

The experiments were performed on male Wistar rats injected unilaterally with 6-OHDA (8 µg/4 µl) into the medial forebrain bundle (MFB). Control, sham-operated rats received vehicle instead of 6-OHDA. Two weeks later, 6-OHDA-lesioned rats exhibiting more than 100 contralateral rotations per 1 h in the apomorphine (0.25 mg/kg) test, indicative on an extensive loss of the nigrostriatal neurons, were treated chronically with molsidomine (2 mg/kg *i.p.*) and L-DOPA (25 mg/kg *i.p.*), once daily for successive 14 days. The concentration of cGMP in the ipsi- and contralateral STR was determined 1 hour after the last drug doses.

Our data show that 4 weeks after unilateral injection of 6-OHDA into the MFB, the cGMP concentrations in the ipsi- and contralateral STR were significantly increased compared to the corresponding sides of the sham-operated group receiving *i.p.* vehicle. Chronic treatment of unilaterally 6-OHDA-lesioned rats with molsidomine alone caused further increases in the cGMP levels on both sides of the STR vs. 6-OHDA-lesioned group treated *i.p.* with vehicle. However, chronic administration of L-DOPA alone to these rats enhanced the cGMP concentration in the contralateral STR while combined injection of molsidomine + L-DOPA increased significantly its level in the ipsilateral STR compared to the corresponding sides of 6-OHDA-lesioned group receiving *i.p.* vehicle.

The obtained results suggest that chronically administered molsidomine, can affect the occurrence of dyskinesias by modulating the striatal levels of cGMP.

This work was supported by a grant from the National Science Centre of Poland, based on decision No. UMO-2011/01/B/NZ4/01581.

[A43]

Shifted activation of NF-κB transcription factor in various cell subpopulations in rat model of cerebral ischemic stroke

Lietzau G.¹, Kowianski P.^{1,2}, Karwacki Z.³, Dziwiatkowski J.¹, Domaradzka-Pytel B.¹, Morys J.¹

¹Department of Anatomy and Neurobiology, Medical University of Gdansk, Gdansk, Poland

²Department of Health Sciences, Pomeranian University of Slupsk, Slupsk, Poland

³Department of Neuroanesthesiology, Medical University of Gdansk, Gdansk, Poland

Introduction: NF-κB plays a regulatory function in the CNS during development and in various pathological conditions. Dual, antagonistic role of NF-κB in cerebral stroke is related with diverse composition of the complexes of this transcription factor. NF-κB dimers containing RelA/p65 subunit exacerbate brain damage by inducing pro-inflammatory and pro-apoptotic genes, whereas c-Rel-containing dimers promote cell survival by inducing anti-apoptotic gene transcription. While NF-κB presence in neuronal nuclei was confirmed by numerous studies, its activation in glial cells still remains the subject of discussion.

Aim: The aim of this study was answering the following questions: 1) in which cell subpopulations and when activation of RelA/p65 and c-Rel subunits of NF-κB occurs

in the CNS after permanent cerebral ischemia?, 2) are there any changes in the DNA-NF- κ B binding activity during the first 24 hours of ischemic stroke?

Material and methods: Permanent middle cerebral artery occlusion (pMCAO) was performed in 40 middle-aged (72 weeks old) male Wistar rats. Animals were randomly assigned to 4 experimental groups with different post-ischemic survival period: 3 h, 6 h, 16 h and 24 h. Activation of NF- κ B was assessed by immunohistochemistry and immunoblotting. EMSA was performed to analyze the DNA binding activity of NF- κ B.

Results: After pMCAO activation of both subunits of NF- κ B was observed in neurons, astrocytes and microglia, but not in oligodendroglial cell lineage. Activation of "neuroprotective" c-Rel subunit occurred during the first 3 h, and was observed mainly in neurons. During the whole study period, an increased expression of RelA/p65 was found in the damaged tissue. Activation of this subunit in neurons was present in all experimental groups, however – in astroglia and microglia it was observed predominantly at 16 and 24 h after pMCAO. Substantial loss in the number of neurons during the first 24 hours of stroke was also present. In our study, transient decrease in NF- κ B-DNA binding activity at 6 h, and subsequent increase – at 16 h and 24 h were detected.

Conclusions: Regulatory function of NF- κ B transcription factor in the acute ischemic stroke is related with its presence in the various cellular subpopulations of the CNS. Prevalence of RelA/p65 over c-Rel subunit explains the dominance of pro-apoptotic action of NF- κ B in this pathological process. Changes in NF- κ B-DNA binding activity may result from the increased cell death of RelA/p65- and c-Rel-immunoreactive neurons during the first hours of ischemic insult and delayed activation of RelA/p65-containing dimers in astrocytes and microglial cells, that occurs no sooner than 16 hours after ischemia induction.

[A44]

Prenatal exposure to 4-para-nonylphenol disrupts estrogen receptor α , but not estrogen receptor β and retinoid X receptor signaling in mouse embryonic neuronal cells

Litwa E., Rzemieniec J., Wnuk A., Lason W., Kajta M.

Institute of Pharmacology Polish Academy of Sciences,
Department of Experimental Neuroendocrinology, Cracow, Poland

Nonylphenol (NP) belongs to a wide group of endocrine disrupting chemicals present in plastic wrappings, bottles, and textile paints. Endocrine disrupting chemicals are known to mimic, antagonize or modify the endogenous hormonal activity and cause deleterious effects on living organisms. Due to the structural similarity, nonylphenol is able to imitate the natural hormone 17-beta-estradiol by competing for the binding site of the estrogen. Although half life of NP in humans is very short, humans are continuously exposed to nonylphenol, especially by food intake. Recent data demonstrated that it accumulates in the brain in much higher concentrations than in other tissues. There is a growing body of evidence that exposure to endocrine disruptors during pregnancy or perinatally, impairs mental development of infants and children. However, mechanisms of action of nonylphenol on the brain tissue at early stages of neural development need to be clarified. In the present study we investigated the impact of prenatal exposure to nonylphenol on estrogen receptors (ER α , ER β) and retinoid X receptor (RXR) signaling in mouse hippocampal cells. CD1 Swiss Albino pregnant mice were exposed to nonylphenol from 6th to 16th gestation days. Hippocampal cells from control and NP-treated mouse embryos were cultured according to the standard procedures, and subjected either to NP or its solvent at 7 day *in vitro*. Hippocampal cell cultures derived from control and NP-treated mouse embryos expressed mRNAs and proteins for ER α , ER β , and RXR, as detected by real time-PCR and specific ELISAs. Addition of nonylphenol (5 μ M) to the cultures stimulated mRNA expression of the receptors, except for ER α mRNA that remained unchanged in the cultures of hippocampal cells from NP-treated embryos. Nonylphenol (5.10 μ M) also induced caspase-3 activity in hippocampal cultures and enhanced it up to approximately 170% of the control, independently of prenatal treatment of hippocampal tissue. In both types of hippocampal cultures, ER β and RXR antagonists, PHTPP and HX 531, attenuated the NP-induced effect. However, ER α antagonist MPP inhibited the NP-evoked caspase-3 activity only in cultures derived from NP-treated embryos. These data suggest that prenatal exposure of hippocampal tissue to NP disrupts ER α , but not ER β and RXR signaling pathways, mainly by affecting ER α mRNA expression and anti-apoptotic capacity of the receptor.

This work was supported by the Polish National Center of Science, grant no. 2011/01/N/NZ4/04950.

|A45|

Effects of mGluR5 positive and negative allosteric modulators on brain damage evoked by hypoxia-ischemia in neonatal rats

Makarewicz D.¹, Slomka M.¹, Danysz W.², Lazarewicz J.W.¹

¹Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Merz Pharmaceuticals GmbH, Frankfurt am Main, Germany

In the present study, we tested the hypothesis that the level of brain development determines the effectiveness of functional coupling between metabotropic glutamate receptor 5 (mGluR5) and *N*-methyl-*D*-aspartate receptors (NMDARs) in terms of neuroprotection. We examined the effects of negative and positive allosteric modulators of mGluR5 receptors (fenobam and ADX47273, respectively) on brain damage induced by hypoxia-ischemia (H-I) in 7-day-old rats. The test drugs were administered intraperitoneally 10 min after H-I. Rectal body temperature was measured for 2.5 h, and the weight of both hemispheres was determined 14 days after the insult. In the vehicle-treated groups, H-I reduced the weight of the ipsilateral (ischemic) hemisphere by approximately 33%. Fenobam (10 mg/kg) had no significant effect on brain damage or the weight of the contralateral (control) hemisphere. In contrast, fenobam (20 mg/kg) potentiated ischemic brain damage by 57.4%. ADX47273 (5, 10, and 30 mg/kg) had no significant effect on ischemic brain damage and did not interfere with the weight of the contralateral hemisphere. In all of the experimental groups, we detected no significant changes in rectal temperature. In conclusion, in 7-day-old rats, the bidirectional modulation of mGluR5 by fenobam (10 mg/kg) and ADX47273 (all doses tested) did not result in significant changes in H-I-evoked brain damage, supporting the hypothesis that the coupling of mGluR5 and NMDARs is a phenomenon that depends on brain development and is not fully operational in 7-day-old rat pups.

|A46|

Initial characterization of neurotrimin (Ntm) gene deficient mice

Mazitov T., Innos J., Vasar E.

Institute of Biomedicine and Translational Medicine, University of Tartu, Estonia

Introduction: Neurotrimin protein belongs to the IgLON family of proteins with Lsamp, Obcam, and Kilon. All these molecules are quite similar, highly glycosylated membrane proteins, characterised by three Ig domains and GPI anchor. They act as heterodimers forming protein complexes that stimulate and inhibit neurite outgrowth and guide axons. On the organism level, members of the IgLON family have been implicated in psychiatric disorders, obesity and certain types of cancer. The goal of our research group is to characterize the IgLON family as a whole by means of both, model organisms (knockout mice) and molecular methods. We have provided a thorough characterization of the phenotype of Lsamp-deficient mice and have now extended our research to include Ntm-deficient mice.

Material and methods: Neurotrimin (Ntm) gene deficient homozygous and heterozygous male and female mice and their wild-type littermates were tested in a battery of behavioral tests. Also some preliminary pharmacological experiments were performed to test for possible changes in major monoaminergic systems.

Results: First, we tested the animals for vision, hearing and olfaction and found that they have normal sensory abilities. We also tested Ntm-deficient mice in three anxiety tests (hyponeophagia, plus maze and light/dark box) and in the locomotor activity test (motility box) and found that their behavior does not differ from that of their wild-type littermates. However, we observed statistically significant differences in one of the learning and memory tests, the active avoidance test. Ntm-deficient mice learn slower than their wild-type and heterozygous littermates, probably exhibiting a learning impairment. Differences were also evident in pharmacological pilot studies with ethanol and amphetamine. Ntm-deficient mice were more sensitive to the sedative effect of ethanol, as it took much longer for them to regain the righting reflex after i.p. injection of 4 g/kg of ethanol. Furthermore, the locomotor stimulating effect of 5 mg/kg of amphetamine was weaker in Ntm-deficient mice than in wild type animals.

Conclusions: In this study, the basic phenotyping of neurotrimin (Ntm) gene knockout male and female mice was performed. These mice display normal sensory abilities, normal locomotor activity and normal anxiety level.

However, they have a learning impairment and altered sensitivity to ethanol and amphetamine, probably reflecting changes in the activity of major monoaminergic systems. It is remarkable that *Lsamp* deficient mice have a similar pharmacological phenotype, which is not surprising, however, because *Ntm* and *Lsamp* are close interaction partners and form heterodimers together. The study underlines the importance of studying all the members of the IgLON family of proteins together.

Poster session II

[B1]

Osteopontin gene and clinical course of multiple sclerosis in a Polish population

Michalowska-Wender G.^{1,2,3}, Biernacka-Lukanty J.^{1,3}, Michalak S.^{2,3,4}, Raczak B.^{1,3}, Wender M.²

¹Laboratory of Neurogenetics, Department of Neurology, University of Medical Sciences, Poznan, Poland

²Neuroimmunological Unit, Medical Research Center, Polish Academy of Sciences, Poznan, Poland

³Department of Neurology, University of Medical Sciences, Poznan, Poland

⁴Department Neurochemistry and Neuropathology, University of Medical Sciences, Poznan, Poland

Osteopontin (OPN) is a key cytokine involved in T-cell activation in multiple sclerosis (MS), and is therefore called an early T-cell activation gene. Studies performed in various geographical regions have demonstrated significant differences. In this study we analysed whether the OPN gene has an effect on MS occurrence and clinical course in a Polish population. Our study material consisted of 100 MS patients compared with control material. The clinical course was evaluated by the EDSS score. After extraction of DNA from peripheral blood cells, genotype and allele frequencies in exon 6 and 7 were examined using specific primers by a standard PCR reaction. The distribution of variables was tested with the D'Agostino-Pearson test. Then the statistical analysis based on one-way ANOVA, Mann-Whitney test and logistic regression was performed using a licensed version of MedCalc software.

The genotype distribution and allele frequencies were not similar between patients and controls but differences were not statistically significant. The heterozygous C/T genotype in position 8090 was detected in 21.6% of patients and in 10% of control material. Our results are negative with respect to the impact of the osteopontin gene on susceptibility to MS in Poland. Genetic polymorphism of OPN was established to be associated with MS in Japanese patients. It seems therefore that the haplotype structure may differ from one population to another. However, we have identified the 8090 T/T 9250 C/C osteopontin genotype associated with higher levels of disability in MS patients. Age is an independent factor influencing disability measures. Based on one-way ANOVA testing, the clinically significant genotypes were identified. The EDSS score was higher (3.0; 2.0-4.0, median; interquartile range) in 8090 T/T + 9250 C/C patients than in 8090 C/C + 9250 C/C MS patients (1.75; 1.5-2.0) ($p = 0.0120$). The disability in

8090 C/C + 9250 C/T MS patients was higher (EDSS = 3.75; 2.5-4.5) than in 8090 C/C + 9250 C/C MS patients (1.75; 1.5-2.0) ($p = 0.0137$). In logistic regression analysis of the effect of variables on EDSS scoring in the model, including sex, age and genotype categories, age was found to be independent factor influencing disability ($p = 0.0140$). No differences in age between genotype categories were found.

[B2]

L-histidine administration increases brain carnosine concentration in rats with thioacetamide-induced acute liver failure: a contribution to increased antioxidant capacity (TAC)?

Milewski K., Hilgier W., Albrecht J., Zielinska M.

Department of Neurotoxicology, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Hepatic encephalopathy (HE), a consequence of acute or chronic liver failure, is a complex neuropsychiatric disorder where oxidative/nitrosative stress (ONS) plays a decisive pathogenic role. ONS imposed on astrocytes causes their swelling which is the primary cause of cerebral edema, increased intracranial pressure herniation, and direct cause of death in acute HE. Existing evidence suggest that systemic administration of L-histidine (His) may be a potentially useful therapeutic agent for the brain edema associated with HE (Rama Rao 2007). The effect of His was primarily ascribed to inhibition of excessive Gln transport to mitochondria (the "Trojan Horse" hypothesis). However, His is known for its antioxidant action; a study from our laboratory has shown that His also prevent reduction of glutathione content in brain mitochondria of TAA rats (Ruszkiewicz 2013) and increases the total antioxidant capacity (TAC) of the brain (Ruszkiewicz and Albrecht 2014). One other *in vitro* study demonstrated reduced mitochondrial permeability transition (mPT) and oxidative stress in ammonia-exposed astrocytes pretreated with His (Rao 2010). We speculated that the action of His may be due to the action of its metabolites. Of these, histamine is a component of and thought to be related with inflammation, one of accompanying events in HE progression. Carnosine, a dipeptide may be associated with wide range of neuroprotective properties (Trombley 2000). It was also indicated that carnosine prevents TAA induced liver injury in rats (Mehmetcic 2007). Here we investigated the effects of acute HE on the rat blood and brain content of hista-

mine and carnosine. HE was induced using the thioacetamide model (300 mg TAA/kg 3 times *i.p.* in 24 h intervals). His was administrated *i.p.* 2 h before TAA (100 mg/kg). Brains and blood were collected 24 h after last injection.

Systemic administration of His had no lasting effect on its concentration in cerebral cortex and in blood plasma either in control and TAA rats, neither did it affect the concentration of histamine in there, countering its putative role in potentiating systemic inflammation in TAA rats. In contrast, the brain level of carnosine was significantly elevated after His treatment in both control and HE rats, suggesting that carnosine may be responsible for some beneficial effects of His administration in the setting of HE. Studies to verify this hypothesis are under way in our laboratory.

Supported by the National Science Centre grant No 2013/09/B/NZ4/00536.

|B3|

Antidepressant-like properties of 1,2,3,4-tetrahydroisoquinoline and its methyl derivative in animal models of depression

Mozdzen E., Michaluk J., Wasik A., Romanska I., Antkiewicz-Michaluk L.

Department of Neurochemistry, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Depression is a serious disorder in today's society, with estimate of lifetime prevalence as high as 21% of the general population in some developed countries. As defined by the American Psychiatric Association, depression is a heterogeneous disorder often manifested with symptoms at the psychological, behavioral and physiological levels. As with all diseases, approximations of both the disorder and the actions of corrective medications in laboratory animals are essential for the development of effective therapies. Most of the currently used antidepressant drugs are monoamine based, affecting the inhibition of re-uptake or metabolism of noradrenaline (NA), serotonin (5-HT), or dopamine (DA), because they play a key role in central nervous system (CNS) in pathophysiology of depression.

The present study is aimed to investigate the potential antidepressant properties of an endogenous amine 1,2,3,4-tetrahydroisoquinoline (TIQ) and its methyl derivative, 1-methyl-1,2,3,4-tetrahydroisoquinoline (1-MeTIQ). The experiments were carried out on male C57BL/6J mice (25 ± 2 g) (Charles River Laboratories, Sulzfeld, Germany). The antidepressant-like activity of tested compounds was

evaluated in the behavioral and neurochemical analysis. In the behavioral study, the forced swim test (FST) and tail suspension test (TST) were used as widely used rodent models sensitive to the effects of acute antidepressant treatments. Additionally, the motor function of mice was checked in the locomotor activity test after investigated drugs administration. TIQ and 1-MeTIQ were administrated in three different doses: 10, 25 or 50 mg/kg *i.p.* Imipramine (15 or 30 mg/kg *i.p.*) was used as a reference drug. Control mice were treated with saline. In the neurochemical *ex vivo* study, the levels of NA, 5-HT, DA and their metabolites, the rate of monoamines metabolism and their neuronal activity in different mice brain structures were determined by HPLC methodology with electrochemical detection. All procedures were approved by the Ethical Committee of the Institute of Pharmacology, Polish Academy of Sciences in Cracow.

The results of these studies have shown that TIQ and 1-MeTIQ administrated systematically to mice produced antidepressant-like effect in the FST. They significantly decreased the immobility time ($p < 0.01$), stronger than imipramine and did not affect the locomotor activity in mice. Additionally, TIQs increased the rate of NA metabolism, enhanced the factor of DA re-uptake inhibition with simultaneously decrease in the rate of 5-HT metabolism in mice brain structures. The results obtained indicate that TIQ and 1-MeTIQ possess a potential antidepressant activity in mice, suggesting their possible mechanism of action based mainly on activation of monoaminergic system – inhibition of MAO-dependent oxidation of monoamines and increase in their concentrations in different brain structures. In that light, both tested compounds may be effective for the depression therapy in a clinical setting.

These studies were financed by the project Interdisciplinary PhD Studies "Mol-Med".

|B4|

The actions of 5-methoxy-diisopropyltryptamine ('foxy') in rat prefrontal cortex and striatum

Noworyta-Sokolowska K.

Department of Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

5-methoxy-diisopropyltryptamine (5-MeO-DIPT, 'foxy') is one of the most popular 'club drug' in the United States, Canada, Japan and Europe. It has been noted in case reports that 5-MeO-DIPT induces visual and auditory hal-

lucinations, insomnia, emotional expression similarly to psilocin and psilocibin. That contributed to recognition these drug as a hallucinogen, however its mechanism of action in the brain is not fully understood. A comparison of the chemical structure of 5-MeO-DIPT shows structural similarities with serotonin (5-HT) and 3,4-methylenedioxy-methamphetamine (MDMA). That suggests its possible influence on monoamine systems such as dopamine (DA) and 5-HT as a main mechanism of action of these drug. Moreover, hallucinations noted after ‘foxy’ treatment indicate on possible influence on glutamate (GLU) system.

The aim of the study was to examine the extracellular level of DA, 5-HT, γ -aminobutyric acid (GABA) and GLU after 5-MeO-DIPT administration in freely moving rats.

All experiments were performed in a strict accordance with the Polish legal regulations concerning experiments on animals (Dz. U. 05.33.289). Determination of extracellular levels of DA, 5-HT, GABA and GLU on Wistar-Han male rats (280-300 g) was carried out using microdialysis technique. The probe implantation in the frontal cortex (AP +2.8, DL +0.8, V -5.7) and in the corpus striatum (AP +1.8, DL +2.8, V -7.0) took place 24 h before experiments. Dialysate fractions were analyzed by High Performance Liquid Chromatography (HPLC) with electrochemical detection. The chromatographic data was processed by Chromeleon v. 6.80 (Dionex, USA) software.

Single intraperitoneal administration of 5-MeO-DIPT given at a dose of 20 or 10 \times 2 mg/kg produced increase in DA and 5-HT release in the rat striatum and the frontal cortex. At the same time increase in GLU and GABA extracellular level was observed in the frontal cortex. However, transient increase in GLU release in rat striatum was followed by decrease in its level for the rest of experimental time. A slight but significant decrease in GABA extracellular level was noted in the striatum. An intriguing behaviour such as head shaking, salivation, walking backward, convulsions during first hour after 5-MeO-DIPT administration was observed as well as playing piano, tickles movement, tremor and muscle weakens were seen throughout the study.

Observed increase in DA and 5-HT extracellular level with concomitant decrease in metabolite extracellular level indicates on inhibition of DA and 5-HT transporters located on DA or 5-HT nerve terminals. Increase in extracellular GLU level may cause behavioural symptoms such as convulsions and head shakings and may also indicate on possible hallucinogenic drug action.

Supported by grant from National Centre of Science (NCN) no 2013/09/B/NZ7/04104.

[B5]

Paraquat induces changes in activity and supercomplexes assembly of mitochondrial complex I and mitochondria membranes fluidity in SH-SY5Y cells

Olech L.¹, Leskiewicz M.², Basta-Kaim A.², Dencher N.³, Kuter K.¹

¹Department of Neuropsychopharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

³Physical Biochemistry, Department of Chemistry, Technische Universität Darmstadt, Darmstadt, Germany

Introduction: Paraquat (PQ) is a herbicide which enhances risk of developing Parkinson’s disease (PD). It increases the level of reactive oxygen species (ROS) in cells via the redox cycling performed by many oxidoreductases. Previous studies showed decreased activity of respiratory complexes (Cxs) I and III by PQ. In the inner mitochondrial membrane Cxs form higher order structures named supercomplexes (SCxs), what has crucial effect to its functioning.

Aim: Our aim was to investigate the influence of PQ on activity of mitochondrial Cx I assembled in SCxs and as individual form in a model of neuron-like cells.

Material and methods: Differentiated SH-SY5Y cells (10 μ M retinoic acid in DMEM with 10% FBS for 7 days) were treated with 600 μ M PQ for 24 hours. Vitality (by MTT test), mortality (by LDH test) and endogenous ROS level (using H₂DCFDA) at this time point were analyzed. Cells were also incubated next 24 hours with fresh medium without PQ. Mitochondria were isolated directly after incubation with PQ as well as after additional time with withdrawal of toxin. After mitochondria solubilization blue-native electrophoresis with further in-gel enzymatic activity measurements of Cx I were performed. Mitochondrial membranes fluidity was measured using fluorescence anisotropy method.

Results: Incubation of cells with PQ caused decrease in vitality and had no influence on mortality. PQ elevated level of endogenous ROS. A significant decrease in activity of Cx I was observed in the three highest molecular weight SCxs. Withdrawal of PQ led to progressive loss in activity of Cx I in the same SCxs and further decreased the activity of complex I in lower mass SCxs. In this time point a slightly increased fluidity of mitochondrial membranes was also observed.

Conclusions: Elevated ROS level may be related to decrease in activity of Cx I in the highest molecular weight

SCxs. PQ caused progressive perturbation in mitochondria even after withdrawal of toxin.

The study was supported by the Statutory Funds of the Institute of Pharmacology, PAS, Kraków, Poland and Technische Universität Darmstadt and DAAD.

[B6]

Proteasome inhibitor abrogates the neuroprotective effect of hyperbaric oxygen preconditioning in experimental global cerebral ischemia

Ostrowski R.P., Stepień K., Pucko E., Wojda R., Matyja E.

Department of Experimental and Clinical Neuropathology, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Hyperbaric oxygen preconditioning (HBO-PC) has been known to provide neuroprotection against cerebral ischemia. However, it is not known whether HBO-PC is effective against severe global brain ischemia, induced by bilateral common carotid artery (CCA) occlusion and profound systemic hypotension due to exsanguinations (modified two vessel occlusion cerebral ischemic model; 2VO). Moreover, the role of proteasome in the mechanisms of HBO-PC has not been as yet investigated.

The aim of this study was to investigate whether brain injury that results from carotid occlusion and deep hypotension is reduced with HBO preconditioning, and to determine the involvement of proteasome in the preconditioning mechanism.

The experiments were performed on 48 male Wistar rats within body weight range of 280-350 g, allocated to the following groups: naive control, sham, rats with 2VO untreated (2VO group), rats with 2VO preconditioned with HBO (HBO-PC group) and rats with 2VO and the proteasome inhibitor MG132, injected *i.p.* at a dose of 1 mg/kg, at 30 min prior to every session of HBO-PC (MG132 group). HBO-PC procedure used 100% oxygen at 2.5 ATA 1 hr daily for 5 consecutive days with the last session at 24 hrs before ischemia. The ischemia was induced under ketamine and xylazine anesthesia by drawing blood from femoral artery to decrease mean arterial blood pressure to 30-35 mm Hg, followed by CCA bilateral occlusion for 5 min. Rats were examined for sensorimotor and T-maze performance deficits and the mortality was calculated. They were euthanized at 24 hrs for WB or at 7 days for brain histology.

After ischemia (2VO group), the number of surviving neurons in CA1 zone of the hippocampus was equivalent to only 5.7% of CA1 neurons in the sham control. However, in the HBO-PC group, the number of well preserved neurons was as high as 20.8% of that in the sham control. When the group preconditioned with HBO also received MG132, the number of surviving neurons equaled 2.7% of control. HBO-PC tended to reduce mortality (25% vs. 38% amongst untreated), although insignificantly. In the MG132 group, 45% of mortality was noted. HBO-PC resulted in 63.9% average T-maze scores as compared to 27.8% in the unpreconditioned rats after ischemia. The proteasome inhibitor combined with HBO-PC reduced the score to 33.3%. On Western blot, the preliminary results showed an increased level of apoptotic protein Bim (BCL-2-interacting mediator of cell death) in the brain after ischemia, while it was reduced in the brains of animals preconditioned with HBO. MG132 reversed the HBO-PC-induced decrease in brain levels of Bim.

In conclusion, the HBO-PC-induced brain protection against severe ischemic brain insult appears to be proteasome-dependent. In order to further verify the underlying mechanism, future studies will examine ubiquitin-ligases and proteasomes leading in concert to a depletion of pro-apoptotic proteins upon HBO-PC.

The authors want to thank Małgorzata Poździk, MSc for the excellent technical assistance.

[B7]

Mutated human beta-amyloid precursor protein gene (*APP-sw*) causes aberrant autophagy in NGF-differentiated neural PC-12 rat pheochromocytoma cells

Pajak B.¹, Kania E.¹, Orzechowski A.^{1,2}

¹Electron Microscopy Platform, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Department of Physiological Sciences, Faculty of Veterinary Medicine, Warsaw University of Life Sciences – SGGW, Warsaw, Poland

Pheochromocytoma PC-12 cells are immune to physiological stimuli directed to evoke programmed cell death. Besides, metabolic inhibitors are incapable to sensitize PC-12 cells to extrinsic or intrinsic apoptosis unless they are used in toxic concentrations. Surprisingly, these cells become receptive to cell deletion after human *APP-sw* gene expression. PC-12 cells were nucleofected with a plas-

mid containing amyloid beta precursor protein wild type APP(-wt), or swedish mutation APP(-sw) together with resistance genes to antibiotics (gentamicin and ampicillin) expressed under green fluorescent protein (GFP) gene promoter. Next, cells were subjected to 48 h differentiation programme (4% FBS + 2% HS + NGF in DMEM) followed by 48 h treatment with selected substances dissolved in 0.1% BSA + NGF in DMEM. Afterwards, cell cultures were subjected to molecular analysis. Presence of transgene and gene activity in nucleofected cells was evaluated by PCR and Western blot analysis, respectively. Transfected neural rat PC-12 cells were examined to assess cytotoxicity of normal APP processing product beta-amyloid via MTT and Neutral Red assay. We observed reduced cell viability in *GFP* vector + *APP-sw*-nucleofected cells (drop by 36%) but not in *GFP* vector-, or *GFP* vector+ *APP-wt*-nucleofected cells. Lower viability was accompanied by higher expression of A β 1-16 and elevated secretion of A β 1-40 (in average 53.58 pg/mL). At the ultrastructural level autophagy-like process was demonstrated to occur in *APP-sw*-nucleofected cells with numerous autophagosomes and multivesicular bodies but without autolysosomes. Human *APP-sw* gene is harmful to PC-12 cells and cells are additionally driven to incomplete autophagy-like process. When stimulated by TRAIL or nystatin, CLU protein expression accompanies early phase of autophagy.

This research was funded by National Science Centre (Poland), Grant No. UMO-2011/03/B/NZ7/02328.

[B8]

Mysterious neuropeptides: phoenixin and spexin. Are they really promising from the clinical viewpoint?

Palasz A., Rojczyk-Golebiewska E., Wiaderkiewicz R.

Department of Histology, Medical University of Silesia, Katowice, Poland

Recently, a lot of new regulatory peptides has been identified and described. Newly discovered and still very mysterious neuropeptides such as phoenixin and spexin appear to be a uniquely intriguing factors. Extremely poorly studied neuropeptide; phoenixin (PNX) has almost the same pattern of hypothalamic immunoreactivity as better known nesfatin-1. This intriguing endogenous factor may control the reproductive hormone secretion by the modulation of the pituitary GnRH receptor expression. Probably, phoenixin sensitizes the pituitary to releasing factors,

rather than directly promote the exocytosis of secretory vesicles to endocrine cells. Immunohistochemical studies revealed PNX positive cells in the rat hypothalamus, dorsal horn, spinal trigeminal tract, nucleus of the solitary tract and even in the trigeminal and nodose ganglia. It was also observed that exogenously administered PNX may preferentially suppress visceral as opposed to thermal pain. Recent reports suggest that the mechanism of signal transduction activated by PNX is MAPK/ERK pathway. Phoenixin may play significant role in reproduction and energy homeostasis at the level of the hypothalamic centres. Recent data may also implicate phoenixin in reproductive or metabolic disorders. Spexin which was identified through bioinformatics search strategy, a novel gene containing spexin has been named NPQ. These peptides have no homologies to other proteins. The rat spexin differs from the human and mouse molecules only by one amino acid at the C-terminal domain. In the brain, many spexin-positive neurons were visible, while neuroglia was usually negative. The hippocampal neurons showed moderate immunoreactivity as well as cerebellar Purkinje cells and brain stem neurons. Spexin seems to have multiple physiological functions in mammals. Studies using goldfish revealed the involvement of SPX in reproduction and food-intake regulation. Treatment of animals with SPX decreased the secretion of LH and suppressed appetite. Recent findings also demonstrate the central control of SPX in cardiovascular/renal function and nociception. Spexin may be a novel factor involved in weight regulation, with potential for obesity therapy.

[B9]

Mitochondria dysfunction and oxidative stress in neurodegenerative process influenced by the designer drug of abuse – N-benzylpiperazine

Persona K.¹, Polus A.², Goralska J.², Gruca A.², Dembinska-Kiec A.², Piekoszewski W.¹

¹Jagiellonian University, Faculty of Chemistry, Department of Analytical Chemistry, Cracow, Poland

²Jagiellonian University – Medical College, Department of Clinical Biochemistry, Chair of Clinical Biochemistry, Cracow, Poland

“Designer drugs of abuse” is a common name for substances advertised as “legal alternative to illegal compounds”. Despite of the introduction of prohibition in trade of medicinal products which mimic the effects of ille-

gal drugs, the problem of "legal highs" is still significant. Knowledge about the effects of action of such substances is scarce and mainly derived from the description of cases form toxicological wards. One of model compounds from the group of designer drugs of abuse is *N*-benzylpiperazine (BZP) – a psychoactive agent for the central nervous system with stimulating and sympaticomimetic properties. The mechanism of action of BZP is compared to the one of MDMA, while the effects are defined as 10-times weaker than the effects of d-amphetamine.

The aim of the study was to evaluate the impact of BZP on brain cells apoptosis including the potential role of mitochondria, the formation of oxidative stress, and disruption of cells energy state in presumptive neurodegenerative properties of this drug.

Experiments were performed using a line of human glial tumor cells (LN-18) incubated with BZP for 24 h. The choice of doses of BZP in the range of 10 mg/mL to 300 mg/mL were made through evaluation of LDH released into the culture medium as well as through consideration of the concentration of drug in the biological material taken from BZP users. Analysis of parameters associated with mitochondrial dysfunction and oxidative stress formation, such as: changes in mitochondrial membrane potential ($\Delta\Psi_m$) and reactive oxygen species (ROS) were measured with the use of flow cytometry technique (FACS), whereas changes in ATP level were examined using luminescent method. The activity of caspases: 3, 8, 9 was analyzed in fluorescence measurements. Increase of oxidative DNA damage was evaluated by ELISA test through determination of DNA oxidative damage marker: 8-OHdG.

The conducted research demonstrated statistically significant changes in increased mitochondrial membrane potential and increased ROS production, along with decreased ATP level. Increased changes in activation of caspase-3 which are a confirmation of cell apoptosis undergoing under the influence of BZP were observed, however no statistically significant changes in activation of caspase-8 and -9 were stated. Elevated ROS generation resulted in oxidative DNA structure changes, which was confirmed by increased levels of 8-OHdG in the cells.

The results indicate that BZP at concentrations comparable to the ones detected in biological material taken from people taking this drug can induce apoptosis in glial cells by affecting the mitochondrial dysfunction and resulting in the oxidative stress generation. The results may also indicate neurodegenerative properties of BZP.

K. Persona acknowledges financial support from the Interdisciplinary PhD Studies project "Molecular sciences for medicine" (co-financed by the European Social Fund within the Human Capital Operational Programme).

K. Persona received funds to prepare her doctoral dissertation from the National Science Centre in the frame-

work of a doctoral scholarship funding (ETIUDA) based on the decision no. DEC-2014/12/NZ7/00262.

|B10|

Identification and classification of transcriptional variants and non-coding RNAs induced by antidepressants in the mouse striatum

Piechota M.¹, Golda S.¹, Jantas D.², Przewlocki R.^{1,3}, Korostynski M.¹

¹Department of Molecular Neuropharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

³Department of Neurobiology and Neuropsychology, IPS, Jagiellonian University, Cracow, Poland

The therapeutic effects of antidepressants are mediated by neuroplastic changes in the brain systems controlling drive and motivation. Despite well-known pharmacological actions of antidepressants, we are still far from having a clear picture of the molecular programs initiated by these drugs. We used next-generation sequencing and bioinformatics approach to build a genome-wide inventory of protein coding and non-coding transcripts altered by antidepressants. Genome expression was analyzed in mouse striatum 4 hours after the administration of mianserine (20 mg/kg, *i.p.*) or tranylcypromine (20 mg/kg, *i.p.*) treatment. We identified 153 gene transcripts of various biotypes regulated (FDR < 0.1%) in response to antidepressants. These transcripts are expressed in five different patterns and are enriched in protein coding variants (77%). Expression of different types of drug inducible transcriptional variants was verified using qPCR, including retained intron (*Clk1*), alternative TSS (*Mctp1*), small RNA (*Mir1983*) or long non-coding RNA (*Neat1*). We also indicated that two key gene regulatory systems are active in separate cellular compartments of the striatum, CREB1/SRF in neurons and GR/CTCF in glia. The results suggest that critical step of molecular regulation of neuroplasticity takes place on the level of transcriptional units rather than genes. This regulation may constitute a transcriptional trigger for development of long-lasting effects of antidepressants in the brain.

This work was supported by POIG De-Me-Ter 3.1 and NCN grant 2011/03/D/NZ3/01686.

[B11]

The role of CXCL9 and CXCL12 in nociceptive transmission under diabetes

Pilat D., Zychowska M., Rojewska E., Mika J.

Department of Pain Pharmacology, Institute of Pharmacology,
Polish Academy of Sciences, Cracow, Poland

Neuropathic pain is one of the most common late complications of diabetes. It is known that neuronal mechanisms can be modulated by pronociceptive mediators released by non-neuronal cells which become activated in the spinal cord during diabetes. However, the signals that mediate the spread of nociceptive signaling remain to be established. Chemokines seems to be crucial for mediating inflammation, although their role in nociceptive processes still needs to be evaluated. Both CXCL9 and CXCL12 are thought to mediate inflammatory processes in people suffering from diabetes, taking into account that both chemokines have increased plasma levels in patients. The role of those chemokines under diabetic neuropathic pain is unknown. The aim of our study was to determine whether in the mice model of diabetic neuropathy there is a change in the level of CXCL9 and CXCL12, and that intrathecal administration of these chemokines does affect nociceptive processes in naive mice. Male Albino-Swiss mice were rendered diabetic by an intraperitoneal injection of streptozotocin (STZ). The experiments were carried out 7 days after the injection of streptozotocin. To analyze if CXCL9 and CXCL12 may be involved in nociceptive transmission, we used RT-qPCR and antibody array techniques. Additionally these chemokines were administrated by an intrathecal injection to the naive mice. Behavioral tests conducted to measure hyperalgesia (cold plate test) and allodynia (von Frey test) were carried out 1, 4 and 24 hours after the injection. All experiments were carried out according to IASP rules (Zimmermann 1983). We have demonstrated that a single streptozotocin injection in mice can induce significant neuropathic pain along with an increase in plasma glucose levels. Seven days after the administration of STZ, an upregulation of CXCL9 and downregulation of CXCL12 mRNA were observed and measured by RT-qPCR. In parallel we observed an upregulation of protein level of both chemokines, as measured by an antibody array. The intrathecal administration of CXCL9 and CXCL12 showed its pronociceptive properties such as allodynia and hyperalgesia. In summary, we provided evidence for a development of allodynia and hyperalgesia following the administration of CXCL9 and CXCL12 in a naive mice. Therefore, we suggest that both studied chemokines, which are

upregulated in diabetic neuropathy, are one of the reasons of pain development.

This work was supported by statutory funds and NCN grants 2012/05/N/NZ4/02416 and 2011/03/B/NZ4/00042.

[B12]

Parthenolide down-regulates pronociceptive factors (IL-1beta, IL-18 and iNOS) and enhances morphine effectiveness under neuropathic pain

Piotrowska A., Popiolek-Barczyk K., Kolosowska N., Makuch W., Rojewska E., Pilat D., Mika J.

Department of Pain Pharmacology, Institute of Pharmacology,
Polish Academy of Sciences, Cracow, Poland

Neuropathic pain is caused by damage of the nervous system. This type of pain is resistant to conventional analgesics, including the effects of morphine, what comprises a serious clinical problem. Treatment of neuropathic pain remains a challenge and requires combination therapy, because of not fully understood pathomechanism, as well as many components involved in nociceptive transmission. The goal of our studies was to determine the influence of parthenolide (PTL; an inhibitor of the NF- κ B) on neuropathic pain development and changes in pronociceptive factors. Moreover, it was interesting to examine a possible influence of PTL administration on effectiveness of morphine in rat model of neuropathic pain. Research was carried out on Wistar rats implanted with intrathecal (*i.t.*) catheters. Neuropathic pain was developed using Bennett's model (chronic constriction injury (CCI) to the sciatic nerve). PTL (5 μ g/5 μ l) was dissolved in 50% DMSO and preemptively administered *i.t.* 16 h and 1 h before CCI and then once daily for 7 days. Two behavioral tests were conducted to measure allodynia (von Frey test) and hyperalgesia (cold plate test). Morphine was injected once, on day 7 after CCI and 60 min after morning PTL administration. Experiments were carried out according to IASP recommendations and were approved by the local Bioethics Committee. Behavioral studies have shown that chronic *i.t.* administration of PTL attenuated the allodynia and hyperalgesia on days 3, 5 and 7 after CCI. Western blot results demonstrated down-regulation of pronociceptive cytokines: IL-1beta, IL-18 and iNOS, parallel to down-regulation of p38MAPK as measured on day 7 after chronic PTL administration. Chronic treatment of PTL enhanced morphine analgesia, which was proven

by von Frey and cold plate tests. In summary, these results suggest that PTL reduces neuropathic pain by modulating the signaling pathways regulated by p38MAPK and by down-regulation of the expression of pronociceptive factors produced by the microglia cells. PTL is especially promising because it is already used in human therapy; our work provides the first evidence for its analgesic potency and beneficial properties on morphine effectiveness during neuropathic pain.

This work was supported by grant OPUS NCN 2011/03/B/NZ4/00042, and statutory funds. K. Popiolek-Barczyk is a Ph.D. student and has a scholarship from the KNOW sponsored by the Ministry of Science and Higher Education, Poland.

[B13]

The neuroprotective role of PKC β II in ischemia-reperfusion injury

Poleszak O., Beresewicz M., Kruszynska D., Zablocka B.

Molecular Biology Unit, Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Protein kinase C β (PKC β) is a serine/threonine kinase whose role in neuronal cells survival/ protection needs further investigation. Previous studies revealed that ischemic injury induce a rapid and long lasting increase of PKC β in mitochondrial fraction but mainly in ischemia-resistant part of hippocampus, what may bespeak neuroprotection. Hence, here we set out to determine the role of PKC β isozymes β I and β II in two models of brain ischemia (i) *in vitro* in rat organotypic hippocampal cultures subjected to excitotoxic injury which mimics ischemic injury and (ii) *in vivo* – transient brain ischemia in gerbils. Therefore we tested: (i) the effect of ischemic injury on the amount of both PKC β isozymes and activity of PKC β II in mitochondria isolated from ischemia-vulnerable (CA1) and ischemia resistant (CA2-4, DG) hippocampal regions, (ii) the effect of inhibition of PKC β isozymes in organotypic hippocampal cultures subjected to excitotoxic injury and (iii) by pull down chromatography followed by mass spectrometry the potential PKC β II mitochondrial partners to elucidate PKC β II-mediated neuroprotective mechanisms. In our studies we have observed that transient ischemic episode induces a significant elevation of PKC β II protein level and activity in ischemia-resistant part of hippocampus in mitochondrial fraction. Moreover, exclusive inhibition of PKC β II is sufficient to obtain 100% of damage in hippocampal

organotypic slices. Thus results suggest that it is rather PKC β II than PKC β I which may be a key element responsible for neuroprotection. To elucidate the mechanisms of PKC β II-mediated neuroprotection its mitochondrial partners were identified. The analysis recognised a number of proteins belonging to mitochondrial inner and outer membrane as well as mitochondrial matrix. They are involved in regulation of different metabolic pathways such as fatty acid oxidation, glucose metabolism, electron transport chain and others. Using phosphosite database (<http://www.phosphosite.org>) we verified that most of identified PKC β II-partners have putative PKC phosphorylation sites indicating that regulation of these metabolic pathways by PKC β II phosphorylation may lead to PKC β II-mediated neuroprotection. The functional significance of the PKC β phosphorylation is being studied.

The project is supported by NSC grant 2012/05/B/NZ/00415 and KNOW-MMRC project.

[B14]

Morphology of synapses and expression of selected proteins involved in synaptic transmission in the mouse azoxymethane model of hepatic encephalopathy

Popek M.¹, Frontczak-Baniewicz M.², Zablocka B.³, Albrecht J.¹, Zielinska M.¹

¹Department of Neurotoxicology, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Electron Microscopy Unit, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

³Molecular Biology Unit, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Hepatic encephalopathy (HE) is characterized by imbalance between excitatory and inhibitory neurotransmission in disfavor of the former which becomes potentiated on transition from acute to chronic stages of the disease. The established view is that ammonia, a key neurotoxin, primarily affects astrocytes, leaving neurons relatively spared. However, impairment of the expression and/or function of specific astrocytic proteins (e.g.: glutamate transporters GLT-1, GLAST, Na-K-Cl co-transporter or potassium channel Kir4.1 or trombospondin-1) results in synaptic dysfunction, mainly involving the NMDA class of Glu receptors. However, there is limited body of evidence concerning the morphology of the synapses in HE-affected brain, particularly the impact of HE on the activity of pro-

teins involved in the NMDA receptor function, trafficking, or composition.

In the present study we used C57Bl6 mice subjected to a hepatotoxic insult (azoxymethane (AOM) administered *i.p.* at 100 mg/kg of body weight) as an experimental model of acute liver failure. We commenced with examining synaptic ultrastructure in cerebral cortex of AOM mice. Next, we compared the subunit composition of NMDA receptor in control and AOM mice, by measuring the expression of the two key subunits, NR1 and NR2A. In order to assess the synaptic hardware, mRNA and/or protein levels of postsynaptic density protein 95 (PSD-95), protein kinase C zeta (PKC ζ) and PKM ξ , an independent fragment of catalytic domain of PKC ζ were quantified using real-time and Western blot analysis.

Cerebral cortex of AOM mice was characterized by the presence of increased numbers of enlarged synapses showing abundance of synaptic vesicles in the presynaptic zone. The expression of NR1, a major NMDA receptor subunit in a mice cerebral cortex was slightly increased, whereas NR2A subunit remained unchanged in AOM model. The amount of PSD-95, the protein involved in anchoring of NMDA receptor subunits, was increased by ~40% compared to control. The expression of PKC ξ involved in the synaptic events was increased by ~30% and ~40% at the mRNA and protein level, respectively. However, expression of PKM ξ , a protein with a unique role in LTP maintenance, was unchanged. Of note, PKC ξ protein level was reduced by ~20% in P2 membrane fraction and elevated by ~30% in S2 cytosolic fraction in AOM mice, suggesting impaired trafficking of PKC ξ to the plasma membrane. While implications of the above described changes for neurotransmission remain to be analyzed in further detail, this report is to our knowledge the first to demonstrate that changes in the synaptic hardware which may have direct bearing on its functioning, including LTP, do occur in the acute stage of HE.

Supported by the Polish-Norwegian grant “CORE”, Grant Number: Pol-Nor/196190/26/2013 and the Leading National Research Centre (KNOW).

[B15]

Expression and the level of selected apoptotic proteins and *Trp53* mutations in PS/APP mice

Postrach I.¹, Rozycka A.², Strugala A.³, Oczkowska A.¹, Florczak-Wypianska J.⁴, Dezor M.¹, Jagodzinski P.P.², Kozubski W.⁴, Dorszewska J.¹

¹Laboratory of Neurobiology, Department of Neurology, Poznan University of Medical Sciences, Poznan, Poland

²Department of Biochemistry and Molecular Biology, Poznan University of Medical Sciences, Poznan, Poland

³Department of Molecular and Systems Biology, Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland

⁴Chair and Department of Neurology, Poznan University of Medical Sciences, Poznan, Poland

Alzheimer’s disease (AD) is the most common, incurable and progressive neurodegenerative disease. There is a number of factors involved in the pathogenesis of AD, i.e. amyloid precursor protein (*APP*) gene. Mutations in the *APP* gene contribute to the familial AD (FAD), by changes in the processing of this gene resulting in increase of β -amyloid ($A\beta$) level. In double transgenic mouse model of AD, mutation of APP695 is associated with early-onset of FAD. In aged transgenic mice show elevated level of $A\beta$, which may cause the generation of free radicals. Free radicals may oxidize macromolecules in cells, such as DNA. 8-Oxoguanine DNA glycosylase 1 (OGG1) is a main DNA repair enzyme that excises 8-oxo-2'-deoxyguanosine from DNA. DNA damage may induce apoptosis in a wild-type p53-expressing cell. Bcl-2 is an anti-apoptotic protein, which prevents caspase-3 activation by Bax protein. Regulation of Bcl-2 family gene expression is not completely understood in AD.

The aim of the study was the analysis of *Trp53* gene mutations and expression, and analysis of p53, OGG1, tumor necrosis factor alpha (TNF-alpha), Bax and Bcl-2 protein levels in Alzheimer’s disease model PS/APP and control mice.

The studies were performed on female double transgenic PS/APP of B6.Cg-Tg(APP695)3DBo Tg(PSEN1dE9)S9Dbo/J strain (Jackson Lab., USA; about 8-months old) and young adults (8-12-weeks old) and aged-match control C57BL/6J strain mice. Brains of the animals were isolated and divided into three structures: cerebral gray matter (GM), subcortical white matter (WM) and cerebellum (C). The analysis of *Trp53* mutations was performed by using DNA sequencing and expression of this gene was carried out with use of RT-PCR and the level of p53 protein and other analysed apoptotic proteins was performed by Western blotting.

Our studies showed that the frequency of mutations was almost fourfold higher in PS/APP mice (A929T, A857G; 44%), compared to controls (T854G, G859A; 14%). However, it was shown that the expression of *Trp53* gene increased only in GM and C of PS/APP mice ($p < 0.05$) as compared to aged-match controls. Moreover, in GM and WM of PS/APP mice without *Trp53* mutations occurred a tendency to increase p53 protein levels together with elevated *Trp53* gene expression. Simultaneously, in PS/APP mice the presence of *Trp53* mutations did not lead to changes in expression and levels of Bax, Bcl-2, OGG1 and TNF- α proteins. In PS/APP mice we showed the tendency to decreased ratio of Bax:Bcl-2.

It seems that mutations of *Trp53* gene may lead to changes in expression especially of this gene and levels of p53 protein in PS/APP mice.

[B16]

Comparison of the neuroprotective potential of neuropeptide Y (13-36) and ACPT-I in spontaneously hypertensive rats subjected to transient focal cerebral ischemia using functional tests

Przykaza L.¹, Domin H.², Boguszewski P.³, Smialowska M.², Kozniowska E.¹

¹Laboratory of Experimental Neurosurgery, Department of Neurosurgery, M. Mossakowski Medical Research Centre Polish Academy of Sciences, Warsaw, Poland

²Department of Neurobiology, Institute of Pharmacology, Polish Academy of Sciences, Krakow, Poland

³Laboratory of Limbic System, Nencki Institute of Experimental Biology Polish Academy of Sciences, Warsaw, Poland

The aim of this study was to evaluate and compare neuroprotective potential of neuropeptide Y (13-36) (NPY₁₃₋₃₆) and (1S,3R,4S)-1-aminocyclopentane-1,2,4-tricarboxylic acid (ACPT-I) against ischemic brain damage in spontaneously hypertensive rats (SHR), using functional tests and macroscopic morphological evaluation of brain infarction volume. NPY₁₃₋₃₆ is a specific agonist of presynaptic neuronal Y2 receptors. Numerous previous studies have shown that activation of Y2 receptors inhibits glutamate release and protects against excitotoxicity *in vitro* and *in vivo*. ACPT-I is an unspecific agonist of group III metabotropic glutamate receptors (mGluR), which are known also to decrease excitatory transmission. ACPT-I has been shown to suppress seizure activity in animal epilepsy models and to protect neurons against kainate-induced excitotoxicity *in vitro* and *in vivo*.

Although there is a theoretical basis for a neuroprotective effect of both compounds against ischemic brain damage and a decrease in the size of ischemic infarct after inhibition of glutamate release has been demonstrated, no data is available evaluating post-ischemic functional state after treatment with either NPY₁₃₋₃₆ or ACPT-I. Forty one male SHRs, weighing 270-300 g, were used for this study. Transient focal cerebral ischemia was induced for 90 min by the intraluminal occlusion of the right middle cerebral artery (MCA). Microflow (LDF) in the brain cortex supplied by this artery was continuously monitored with a help of a laser-Doppler flowmeter starting before the occlusion of the vessel. NPY₁₃₋₃₆ (10 μ g) or vehicle were administered into the right lateral ventricle either 30 min after MCA occlusion or 30 min after reperfusion. ACPT-I (30 mg/kg) or vehicle were administered intraperitoneally in separate groups of rats according to the same paradigm. Behavioral tests (CatWalk, open field and forelimb placing) were performed before ischemia and 72 hours after reperfusion. The area of infarction was evaluated after the completion of behavioral tests on brain tissue slices stained with 2,3,5-triphenyltetrazolium chloride (TTC) using a computer based image analysis system. Both NPY₁₃₋₃₆ and ACPT-I significantly diminished volume of infarction and improved selected gait parameters even when administered during reperfusion. The forelimb placing reflex was improved, however, only when neuroprotective compounds were administered 30 min after MCA occlusion. Moreover, ACPT-I occurred to be more efficient in improving rats mobility in the open field. Thus, these results suggest that ACPT-I exerts better protection than NPY₁₃₋₃₆ in this model of ischemia.

The study was supported by grant No. N N401 091037 from MS&HE.

[B17]

Genes associated with minocycline efficacy in neuropathic pain

Rojewska E.¹, Korostynski M.², Przewlocki R.², Przewlocka B.¹, Mika J.¹

¹Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

²Department of Molecular Neuropharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Microarray analysis give us an opportunity to examine the molecular mechanisms of neuropathy and to find a new targets for neuropathic pain treatment. Last studies show multiple endogenous factors, which are involved in the chronic

pain development. It is known that the neuroimmune interactions play important role in this process and they may be one of the reasons for the loss of efficacy for many analgesics. Our previous data indicated the importance of spinal microglia activation-related genes in the development of neuropathic pain and their participation in the persistence of its symptoms. Repeated minocycline (30 mg/kg, *i.p.*) was shown to relieve neuropathic pain symptoms by inhibiting the spinal microglia activation and, in consequence, reduced release of numerous cytokines. The genes associated with minocycline efficacy under neuropathy should provide insight into the etiology of pain. Therefore, the aim of the our study was to determine, at the spinal cord level, which of the genes that were changed after sciatic nerve injury are modulated by minocycline. The microarray experiments helped us to identify numerous transcripts that are important for the development of neuropathy and may be possible targets for drug therapy. The DNA microarray method allowed us to choose 93 transcripts (transcription pattern T) from 22 500 studied transcripts, due to their abundance level changes after sciatic nerve injury, as well as 39 transcripts that were modulated by repeated administration of minocycline. To further research we have selected two gene expression patterns T1 and T2. The first transcription pattern T1 consisted of 10 transcripts with mRNA abundance levels rising after the injury and minocycline inhibit the effect of the injury; and the T2 transcription pattern have 6 transcripts with mRNA abundance levels dropping following CCI and minocycline reverses the effect. Additionally, we studied changes in selected gene expression by RT-PCR. In summary, our gene analysis offers the first step towards future research into genes that can be modulated and may be good targets for effective and safe therapies for the treatment of neuropathic pain.

Supported by statutory funds and NCN grant 2011/03/B/NZ4/00042.

|B18|

Anti-ischemic effects of 3,3'-diindolylmethane in mouse hippocampal cells at early developmental stages

Rzemieniec J., Litwa E., Wnuk A., Lason W., Kajta M.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Cerebral ischemia is a condition in which insufficient blood flow results in poor oxygen and glucose supply to

the brain, thus causing brain damage or stroke. During the neonatal period of life, ischemia appears as a major risk factor of cerebral palsy, seizures and cognitive impairments. Ischemic episodes have been increasingly recognized in children, but diagnosis and management is difficult because of diversity of underlying risk factors and the absence of a uniform treatment approach. 3,3'-diindolylmethane (DIM) is a plant-derived, acid-catalyzed condensation product of indole-3-carbinol that exhibits property of selective aryl hydrocarbon receptor modulator, and shows anti-carcinogenic capacity by inducing apoptosis in various cancer cells. Recent studies also indicated neuroprotective potential of DIM against brain inflammation and neurodegenerative diseases. The aim of present study was to evaluate neuroprotective effect of DIM in mouse hippocampal cells exposed to ischemia at early developmental stages. In this study, ischemic conditions were mimicked by exposure of embryonic hippocampal cells in primary cultures at 2, 7 and 12 days *in vitro* to 6 h lasting oxygen and glucose deprivation (OGD). Caspase-3 activity and lactate dehydrogenase (LDH) release were measured at 18 h of reoxygenation. In this study paradigm, OGD induced LDH release in mouse hippocampal cultures and increased its level by 23, 165 and 40% at 2, 7 and 12 days *in vitro*, respectively. OGD also caused about 30% enhancement in caspase-3 activity, but only in neuronal cells at later studied days *in vitro*. DIM used in wide concentration range (0.01-10 mM) inhibited the ischemia-induced LDH release in mouse neuronal cells at all analyzed stages. In addition, DIM in concentrations of 0.01 and 10 mM reduced the OGD-induced caspase-3 activity in 7 and 12 days old hippocampal cultures. This study demonstrated strong protective and anti-apoptotic effects of DIM in neuronal cells exposed to ischemia/OGD at early developmental stages. According to our data, the selective aryl hydrocarbon receptor modulator may represent novel therapeutic approach against ischemic brain injury.

This study was supported by the Polish National Center of Science grant No. 2011/01/N/NZ3/04786.

|B19|

Induction of oxidative stress in rat brain after repeated oral exposure to silver nanoparticles

Skalska J.¹, Frontczak-Baniewicz M.², Lenkiewicz A.¹, Struzynska L.¹

¹Laboratory of Pathoneurochemistry, Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Electron Microscopy Platform, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

The unique physical, chemical and biological properties of silver nanoparticles (AgNPs) have led to their extensive use in many industrial applications, including food industry, textile engineering and silver-based consumer products (e.g. cosmetics, cleansers, medical devices).

However, silver nanoparticles can interact with many substances in the human body due to their high reactivity, which might result in unknown hazards. Presently, only a few studies are dedicated for investigating their toxic potential. Most of them have been conducted using *in vitro* models.

The aim of the present study was to investigate whether commercially available small-sized silver nanoparticles induce oxidative stress in male adult rats and to compare their effect with the effect caused by silver ions.

Animals were exposed to oral application of 10 ± 4 nm AgNPs or silver ions in concentration 0.2 mg/kg for 14 days. Using transmission electron microscopy (TEM), nanosized granules were detected in the brain of exposed rats. To assess oxidative stress in the brain homogenates from exposed animals, ROS level and membrane peroxidation were measured. Additionally, the influence of AgNPs and silver ions on antioxidant enzymes activity (catalase, superoxide dismutase, glutathione peroxidase) was evaluated. Despite this, the ratio of reduced to oxidized glutathione was investigated, so was the level of protein glutathionylation. Our results show, that exposition to AgNPs or silver ions resulted in increase in ROS level, lipid peroxidation and antioxidant enzymes activity in the brain homogenates, whereas the ratio of reduced to oxidized glutathione significantly decreased.

These results indicate, that the induction of oxidative stress under condition of AgNPs or silver ions exposure, is almost certainly one of the mechanisms of their neurotoxicity.

|B20|

Prenatal stress enhances the production of pro-inflammatory cytokines: studies in hippocampal organotypic cultures

Slusarczyk J., Trojan E., Glombik K., Basta-Kaim A.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Inflammatory processes may be one of the most important causes of the development and progression of affective disorders. Due to a cytokine theory of depression changes in production of cytokines may play the role in the etiology of these disease. It has been observed that IFN- γ , IL-1 or IL-2 administration may lead to manifestation of symptoms and behavioral changes resembling those observed in major depression in humans. On the other hand, stress is a very important factor investigated in context of mood disorders. Exposure to stressful events, especially in the critical time of development as prenatal or early postnatal period, influences brain homeostasis by affecting the nervous, endocrine and immune systems and may lead to disturbances in the neurodevelopment in adulthood.

In our studies, we determined the effect of prenatal stress procedure – an animal model of depression – on the cytokine levels in hippocampal slice cultures. This *ex vivo* procedure allows preservation of natural connections between the immune, nervous and endocrine systems, cytoarchitecture and functional neuronal-glial interactions. The hippocampus was chosen for the study due to its role in the pathogenesis of depression.

Pregnant Sprague-Dawley rats were subjected daily to three (at 9.00, 12.00, 17.00 o'clock) stress sessions from 14th day of pregnancy until delivery. Control pregnant females were left undisturbed in their home cages. Hippocampal slice cultures were prepared from 7-8 day old rats, according to the method previously described by Stoppini *et al.* (1991) with slight modifications. Hippocampi, isolated from control and prenatally stressed animals were sectioned into 350 μ m transverse slices, separated and then transferred onto porous membrane inserts of 6-well culture plates. Each well contained 1 ml culture medium composed of 50% DMEM; 25% HBSS; 25% Horse Serum; 5 mg/ml glucose; amphotericin B; penicillin–streptomycin). Cultures were initiated in a regular, 25% horse serum-containing medium which was gradually replaced by a serum-free, defined-solution-based medium. Cultures were maintained for 7-9 days under standard conditions. The levels of cytokines (TNF- α , IL-1 β , IL-6) in supernatants were quantified by specific ELISA kits.

Prenatal stress procedure enhanced the release of IL-1 β and TNF- α in hippocampal slices. Moreover, we observed an increased production of IL-6 in hippocampal organotypic cultures obtained from animals after prenatal stress procedure.

Our results, using the organotypic cultures, show that prenatal stress may have an important influence on hippocampus, the brain structure, which plays a key role in the pathogenesis of depressive disorders. Increased inflammatory status after stress may affect these structure especially the neuron-glia interaction.

This work was supported by grant no. 2013/09/B/NZ7/04096.

[B21]

Comparison of pronociceptive system's activity in two neuropathic pain models in mice

Starnowska J., Rojewska E., Mika J., Przewlocka B.

Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Neuropathic pain may be caused by various conditions that result in nerve damage, including traumatic injury or disease, like diabetes. Typical painful symptoms, such as hyperalgesia (enhanced response to painful stimuli) and allodynia (painful response to neutral stimuli), are thought to be related to constant over-activation of pronociceptive system. Identification of biochemical components that comprise on escalated nociceptive mechanisms may help in finding an efficient remedy premised on re-balancing pronociceptive system's functions. In our study we compared two mice neuropathic pain models that refer to the common clinical neuropathy causes: Bennett's chronic constriction injury model (CCI) with sciatic nerve damage, and streptozotocin-induced diabetic neuropathy. Male CD-1 Albino Swiss mice were tested at 7th, 14th, 21st and 28th day after CCI surgery or streptozotocin (STZ) intraperitoneal administration to assess allodynia and hyperalgesia development, with usage of von Frey and cold plate tests, respectively. Also the glucose level in animals' blood was measured to assess diabetes development. All animals that have not developed neuropathy after nerve injury were excluded from the study as well as animals with no diabetes in STZ model. The curve of allodynia and hyperalgesia development is different for these two models. While hyperalgesia and allodynia intensity is similar on day 7 in both models, it differentiates markedly on day 14 and on day 28.

In the latter time point much weaker effects are obtained in a model of CCI and, in contrast, stronger in the STZ model. This proves that neuropathy symptoms in diabetes build up over time, while symptoms in CCI model have opposite tendency. At every time point aforementioned probes of spinal cord tissue were taken to be further analyzed with qRT-PCR method for the activity of pronociceptive systems by analysis of the changes in kappa opioid, B2, and/or MC4 receptors which activation produced pronociceptive effect mediated by endogenous peptides, such as dynorphin, bradykinin and/or alpha MSH. Such analysis is to identify peptides and biochemical routes that are crucial in neuropathic pain development and maintenance. Further studies may test the possibilities of modulating the activity of pronociceptive peptides with their receptors' blockage, with an outlook to develop novel neuropathic pain therapies.

The work was supported by grant Maestro NCN 2012/06/A/NZ4/00028. J. Starnowska is a Ph.D. student and has a scholarship from the KNOW funds sponsored by Ministry of Science and Higher Education, Poland.

[B22]

Morphological features of astrocytic reaction after transient cerebral ischemia in the rat

Steliga A.¹, Lietzau G.², Waskow M.¹, Karwacki Z.³, Wojcik S.², Klejbor I.^{1,2}, Morys J.², Kowianski P.^{1,2}

¹Department of Health Sciences, Pomeranian University of Slupsk, Slupsk, Poland

²Department of Anatomy and Neurobiology, Medical University of Gdansk, Gdansk, Poland

³Department of Neuroanesthesiology, Medical University of Gdansk, Gdansk, Poland

Introduction: Cerebral ischemia is one of the most powerful pathological stimuli triggering reaction of glial cells, known as reactive astrogliosis. Fundamental features of this reaction depend on type, duration and severity of the pathological stimulus, as well as affected region in the brain [1]. Astrocytes are one of the most important cell populations of the central nervous system, generating the response to the damaging factor. Astrocytic reaction to ischemic stroke is predominantly related with restriction and isolation of the damaged tissue from the intact brain's structures, as well as ability to restore the other damaged cell populations.

Aim: The assessment of selected morphological features of reactive astrocytes, indicating their potential for division,

proliferation and reduction of tissue damage consequences was performed in the rat model of ischemic stroke.

Material and methods: Forty adult male Wistar rats were used in this study. Transient focal cerebral ischemia was evoked by placement of the surgical thread into the internal carotid artery and 1 hr occlusion of the middle cerebral artery. The postoperative survival period was up to 6 weeks. Immunocytochemical stainings for GFAP, Ki-67, and Pax6 were performed, with subsequent confocal microscopic study.

Results: Both studied transcription factors, namely Ki-67 and Pax6 were found in GFAP-immunoreactive (ir) astrocytes, localized within the damaged brain. The intensity of double-labeling for the studied factors increased gradually after the first week of reperfusion. Intense hypertrophic reaction developed. An increased GFAP-ir of the astrocytic fibers surrounding the lesion was observed.

Conclusions: Reassuring, our results concerning astroglial reaction indicate regaining of the morphological features, characteristic for the earlier developmental stages, by this cell lineage, as well as development of proliferative and division capacities in response to transient ischemia. The intense and multidirectional nature of the astrocytic response should be taken into account in future studies on the ischemic stroke.

References

1. Steliga A, Waśkow M, Karwacki Z, Wójcik S, Lietzau G, Klejbor I, Kowiański P. Transcription factor Pax6 is expressed by astroglia after transient brain ischemia in the rat model. *Folia Neuropathol* 2013; 51: 203-213.

[B23]

Sensitivity of glioma cell lines to modified isothiurea derivative (ZKK-3) under anoxia conditions

Stepien K.¹, Ostrowski R.P.¹, Pucko E.B.¹, Kazimierzuk Z.², Matyja E.¹

¹Department of Experimental and Clinical Neuropathology, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Institute of Chemistry, Warsaw Life Sciences University, Warsaw, Poland

Introduction: Gliomas are the most common primary brain tumours. Despite the improvement of the surgical techniques, radio- and chemotherapy, the treatment of malignant gliomas remains non effective. Hypoxia is

thought to play the major role in treatment failure of glial tumours. The aim of this study was to examine how an anoxic gas application impacts on the growth and proliferation of glioma-derived cell lines treated with selected cytostatic agent with potential anti-tumour properties.

Material and methods: The experiment was performed on human glioblastoma T98G cell line (WHO grade IV), cell line derived from low-grade paediatric brain tumour – subependymal giant cell astrocytoma (SEGA, WHO grade I) and normal human astrocytes (HA). For the experiment, the modified isothiurea derivative, N,N'-dimethyl-5-(2,3,4,5,6-pentabromobenzyl)-isothiouronium bromide (ZKK-3), was chosen. Cells were cultured in a medium supplemented with ZKK-3 (25 and 50 µM) or without the tested agent. Cells were divided into two experimental groups. One of them was subjected to 24 hours oxygen deprivation after ZKK-3 application, while the second one was additionally exposed to 18 hours of anoxia pretreatment (N₂/CO₂ gas mixture with 95% N₂ and 5% CO₂). Control cultures were maintained in normoxic conditions. The viability of cells was assessed using Trypan blue dye exclusion. Cell proliferation was evaluated by hemocytometer and Multisizer 3 Beckman Coulter.

Results: Oxygen deprivation caused the decrease of viability of SEGA neoplastic cells and normal human astrocytes. However, there was no significant alteration in the number of cultured T98G cells between cells exposed to 24 hours of anoxia and those maintained in normoxic conditions. The reduction of T98G cell lines proliferation rate was observed only while 18 hours of anoxic gas pretreatment was applied. The anoxic conditions resulted in considerable decrease of sensitivity of T98G glioma cells to the low concentration of ZKK-3, especially in the group pretreated with anoxia. For 50 µM of tested compound, the impact of anoxia on T98G cell line viability was not observed.

Conclusions: Reduction of anti-tumour properties of ZKK-3 related with oxygen deprivation condition, observed for lowconcentration of tested agent, was probably associated with the drug resistance effect. However, the 24h-anoxia was insufficient to induce drug resistance at high 50 µM concentration of tested compound. It might suggest, that limited access to the oxygen slows down the metabolism of astroglial cells due to anaerobic respiration. This phenomenon probably results in decrease of proliferation rate in anoxia-treated cell lines. The results spotted in anoxia-treated T98G and SEGA cell line suggest that the survivability of neoplastic cells upon anoxia conditions might depend on tumour type/grading.

The research was supported by the KNOW-MMRC project and partially by the Foundation for the Development of Diagnostic and Therapy.

|B24|

An analysis of the relationship between anxiety and social rank

Strzelczuk M.¹, Krotewicz M.¹, Bajaj M.¹, Krotewicz M.K.²

¹Department of Neurobiology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland

²Department of General and Colorectal Surgery, Medical University of Lodz, Lodz, Poland

Social rank limits social space for conspecifics. This phenomenon occurs in human and non-human forms of social community. Social rank means that the individuals high in the social rank have more social opportunities compared to those low in the social hierarchy. Several studies revealed that maintenance of social position is stressogenic and altered action of physiological systems. In our previous studies we observed that the brain noradrenergic system is implicated in the control of social rank studied in a social competition test what indicates that social rank have important consequences for the hypothalamo-pituitary-adrenocortical (HPA) system activity of the conspecifics. It is known that the brain noradrenergic neurons activity is closely related to the control of the HPA system. The obtained data showed that the maintenance of the low social position was correlated with enhance of noradrenaline release. It indicates that the low position in social stratification was due to anxiety induced by the presence of a high ranking animal, winner of a competitive situation. One of the hypothesis suggests that the formation of social rank resulted in the increase of anxiety in low ranking people. Moreover, the anxiety and observed in subordinate people anhedonia lead to mood disorders and other pathologies. The present study was designed to examine 1) anxiety indices of conspecifics holding different social position and 2) correlations between social indices measured in the social competition test and concentration of corticosterone (CORT) in the blood plasma of rats. In the presented experiments two social competition models were used, with or without food deprivation. The statistical analysis of behavioural indices registered during anxiety test showed a lack of differences related to social stratification. However, visible differences were revealed in temporal dynamics of measured behavioural indices in relation to social position. Competition between conspecifics was associated with differences in CORT concentration accordingly to social rank of rats. Additionally, the analysis of social indices showed a positive correlation between CORT concentration and index of vertical behaviour of dominants.

|B25|

Effects of glutamate receptors antagonists on morphological and biochemical changes in the myelin of EAE rats

Sulkowski G.¹, Dabrowska-Bouta B.¹, Frontczak-Baniewicz M.², Struzynska L.¹

¹Laboratory of Pathoneurochemistry, Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Electron Microscopy Platform, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Experimental autoimmune encephalomyelitis (EAE) is an animal model useful in investigations of the pathomechanisms of multiple sclerosis (MS). MS is an inflammatory demyelinating disease of the central nervous system (CNS) that often affects young adults between 20-40 years of age. A characteristic sign of the disease is the damage and loss of oligodendrocytes and myelin sheaths in the white matter of the spinal cord and brain. So far the etiology of MS is unknown. However, presumably one of the mechanisms involved in the disease, might be excitotoxicity. The elevation of glutamate in cerebro-spinal fluid as well as changes in the expression of glutamate receptors (iGluRs and mGluRs) and excitatory amino acids transporters (EAATs) were observed in brains of MS patients/EAE animals. We tested the effect of glutamatergic receptors antagonists: NMDARs – amantadine and memantine and mGluRs G I – LY 367385 and MPEP, on the neurological condition of EAE rats and biochemical and morphological changes observed in brain's myelin during EAE course in rat brains. The electron microscopy studies showed disturbances in myelin ordered layers in EAE rats. A significant changes in the structure of oligodendrocytes, activation and proliferation of microglia and astroglia were also observed. Moreover, we investigated the levels of mRNA for characteristic myelin protein (myelin markers) such as: MOG, MAG, PLP, and CNP-ase, by real time PCR technique in EAE rats and after treatment with glutamate receptors antagonists. The therapy with some of glutamate receptors antagonists (amantadine and memantine) has shown protective effects in EAE rats. The results suggest that disturbances in progressive brain damage in EAE and mechanisms of myelin degradation may be associated with glutamate excitotoxicity.

This work was supported by the Polish MNiSW grant N N401 620038.

|B26|

Oligodendrocyte progenitors provide trophic support for ischemically injured nervous tissue

Sypecka J., Strojek L., Winiarska H., Sarnowska A.

NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

The main role of oligodendrocytes is to myelinate nerve fibers in the central nervous system. However their high importance for the functioning of nervous tissue is suspected to go beyond the process of myelinogenesis. To address this issue, we designed a study aimed at determining potential neuroprotective properties of oligodendrocytes and their progenitors. For this purpose, rat organotypic hippocampal slices were exposed to a brief oxygen and glucose deprivation (OGD) which allows mimicking an ischemic injury in vitro. Immediately after the OGD procedure, the hippocampal slices were co-cultured together with differentiating oligodendroglial progenitors derived from neonatal rat brains. The cell survival and proliferation rate in the injured slices were estimated by immunohistochemical methods. After confirming the beneficial influence of the neighboring oligodendrocytes on the injured tissue, a set of molecular and biochemical experiments was carried out with aim of determining the mechanism(s) of the observed profound neuroprotective effect. The obtained results revealed that the oligodendrocyte-derived BDNF is the major factor promoting neuron survival in OGD-subjected hippocampal slices, while SCF strongly promote the cell proliferation. Among the newly-born cells, neuroblasts and microglia could be found most frequently, pointing to both the neuroprotective and neuroimmunomodulatory role of differentiating oligodendrocytes. In conclusion, the presented study revealed that oligodendrocytes are able to secrete BDNF and other active factors, thus providing trophic support for the other neural cells. The presented study confirms the hypothesis concerning the complex functions of oligodendrocytes in the nervous tissue.

Supported by grant 0345/B/P01/2010/38.

|B27|

The influence of recombinant α -synuclein protein injection on the expression of selected inflammatory factors in the murine brain

Sznejder-Pacholek A., Joniec-Maciejak I., Wawer A., Schwenkgrub J., Mirowska-Guzel D.

Department of Experimental and Clinical Pharmacology, Medical University of Warsaw, Warsaw, Poland

The studies exploring possible reasons of the neurodegeneration in Parkinson's disease are still far from conclusion. However, it has been already confirmed that α -synuclein protein (ASN), together with other proteins, is the principal component of Lewy pathology and influences strongly the pathogenesis of Parkinson's disease. It was observed that the increased level of ASN protein caused the microglia response. Microglial activation leads to the increased expression of inflammatory genes in the central nervous system (CNS), and the infiltration of peripheral immune cells into the CNS. Recent evidence suggests that reactive microglia cells may actively participate in the damage of dopaminergic neurons. Therefore, we examined the potential role of recombinant monomer of ASN in the inflammatory reaction in murine model of PD, hoping that clarification of the exact role of ASN in the inflammation will help us to understand the molecular mechanism of injuries of dopaminergic cells. ASN was bilaterally administered into substantia nigra (SN) or striatum (ST). We investigated the effect of high ASN monomer concentrations on microglia response: ionized calcium-binding adapter molecule 1 (IBA1) expression by western blot method, as well as some pro-inflammatory cytokines expression (interleukin: IL-6, IL-12) by Real-Time PCR and the lymphocytes T (CD3+, CD8+, CD4+) infiltration in the ST and SN of mice. The infiltration of the lymphocytes was measured by immunohistochemical method. We observed an increase of activated microglia cells (IBA1), a strong infiltration of lymphocytes T (CD8+ and CD4+) and higher level of mRNA for pro-inflammatory cytokines (IL-6, IL-12) after bilateral injection of ASN.

Our study provided more evidence for the involvement of ASN in the inflammatory response in the CNS. However, further studies are required to reveal the detailed mechanism of the influence of ASN in neuro-inflammation in course of Parkinson's disease.

[B28]

Impact of triclosan on aryl hydrocarbon receptor-induced neurotoxicity and apoptosis in mouse neocortical neurons

Szychowski K.A., Wojtowicz A.K.

Department of Animal Biotechnology, Agricultural University of Cracow, Cracow, Poland

Triclosan (TCS) is an antimicrobial agent used extensively in personal care and sanitizing products, such as soaps, toothpaste, and hair products. Over the past half century TCS has been incorporated into ever growing numbers of medical products as well as in household items, such as plastic cutting boards, sports equipment, textiles and furniture. The TCS has been detected in water and sediments and also in human workplace and private houses dust in wide range of concentrations. A number of studies has shown presence of TCS in different human tissue such as blood, adipose tissue, liver, brain, breast milk and in urine. There are limited studies evaluating the toxicity of TCS and data concerning bioaccumulation of TCS in living organisms are inconsistent. Some evidence indicate that TCS can be cumulated in human nails but previous studies have demonstrated that TCS is also metabolized to sulfate and glucuronide by the liver and extracted in methylated form in urine. It is well known that TCS has a similar chemical structure to dioxin and in the presence of sunlight may be transformed to produce up to four dioxin compounds such as 2,8-DCDD, 2,3,7-TCDD, 1,2,8-TricDD, and 1,2,3,8-TCDD. Aryl hydrocarbon receptor (AhR) is involved in the cell response to the numerous xenobiotics especially to dioxin. It has been demonstrated that this receptor is involved in regulation of genes expression which control growth and differentiation of neurons, and plays a key role in their metabolism and adaptation. The apoptosis dependent on the activation of AhR has been observed in primary cells cultures of the neocortex, hippocampus and cerebellum. The aim of the present study was to investigate the involvement of AhR in the cytotoxic and proapoptotic effects of TCS in mouse neocortical neurons. The cultures of neocortical neurons were prepared from Swiss mouse embryos on 15/16 day of gestation. The cells were cultured in phenol red-free Neurobasal medium with B27 and glutamine. On 7th day *in vitro* AhR siRNA was applied to neurons for 12 h at 50 nM in antibiotic-free medium containing the siRNA transfection reagent interferin. After silencing process the experimental cell culture groups were treated with AhR agonist β -naphthoflavone (β NF) and TCS for 24 h. After silencing

procedure and exposure to β NF or TCS the release of the lactate dehydrogenase (LDH) in the culture medium and caspase-3 activity in cells was measured.

Our preliminary data demonstrated that in the presence of β NF as well as TCS the increase in neuronal apoptosis and LDH release into the culture medium were observed. In neurons undergoing silencing procedure in the presence of β NF no significant changes in cytotoxicity or induction of apoptosis were observed. Additionally, the cytotoxic as well as proapoptotic effect of TCS was partially reduced after silencing process. In summary, the presented study demonstrated that AhR is involved in TCS mechanism of action.

Support by statutory funds of Agricultural University in Cracow DS 3242/14.

[B29]

The impact of olanzapine on the dehydrogenase lactate release in cortical organotypic cultures – study in an animal model of schizophrenia

Trojan E., Slusarczyk J., Glombik K., Kurek A., Basta-Kaim A.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Introduction: Schizophrenia is a common and disabling psychiatric brain disorder of neurodevelopmental origin, in which both genetic and environmental factors appear to play an important role. Many data from epidemiological and clinical studies indicate that viral or bacterial infection during pregnancy may enhance the risk of schizophrenia in adulthood. Therefore among the animal models used to study the etiology of schizophrenia, the systemic administration of the bacterial endotoxin lipopolysaccharide (LPS) during pregnancy is often applied.

Aim: The aim of this study was to assess the effect of prenatal LPS administration on the lactate dehydrogenase (LDH) activity in cortical organotypic cultures in both – basal and inflammatory conditions. Moreover, we determined the role of antipsychotic drug – olanzapine administration on the above mentioned parameter and the possible mechanism of this drug action in cortical organotypic cultures obtained from control and prenatally LPS treated offspring.

Material and methods: Female Sprague-Dawley rats were paired with syngenic males. Pregnancy was confirmed

the next morning. LPS (at the dose of 1 mg/kg, Serotype 026:B6) was injected sc every second day from the 7th day of pregnancy till delivery. The cortical organotypic cultures were prepared from the brains isolated from 6-7 day old offspring. All biochemical analyses were performed after culture stabilization. The activity of lactate dehydrogenase was measured in the medium by colorimetric LDH assay. The tests were performed in cortical organotypic cultures obtained from control and prenatally LPS-treated newborns in both basal conditions and following additional endotoxin (100 ng/ml) stimulation. In the next set of experiments the influence of olanzapine (at a dose of 5 μ M) or/and selective phosphatidylinositol 3-kinase (PI3K) inhibitor – wortmannin (at the dose of 10 μ M) – administration was evaluated.

Results: Our data demonstrated that prenatal LPS administration during last two weeks of pregnancy increased LDH release from the cortical organotypic cultures. Moreover in the cortical cultures obtained from pups prenatally treated with LPS the potentiation of LDH release after acute LPS treatment with comparison to cultures of controls was observed. On the other hand acute administration of olanzapine resulted in a decrease of cell mortality. The involvement of PI3K/Akt pathway in the olanzapine action was confirmed by the use of wortmannin, an inhibitor of phosphatidylinositol 3-kinase which attenuated the protective olanzapine action.

Conclusions: This study provided evidence of the protective effect of olanzapine in cortical organotypic cultures in animal model of schizophrenia based on prenatal LPS administration. Furthermore, it may be postulated that inhibitory impact of this drug on the LDH release is mainly connected with enhancement of PI3k pathway.

This study was supported by statutory funds Institute of Pharmacology PAS.

[B30]

Increased concentration of α -synuclein alters metabolism and transport of dopamine in mice

Wawer A.¹, Szejder-Pacholek A.¹, Joniec-Maciejak I.¹, Schwenkgrub J.¹, Mirowska-Guzel D.^{1,2}

¹Department of Experimental and Clinical Pharmacology, Medical University, Warsaw, Poland

²Institute of Psychiatry and Neurology, Warsaw, Poland

Parkinson's disease (PD) is the second most common progressive neurodegenerative disorder after Alzheimer's disease. It is characterized by two major processes: a pro-

gressive loss of dopaminergic neurons and an accumulation of intraneuronal Lewy bodies containing misfolded fibrillar α -synuclein (ASN). Neurodegeneration occurs mainly in substantia nigra (SN), accompanied by a decreased concentration of dopamine (DA) and its metabolites in striatum (ST). Various etiological triggers have been linked to PD, including genetic mutations and environmental toxins, but the pathway that leads to neurodegeneration remains unknown.

The aim of our study was to examine the influence of recombinant monomers of ASN on dopaminergic system activity. 1-year-old male C57Bl mice were used in this study. ASN was bilaterally administered into SN or ST. Animals were sacrificed at 4 weeks following ASN administration.

To evaluate the influence of the ASN on the neurodegeneration concentrations of striatal DA and its metabolites were measured by high performance liquid chromatography (HPLC). The expression of dopamine transporter (DAT) was measured by western blot method.

We observed changes in dopamine metabolism after ASN monomers administration. Our study also showed statistically significant influence of ASN on dopamine reuptake via DAT.

Further research must be conducted to better understand the crucial role of ASN in the neurodegenerative process in PD.

[B31]

Adipokines and other factors connected with dementia

Wehr H., Bednarska-Makaruk M., Graban A.

Department of Genetics and I Department of Neurology, Institute of Psychiatry and Neurology, Warsaw, Poland

Several circulating factors manifesting activity in neurochemical events were determined in persons with dementia and compared with control subjects. The whole group comprised 327 persons: 207 females and 120 males, aged 72.7 ± 8.7 years. 68 of them were diagnosed as probable Alzheimer's disease (AD), 33 as dementia of vascular origin (VaD), 47 as mixed dementia (MD), 83 as mild cognitive impairment (MCI) and 96 were controls without cognitive impairment (C). The levels of several adipokines: adiponectin, leptin, resistin and of the brain derived neurotrophic factor (BDNF) and interleukin-6 (IL-6) were determined in blood serum.

As connection of dementia with carbohydrate metabolism disturbances is observed, therefore glucose levels, glucose tolerance and insulin levels were also determined.

Statistically significant higher adiponectin and IL-6 levels, as compared with controls and glucose intolerance, were stated in MD and higher resistin levels and glucose intolerance in VaD. Therefore these preliminary results suggest the existence of disturbances in such forms of dementia which are connected with vascular pathology.

[B32]

Purinergic P2 receptors activation mediates neuroblastoma SH-SY5Y cell death evoked by extracellular α -synuclein

Wilkaniec A., Cieslik M., Gassowska M., Lenkiewicz A.M., Czapski G.A., Adamczyk A.

Department of Cellular Signaling, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

The purinergic P2 receptors for adenosine 5'-triphosphate (ATP) are abundantly present in the central nervous system and play important roles in neurotransmission, neuromodulation, as well as in glial cells function. P2 receptors belong to two structurally distinct families, the G protein-coupled P2Y receptors and the ligand gated ion channels – P2X receptors. Recently, extracellular ATP and specific purinergic receptor subtypes have been shown to be involved in neurodegenerative disorders including Parkinson's and Alzheimer's diseases. However, till now the mechanisms underlying the disturbances in ATP-mediated neurotransmission are not well established. Recent studies indicated that in neurodegenerative disorders, α -synuclein (ASN) has been released and taken up by cells that may facilitate its progressive pathological spreading in the brain. Our own *in vitro* studies support the idea of trans-cellular spread of secreted ASN and showed its significant cytotoxic effect when applied extracellularly. Therefore, the aim of this study was to examine the role of purinergic P2 receptors and ATP in the neurotoxicity induced by extracellular ASN.

The experiments were performed in human SH-SY5Y neuroblastoma cells differentiated with the all-trans retinoic acid (ATRA) using immunochemical, spectrophotometric, radiochemical and spectrofluorometric methods. Our study showed that SH-SY5Y cells express P2X₄, 5, 7 and P2Y₁ receptors. We have investigated the effect of ASN on gene expression and protein level of all these receptors, but we did not observe any significant changes. However, exposure of SH-SY5Y cells to ASN (10 μ M) induced significant increase in cytosolic calcium load, that

was mainly caused by the activation of extracellular Ca²⁺ influx. The stimulation of P2 receptors by ATP (5 mM) was found to trigger the similar but more pronounced effect than ASN. Furthermore, we found that ASN-evoked Ca²⁺ influx was partially eliminated by the selective purinergic P2 receptors antagonist, PPADS (200 μ M). In order to clarify the molecular mechanisms mediating the activation of P2 receptors, we examined the effect of ASN on extracellular ATP concentration and found that ASN promoted the release of ATP from SH-SY5Y cells. Moreover, both exogenous ASN and ATP induced significant increase in SH-SY5Y neuroblastoma cells death by about 50%. The toxic effect of these compounds was significantly prevented by PPADS, indicating that activation of P2 receptors is involved in cytotoxicity evoked by extracellular ASN in the human neuroblastoma SHSY-5Y cell line. Taken together, these results raise the possibility that the P2 signaling pathway could be a therapeutic target for treating neurodegenerative disorders.

Supported by NCN grant 2013/09/D/NZ3/01359.

[B33]

Impact of retinoid X receptor on the apoptotic and toxic effects of dichlorodiphenyldichloroethylene (DDE) in mouse embryonic neuronal cells

Wnuk A., Rzemieniec J., Litwa E., Lason W., Kajta M.

Department of Experimental Neuroendocrinology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

One of the most toxic compounds is dichlorodiphenyldichloroethylene (DDE) which is metabolite of the pesticide dichlorodiphenyltrichloroethane (DDT). DDE may easily cross the skin and blood-brain barrier. It accumulates in human organisms, especially in adipose and brain tissues. Recent epidemiological data exhibited the association between exposure to pesticides and increased ratio of neural degeneration. Enhanced level of DDE was detected in brain and blood samples of people suffering from Alzheimer's disease. Available data suggest that DDE induces apoptosis in bird and murine neurons. Although neurotoxic effects of DDE have been demonstrated, the role of retinoid X receptor (RXR) in apoptotic and toxic actions of DDE needs to be clarified. This study aimed to provide evidence on involvement of RXR in DDE-induced

apoptosis and neurotoxicity in mouse neuronal cells in primary cultures.

Our study demonstrated that DDE (10 and 100 μ M) activated caspase-3 in the mouse neuronal cell cultures, which was found within 6 hours after the treatment and accompanied by significant increase in lactate dehydrogenase-release (LDH) at 6 and 24 h of experiment. These effects were not tissue-specific, i.e. hippocampal, neocortical and cerebellar cells responded to DDE in similar way. In addition, the apoptotic and neurotoxic effects of DDE were followed by approximately 2-fold increase in RXR protein level, as evidenced by ELISA and Western blot analyses. DDE-induced stimulation of RXR-signaling was verified using selective antagonist HX531 and specific siRNA. HX531 inhibited the effects evoked by DDE and reduced both caspase-3 and LDH release. Furthermore, in RXR siRNA-transfected cells DDE did not retain its apoptotic and toxic potency, thus suggesting a crucial role of the receptor in propagation of DDE-induced apoptosis and neurotoxicity.

This work was supported by the Operating Program of Innovative Economy 2007-2013, grant No. POIG.01.01.02-12-004/09.

[B34]

Contribution of tachykinin NK₁ and μ -opioid receptors to cardio-respiratory effects of the analgesic chimeric peptide in anaesthetized rats

Wojciechowski P.¹, Szereda-Przestaszewska M.¹, Lipkowski A.W.²

¹Laboratory of Respiratory Reflexes, Department of Experimental Pharmacology, Warsaw, Poland

²Department of Neuropeptides, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Clinical use of opioid analgesics is limited by a variety of side effects such as respiratory arrest, arterial hypotension, drug tolerance and dependence. Therefore, many attempts are being made to create an effective analgesic that is free from above-mentioned side effects. Recently synthesized peptide – AWL3106 – is composed of two pharmacophores: μ opioid receptor agonist – dermorphin, and tachykinin NK₁ receptor agonist – substance P₇₋₁₁. This novel peptide shows strong analgesic activity of 5 and 30 folds the potency of morphine after intravenous and intrathecal administration, respectively.

The purpose of the present study was to show the contribution of each pharmacophore to cardio-respiratory pattern evoked by AWL3106.

Cardiorespiratory parameters were measured in 27 anaesthetized spontaneously breathing rats. Animals were treated with an intravenous injection of AWL3106 (0.3 μ mol/kg) in the following scheme:

- in control conditions ($n = 9$) and after blockade of opioid receptors with naloxone (2 mg/kg) ($n = 9$),
- following simultaneous blockade of opioid (naloxone) and tachykinin NK₁ receptors with SR140333 (2 μ mol/kg, $n = 9$).

Control intravenous injection of AWL3106 induced an apnoea of mean duration of 5.1 ± 0.7 s ensued by breathing of 19% reduced frequency (f) and 22% augmented tidal volume (V_T) with no signs of ventilatory depression, as well as 59% fall in mean arterial pressure (MAP) measured 2 min post drug.

Naloxone injected 2 min prior to AWL3106 challenge prevented the occurrence of apnoea as well as significantly reduced the maximal fall in breathing rate and an increase in tidal volume compared to the control injection. Minute ventilation still remained unaffected. Opioid receptor blockade failed to prevent the decrease in MAP. Subsequent blockade of NK₁ receptors resulted in suppression of all cardiorespiratory effects induced by AWL3106.

The present study showed that activity of opioid pharmacophore was essential to induce apnoea, whereas simultaneous activation of opioid and tachykinin receptors with AWL3106 affected both components of respiratory pattern (V_T and f). Arterial hypotension was associated mainly with activation of tachykinin NK₁ receptors.

[B35]

Neurochemical changes underlying modified swim test as an animal model of negative symptoms of schizophrenia

Wozniak M., Wieronska J., Pilc A.

Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Modified swim test has been recently introduced as an animal model of negative symptoms of schizophrenia, especially the blunted effect. The model is based on the repeated treatment with antagonist of NMDA receptor – MK-801 (0.4 mg/kg, one a day for 13 days) which increases the immobility in the modified swim test 24 h after the final injection

of chosen compound. These behavioral deficit was reversed by the atypical antipsychotic risperidone (0.1 and 0.2 mg/kg), as well as by agonist of 5-HT_{1A} receptors – (RS) 8-OH-DPAT (0.01; 0.025 mg/kg). We also studied the activity of two positive allosteric modulators of GABA_B receptor: CGP7930 (1; 2.5; 5 mg/kg) and GS39783 (0.5; 1; 2 mg/kg) in the same model. The compounds were active at the doses of 5 mg/kg and 0.5 and 1 mg/kg, respectively. In the second part of the present study, with the use of western blotting analysis, we also investigated the changes in 5-HT_{1A} receptor and GAD65 and GAD67 level in PFC and hippocampus of MK-801-treated mice, as those proteins can be regarded as the markers of the psychosis. The increases in 5-HT_{1A} receptor level was observed both in the PFC and hippocampus, while decrease in GAD65 protein was observed in the hippocampus only. No changes were observed for GAD67 receptor level in either structure investigated. Our findings support the hypothesis that the changes in 5-HT_{1A} receptor and/or GABA level may contribute to the pathophysiology of schizophrenia.

[B36]

Role of the Ras in regulation of BNIP3 expression in glioblastoma cells

Wysokinska E., Kalas W., Pawlak A., Ziolo E., Strzadala L.

Ludwik Hirszfeld Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, Wrocław, Poland

It has been demonstrated that anticancer treatments are able to induce autophagy in cancer cells. The role of autophagy in cancer is not clear, as autophagy may promote cancer cell survival or became the mechanism of growth restriction. In glioma cells this process can be induced by anticancer agents such as arsenic trioxide, ceramide, temozolomide, quercetin or radiation.

Ras proteins, frequently mutated in human cancers, function as molecular switches regulating pathways responsible for proliferation and cell survival. Ras family proteins were shown to play a role in the autophagy process. For example, HRas is able to induce autophagy through ERK signaling pathway in human colorectal, breast and cervical cancer. In other models Ras induces autophagy by upregulating Atg5 or Noxa and MEK/ERK pathway. Bcl-2/adenovirus E1B-19 kDa-interacting protein 3 (BNIP3), likewise Ras protein, is responsible for the fate of cancer cells. BNIP3 is an important mediator of cell survival and a member of the Bcl-2 family of proteins that regulate programmed cell death and autophagy.

In our previous studies, we for the first time demonstrated correlation between Ras, BNIP3 and process of autophagy in human colorectal cancer cell line (DLD-1). We have showed the primary role of KRas in the regulation of BNIP3 protein expression, establishing a link between the expression of oncogenic HRas and upregulation of BNIP3 and their control of autophagy in cancer cells. Additionally we found that upregulation of BNIP3 in DLD-1 cells results in KRas-dependent resistance to 5-Fluorouracil.

Glioblastomas are the most aggressive form of brain tumor that show resistance to treatment with various chemotherapeutic agents and BNIP3 was shown to promote its survival and adaptation to hypoxia by autophagy. The purpose of our studies is to elucidate the role of Ras and BNIP3 in induction of autophagy in U373MG glioma cell line. Here we present the data on regulation of BNIP3 expression by Ras in glioblastoma cells in normoxia and hypoxia. We believe that these studies will reveal the new opportunities for treatment of glioblastoma.

References

1. Swiderek E, Kalas W, Wysokinska E, Pawlak A, Rak J, Strzadala L. The interplay between epigenetic silencing, oncogenic KRas and HIF-1 regulatory pathways in control of BNIP3 expression in human colorectal cancer cells. *Biochem Biophys Res Commun* 2013; 441: 707-712.
2. Kalas W, Swiderek E, Rapak A, Kopij M, Rak JW, Strzadala L. H-ras up-regulates expression of BNIP3. *Anticancer Res* 2011; 31: 2869-75

This work was supported by grant of NCN no. 2011/01/NZ4/00938.

[B37]

Effect of clonidine on kynurenic acid production in rat brain cortex

Zakrocka I., Turski W.A., Kocki T.

Department of Experimental and Clinical Pharmacology, Medical University of Lublin

Clonidine is a central acting antihypertensive agent. Its mechanism of action is connected with activation of presynaptic α_2 -adrenoreceptors and imidazoline receptors in the central nervous system. Clonidine may have also influence on glutamatergic neurotransmission, by attenuating effects caused by NMDA or AMPA receptors activation. Kynurenic acid (KYNA) is the only known endogenous antagonist of excitatory amino acid receptors and of

α 7 nicotinic receptors. Cerebral synthesis of KYNA from its bioprecursor L-kynurenine is catalyzed by aminotransferases localized preferentially within astrocytes. In the brain, two distinct kynurenine aminotransferases, KAT I and KAT II, are responsible for the formation of KYNA. KYNA was shown to display potent neuroprotective properties and its impaired production was implicated in the neurodegenerative diseases. Additionally, it was recently reported that KYNA could play a role in regulation of blood pressure. A significant reduction of blood pressure in animal model of hypertension has been achieved by local intracerebral administration of KYNA.

The aim of this study was to evaluate the effect of clonidine on the formation of KYNA in brain cortex in rat.

Experiments were carried out on brain cortex obtained from male Wistar rats. The effect of clonidine on kynurenic acid concentration was evaluated on cortical slices after 2 hours of incubation in different concentrations of tested drug. To determine kynurenine aminotransferases (KATs) I and II activities cortical tissue was homogenized and dialyzed for 12 hours. Homogenates were incubated for 2 hours with tested compound. Supernatant was applied on the cation-exchange resin (Dowex 50 W+). Eluted kynurenic acid was subjected to the HPLC and quantified fluorimetrically. The experiments were conducted at least three times to achieve comparable results. Experimental procedures have been approved by the Local Ethical Committee in Lublin and are in agreement with European Communities Council Directive on the use of animals in experimental studies.

Clonidine at the concentration of 1, 3 and 5 mM decreased KYNA production in the brain cortical slices to 80% ($p < 0.05$), 70% ($p < 0.001$) and 40% ($p < 0.001$) of control, respectively. whereas at the concentration of 0.5, 0.1 and 0.01 mM was ineffective. Clonidine at the concentration of 0.01-5 mM did not affect the activity of KAT's.

The data presented here indicate that clonidine might decrease cortical level of kynurenic acid and thus suggests the novel mechanism of action of clonidine.

Supported by the grant for young scientists from Ministry of Science and Higher Education, MNmb515/2013.

[B38]

Influence of losartan on kynurenic acid production in the rat brain cortex

Zakrocka I., Turski W.A., Kocki T.

Department of Experimental and Clinical Pharmacology, Medical University of Lublin, Lublin, Poland

A renin–angiotensin–aldosterone (RAA) system is an important regulator of body homeostasis. It is thought to regulate not only water – electrolyte metabolism, including the regulation of thirst, blood pressure maintenance, vasopressin release, but it is responsible for controlling thermoregulation, learning, memory, emotions, reproduction and sexual behaviors as well. Among angiotensin receptors, the most prominent role in pharmacotherapy possesses receptor type 1 (AT-1), RAA was also implicated in the development of neurodegenerative disorders. It was reported that activation of RAA in the brain may have important role in animal seizure models. Recent studies indicate that substances interfering with concentration and pharmacodynamic properties of angiotensin are proven to have neuroprotective effect, especially to ischemic nervous tissue, improving patient's prognosis after vascular brain damage.

Kynurenic acid (KYNA) is the only known endogenous antagonist of excitatory amino acid receptors and of α 7 nicotinic receptors. Brain KYNA displays high affinity towards the glycine site of N-methyl-D-aspartate (NMDA) receptor complex. The cerebral synthesis of KYNA from its bioprecursor L-kynurenine is catalyzed by two distinct kynurenine aminotransferases (KAT I and KAT II). The disturbances of KYNA production have been linked to the occurrence of epilepsy, Huntington's disease, Alzheimer's disease, schizophrenia, AIDS-related dementia and others. The anticonvulsant and neuroprotective role of KYNA in vivo and in vitro is well documented.

In this study we aimed to analyze whether losartan, an AT₁ – receptor antagonist, might modify KYNA production in brain cortex of a rat.

In cortical slices losartan at the concentration of 0.5, 1, 3 and 5 mM decreased KYNA synthesis to 61% ($p < 0.05$); 17% ($p < 0.001$) 6% ($p < 0.001$) and 5% ($p < 0.001$) of control value, respectively. The activity of KAT I was decreased by losartan at the concentration of 0.01; 0.1; 0.5; 1 and 3 mM to 45% ($p < 0.01$); 13% ($p < 0.001$); 12% ($p < 0.001$); 6% ($p < 0.001$); 3% ($p < 0.001$) of control value, respectively. The activity of KAT II was diminished by losartan at the concentration of 0,01; 0,1; 0,5; 1 and 3 mM to 74% ($p < 0.01$); 9% ($p < 0.001$); 6% ($p < 0.001$); 2% ($p < 0.001$); 1% ($p < 0.001$) of control, respectively.

Our data suggest that losartan might decrease kynurenic acid production in the rat brain cortex, most probably via influence on KAT's activity.

Supported by the grant from National Science Centre (NCN) PRELUDIUM 4 No UMO-2012/07/N/NZ4/02088.

[B39]

The role of oxidative stress in cytotoxicity evoked by tetrabromobisphenol a in primary cultures of rat cerebellar granule cells

Ziemska E., Lenart J., Lazarewicz J.W.

Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Tetrabromobisphenol A (TBBPA) and other brominated flame retardants (BRF) are additives to industrial products reducing their combustibility. It was found that TBBPA leaks to environment and contaminates samples collected from humans and wildlife. Recent papers showed cytotoxic effects of BRFs, however relatively few studies addressed the subject of TBBPA neurotoxicity and its mechanisms. Although there are indications that TPPBA-induced excitotoxicity, influx of extracellular Ca²⁺ and particularly its release from intracellular stores play a key role in the mechanism of TPPBA neurotoxicity, there is also data suggesting a primary role of oxidative stress in this process. The aim of this study was to evaluate the latter hypothesis. Experiments were performed using the primary cultures of rat cerebellar granule cells (CGC) at 7th day *in vitro*. To evaluate TBBPA neurotoxicity, the cells were exposed for 30 min to this BFR, and neuronal viability was assessed after 24 h, using fluorescence microscope and propidium iodide staining. During or after 30 min incubation with TBBPA in Locke medium, we estimated selected parameters of oxidative stress including reactive oxygen species (ROS) production and changes in mitochondrial membrane potential by measuring DCF and rhodamine fluorescence, respectively. Moreover, the intracellular glutathione (GSH) level was assessed using fluorescence kit, SOD level with Western blot and catalase activity using absorbance kit. Free radical scavengers GSH, vitamin C or N-acetyl cysteine (NAC), NMDA receptor antagonist MK801 and ryanodine with bastadin 10 inhibiting intracellular Ca²⁺ release *via* ryanodine receptors as well as a megachannel antagonist cyclosporine A (CsA) were used as pharmacological tools to test different mechanisms contributing to oxidative stress and

cytotoxicity. Viability of cells 24 h after 30-min incubation with 25 μM TBBPA decreased to 30% compared to control, while application of all the substances tested, except CsA, increased their survival to 60-80%. Incubation with 25 μM TBBPA increased ROS production by about 50% and significantly decreased the mitochondrial potential. Application of 1 mM GSH completely prevented the former effect but did not affect depolarization of mitochondria. Administration of 7.5 to 25 μM TBBPA dose dependently decreased intracellular GSH level to 40%, and catalase activity to 60% of the control. Application of ROS scavengers, bastadin with ryanodine and MK801 partially prevented these effects. In turn SOD-1 level increased in dose depend manner but SOD-2 declined after TBBPA administration compare to the control. Results of the presented experiments indicate that oxidative stress reflected by changes in ROS production, GSH/SOD level and catalase activity, plays a significant role in TBBPA cytotoxicity in CGC in concert with excitotoxicity and calcium imbalance, but not with activation of megachannels.

Supported by the NCN grant 2012/05/B/NZ7/03225.

[B40]

Bastadin 12 is a new tool useful in the study of mechanisms of intracellular calcium release evoked by thapsigargin and Tetrabromobisphenol A

Ziemska E.¹, Stafiej A.¹, Toczyłowska B.^{2,3}, Lazarewicz J.W.¹

¹Department of Neurochemistry, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland

³Nalecz Institute of Biocybernetics and Biomedical Engineering, Polish Academy of Sciences, Warsaw, Poland

Thapsigargin (Th) is the established Ca²⁺ releaser from intracellular ER stores. Studies of others using mainly skeletal SR membranes demonstrated that Th-induced Ca²⁺ efflux from SR can be inhibited by simultaneous administration of ryanodine (Ry) and the macrocyclic brominated tyrosine derivative bastadin 5 (Ba5) at micromolar concentrations but not by these compounds applied separately. These results have been interpreted as unmasking by Th of Ry-insensitive leak conformation of ryanodine receptors (RyR) which are converted by Ba5 into Ry-sensitive conformation. We previously confirmed these properties of Ba5

using the rat cerebellar granule cell (CGC) in primary cultures. The effects of other bastadin derivative Ba12 on Ca²⁺ homeostasis were never systematically studied. Tetrabromobisphenol A (TBBPA) is a commonly used brominated flame retardant with the recognized neuro- and cytotoxic properties. It has been suggested that release of Ca²⁺ from the intracellular stores mediates TBBPA cytotoxicity. In the present experiments, using CGC and the fluorescent intracellular Ca²⁺ probe fluo-3, we evaluated usefulness of the synthetic Ba12 as a tool that could replace Ba5 in the comparative study characterizing intracellular Ca²⁺ release evoked by thapsigargin and TBBPA. The results demonstrated that 200 nM Th as well as 30 μM Ba12 applied alone induces increase in the intracellular Ca²⁺ level in CGC, while 2.5 and 10 μM Ba12 did not interfere with basal Ca²⁺ concentration. Both 2.5 μM Ba12 or 200 μM Ry administered separately only partially reduced the release of Ca²⁺ induced by Th, whereas these compounds in combination completely inhibited this effect. Ca²⁺ transients in CGC evoked by the muscarinic receptor agonist, 100 μM oxotremorine-M, were significantly inhibited by antagonist of IP3 receptors 100 μM 2-APB, but not by Ba12 and Ry given either separately or together. TBBPA (5, 10 and 25 μM) concentration-dependently increased intracellular Ca²⁺ level in CGC. Administration of Ba12 in combination with Ry completely blocked the increase in intracellular Ca²⁺ concentration evoked by 10 μM TBBPA, while these compounds applied separately were ineffective. In conclusion, these results show that Ba12 can replace Ba5 in testing sensitivity to Ry of Ca²⁺ release from the intracellular stores of cultured neurons and in identifying leak conformation of RyR. They are also consistent with the hypothesis that TBBPA and Th induce release of intracellular Ca²⁺ through a common mechanism involving leak conformation of RyR.

Supported by the Polish National Science Centre grant # 2012/05/B/NZ7/03225.

|B41|

The neuroprotective effect of histone deacetylase inhibitor – sodium butyrate after experimental neonatal hypoxia-ischemia

Ziemka-Nalecz M., Jaworska J., Sypecka J., Zalewska T.

NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

Birth asphyxia remains a frequent cause of perinatal morbidity and mortality. During perinatal hypoxic-ischemic

(HI) brain injury neuronal cells are damaged and lose their functions or die. In the past few years, it has become clear that brain ischemia stimulates neural stem cell proliferation and differentiation in the cerebral neurogenic area – the dentate gyrus (DG) of the hippocampus, which is frequently injured after the perinatal HI. Induction of neural progenitors after HI insult may represent an endogenous mechanism for brain regeneration. Mounting evidence indicates that histone deacetylase inhibitors (HDACi) exert stimulation of endogenous neurogenesis in adult brain. Our aim in this study was to examine the influence of HDACi – Sodium Butyrate (SB) on the generation of neurons and oligodendrocytes in immature brain subjected to hypoxia-ischemia. Furthermore, we tried to elucidate the mechanisms of action of Sodium Butyrate. We utilized an established model of hypoxia-ischemia induced in rats of postnatal day 7 (PND7). After ligation of the left common carotid artery the animals were exposed to hypoxia (7.6% oxygen for 60 min). Sham-operated rats were used as controls. SB (300 mg/kg) was injected subcutaneously for 5 consecutive days starting immediately after HI. To estimate the proliferation profile and phenotype of newborn cells, animals were injected with BrdU (50 mg/kg). 3-14 days after HI the presence of BrdU-positive cells was seen in both hemispheres (ipsi- and contra-lateral) with the strongest proliferation level occurring 3-6 days after the injury. To confirm that BrdU-positive cells represent newly generated neuroblasts and oligodendrocyte progenitors, we used double staining BrdU/DCX and NG2/BrdU, respectively. We found that administration of SB stimulates the generation of neuroblasts and progenitors of oligodendrocytes, and leads to reduced volume of brain lesion caused by HI insult. We also evaluated if neuroprotective effect was associated with increasing acetylation of histon H3 and with changes in the expression of some proteins which control apoptosis.

Supported by NSC grant 2012/05/B/NZ3/00436.

|B42|

Lesions of noradrenergic neurons with DSP-4 inhibit cell proliferation in the dentate gyrus of the hippocampus but do not damage the progenitor cells in adult mice

Zietek M., Swiergiel A.H.

Department of Animal and Human Physiology, University of Gdansk, Gdansk, Poland

DSP-4 is a highly selective neurotoxin of central and peripheral noradrenergic neurons. DSP-4 strongly inter-

acts with the noradrenaline uptake mechanism at the level of the neuronal membrane and permanent degeneration of noradrenergic terminals, though the precise mechanism of its action has not yet been fully elucidated. Several investigators reported that the neurotoxic effects of DSP-4 are more extensive in the region innervated by axons of the locus coeruleus (LC). Moreover, noradrenergic cells from the LC degenerate in Parkinson's disease. We assessed the number of proliferating cells in the dentate gyrus of the hippocampus of adult mice after damage to the projection from the LC as a result of a single administration of DSP-4.

Male Swiss mice (6-7 weeks) were divided into two groups that were examined 4 and 16 days after the administration of DSP-4. In the first group, the brains were dissected four days after a single injection of DSP-4 (50 mg/kg, 0.1 ml, *i.p.*, $n = 4$) or saline (0.1 ml, $n = 8$) and investigated by immunohistochemistry. Cell proliferation was determined, using 5-bromo-2'-deoxyuridine, which is combined with the newly synthesized DNA. The brains of the second group of animals, sixteen days after the administration of DSP-4 (50 mg/kg, 0.1 ml, *i.p.*, $n = 4$) or saline (0.1 ml, $n = 8$), were also analyzed immunohistochemically. The immunohistochemical studies in group I showed significant differences between the experimental and the control animals. Four days after administration of DSP-4 we observed decreased number of proliferating cells in the experimental groups in all examined section as compared to the control groups. In group II, 16 days after administration of DSP-4 an increased number of proliferating cells was found in all experimental groups compared to controls. Unfortunately, no statistically significant differences were confirmed. The results suggest that the lesion of noradrenergic neurons caused by DSP-4 does not damage the progenitor cells which produce granule neurons in the hippocampus.

[B43]

Neuronal and glial progenitors obtained from human iPS cells on 3D modified collagen scaffolds – the tool for the neurodevelopmental and neurotoxicity studies

Zychowicz M.¹, Pietrucha K.², Podobinska M.¹, Augustyniak J.¹, Winiarska H.¹, Buzanska L.¹

¹NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

²Lodz University of Technology, Lodz, Poland

Since the induced pluripotent stem cells (iPSC) can be obtained from the adults, the regenerative medicine can be supported by new tools comprising autologous iPSC-derived cells. The neurodegenerative diseases are in the scope of current research based on the iPSC cells technology, while biomaterials combined with the iPSC derived cells can have a great potential to fulfill the demands of neuroregeneration of CNS.

Here we describe the possibility of neural commitment and differentiation of iPSC as well as HUCB-NSC cells seeded onto the chemically modified collagen scaffolds in order to establish and standardize the *in vitro* “biomimetic” culture system.

Shape-sponge scaffolds were prepared from the porcine collagen dispersion using a lyophilization technique. This process produces a highly three-dimensional (3D) porous scaffold. Due to relatively weak mechanical and thermal properties the collagen sponges were cross-linked using carbodiimide. This way is preferable for modification due to that “zero length” cross-linking chemical catalyst non-incorporation within structure of matrix, and scaffolds exhibit no cytotoxicity.

The iPSC cells obtained by non integrative, virus free methods were expanded in the feeder free culture and differentiated toward neural lineage. The obtained neural stem cells, as well as more differentiated neuronal/glial cell population, were seeded on the collagen modified scaffolds. The RT PCR data have revealed gradual disappearance of pluripotency markers (*Oct4*, *Sox2*, *Nanog*) during the differentiation process and acquiring the neural markers, such as *Nestin*, *Pax6*, *GFAP*, *Map-2*, *Neurogenin*.

The undifferentiated, pluripotent stem cells seeded on the chemically modified, collagen scaffolds revealed limited viability and did not differentiate spontaneously. In contrast iPSC derived neural stem cells and more differentiated neural progenitors survived, adhered in a well-defined flattened manner, and after a few days co-culture on the scaffolds expressed neuronal (*Nestin*, *Beta-Tubulin III*,

MAP-2, Doublecortin) and glial (A2B5, NG2, PDGFR α , GFAP) markers.

Moreover, the modified collagen scaffold can serve as a stem cell niche surrogate and can be used for toxicology studies. The HUCB-NSC cells seeded on these scaffolds were more resistant to the methylmercury treatment compared to the two dimensional culture conditions.

Our results indicate that collagen-based 3D microenvironment preferentially support neurally committed cells as compared to induced pluripotent stem cell, and may serve as an in vitro platform for further medical, pharmacological and toxicological applications.

The work is supported by statutory funds to MMRC and National Science Centre via Grant No DEC-2011/03/B/ST8/05867 and EIT+ BioMed 5.4 Project.

[B44]

The role of CXCL1 and CXCL5 in nociceptive transmission in a mouse model of diabetic neuropathy

Zychowska M., Rojewska E., Pilat D., Mika J.

Department of Pain Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Cracow, Poland

Diabetic neuropathy is the most common complication of diabetes mellitus, however its mechanism is still poorly understood. It seems that the chemokines may play a key role. It was shown that increased serum levels of CXCL1 and CXCL5 are linked to obesity, hyperglycemia, and impaired islets of Langerhans function. The goal of our study was to examine how the level of two chemokines CXCL1 and CXCL5 varies in the course of diabetic neuropathy and how their administration influences nociceptive transmission. All experiments were performed on Albino Swiss mice. Diabetes type 1 model was obtained by single intraperitoneal injection of streptozotocin (STZ; 200 mg/kg; *i.p.*). To verify the development of hyperglycemia, the blood glucose concentration and body weight were measured. In order to evaluate allodynia and hyperalgesia, von Frey test and cold plate tests, respectively, were used. The changes in mRNA amount and protein level of CXCL1 and CXCL5 in lumbar spinal cord were estimated by qRT-PCR and Antibody Array technic, respectively. The influence of single intrathecal (*i.t.*) administration of CXCL1 and CXCL5 on nociceptive threshold were measured using the tail flick, von Frey and cold plate tests, after 1, 4 and

24 hours. Both chemokines were administered in doses 10 ng, 100 ng and 500 ng/5 ml. The experiments were carried out according to the Institute's Animal Research Bioethics Committee and in accordance with IASP rules (Zimmermann, 1983). We observed that the STZ injection in mice increased plasma glucose and decreased body weight, along with inducing significant allodynia and hyperalgesia measured seven days after administration. No changes of CXCL1 and CXCL5 mRNA level were observed, in contrast upregulation of both at protein level was detected at day seven after STZ-treatment. The *i.t.* administration of CXCL1 and CXCL5 shown their pronociceptive properties. Furthermore, both contributed to the development of allodynia and hyperalgesia, however after 24 h only mechanical allodynia was still observed. Summing up, these findings indicate that CXCL1 and CXCL5 play an important role in the diabetic neuropathic pain development, which requires further research.

Supported by the NCN-grants 2012/05/N/NZ4/02416, 2011/03/B/NZ4/00042 and statutory funds. Magdalena Zychowska, a Ph.D. student, is a scholarship holder from the KNOW sponsored by Ministry of Science and Higher Education, Republic of Poland.

[B45]

LPS injections in young rats affect reactive gliosis following seizures evoked in adulthood

Kosonowska E., Kielbinski M., Gzielo-Jurek K., Janeczko K., Setkowicz Z.

Department of Neuroanatomy, Institute of Zoology, Jagiellonian University, Cracow, Poland

Aims: An increased activity of the immune system during an ongoing inflammatory process has adverse effects on the nervous system – it disturbs homeostasis of neural tissue and hence causes activation of two types of glial cells (reactive gliosis): astrocytes and microglial cells. Considering recent scientific reports, prior inflammation may change susceptibility to epileptic seizures. However, there is still not enough studies which indicate which cells and processes it can activate. Lipopolysaccharide (LPS) injection results in a massive antimicrobial defense reaction with an acute excessive activation of microglia. Some studies suggested that activated microglia may play a neuroprotective role by facilitating reparatory and regenerative processes. However microglia may also ini-

tiate and exacerbate neuropathological changes through secreting proinflammatory and cytotoxic molecules. Astrocytes may contribute to brain damage in a similar way but they also produce neuroprotective molecules and exert its neuroprotective actions on the neighbouring neurons. This issues could be important in case of studying the effects of systemic inflammation on seizure susceptibility. The aim of this study was to examine the long term effect of systemic inflammation induced on different postnatal developmental stages on activation of glial cells (astrocytes and microglial cells) in different experimental groups (with or without induction of status epilepticus).

Methods: LPS solution was injected intraperitoneally (2 mg/kg b.w.) to Wistar rats on postnatal day 6 or 30. When became two-month-old, the rats which survived inflammation were injected with pilocarpine to evoke status epilepticus and sacrificed 3 days thereafter. Brain sections were then processed for GFAP and Iba-1 immunohistochemistry to evaluate the area of reactive gliosis in the hippocampus using ImageJ software (*Area Fraction measurement*).

Results & Conclusions: Unexpectedly, the obtained results showed that status epilepticus induced in the adult rats evoked statistically significant reductions of Iba-1- and GFAP-immunopositive areas within the hippocampal formation. However, peritoneal LPS injections performed on the postnatal day 6 could prevent such seizure-induced decreases both in Iba-1 and GFAP tissue immunoreactivity. Following LPS injections on the postnatal day 30 significant decreases were limited to GFAP immunoreactivity only. The detected changes appear to reflect modification in reactive morphological transformation of their cellular substrates, i.e. astroglia and microglia involved in formation of seizure-genic gliosis in the hippocampal formation. Therefore, further detailed investigations are needed to elucidate relations between the reactive phenomena and effects of LPS proinflammatory stimulation at early stages of postnatal development on glial responses to seizures evoked in adulthood.

Supported by NCS GRANT: UMO-2012/05/B/NZ4/02406.

|B46|

The high cervical spinal cord injury and unilateral phrenic nerve ligation strongly reduce sGC/cGMP signaling in lower respiratory pathway followed by the failure of EMG activity in diaphragm

Lukacova N.¹, Hricova L.¹, Gedrova S.¹, Galik J.¹, Pavel J.¹, Sulla I.¹, Stropkowska A.¹, Zavodska M.¹, Oroszova Z.¹, Chalimoniuk M.², Langfort J.²

¹Institute of Neurobiology, Slovak Academy of Sciences, Kosice, Slovakia

²Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw, Poland

The high cervical spinal cord injury leads very often to the interruption of descending bulbospinal respiratory pathway, one of the vital pathways in CNS, which begins in brainstem, innervates phrenic motor neurons located at the C3-C6 cervical level and controls the diaphragm. Such injury leads to breathing rhythm abnormalities and life-threatening weakness of respiratory function. We previously demonstrated NO/soluble guanylyl cyclase (sGC) signalization in upper bulbospinal respiratory pathway. This study is focused on sGC/cGMP signalization in lower respiratory pathway after i) unilateral SCI performed at C2-C3 level, and ii) unilateral phrenic nerve ligation. The phrenic motoneurons within the phrenic nucleus in cervical spinal cord have been identified with retrograde tracer Fluorogold (FG). Two days after FG injection into hemidiaphragm, the ipsilateral phrenic motoneuronal pool was labeled within C3-C5 segments, with the greatest density found at C4 level. The increased intensity of sGCβ1 staining was detected 8 days after phrenic nerve ligation in phrenic motoneurons (C3-C5 segments) on the side of the injury. In addition, the phrenic nerve ligation strongly reduced the level of sGCβ1 in efferent respiratory pathway (diaphragmatic nerve) just below the nerve ligation, what correlates with the alterations of cGMP level in diaphragm. Although we found strong reduction of the cGMP level in both the contra- and ipsilateral hemidiaphragm, the decrease was much stronger on the side of the ligation. The unilateral interruption of the high cervical spinal cord had less impact on the cGMP synthesis in the diaphragm (mostly on contralateral side). It seems that the phrenic nerve activity was after cervical hemisection limited, but not so dramatically than after the phrenic nerve ligation. We assume that pre-phrenic interneurons located between the brainstem and the spinal cord may amplify the phrenic output after high cervical hemisection on contralateral

side. Moreover, the results clearly show remarkable unilateral silencing of the EMG activity after the phrenic nerve ligation and/or C2-C3 hemisection.

Supported by VEGA Grants No. 2/0173/14, 2/0191/13, 2/0187/13 and CE-NOREG SAS.

Editorial note:

The abstracts are printed in the form accepted by the Scientific Programme Committee of the Symposium.