

REVIEW PAPER

ARTYKUŁ PRZEGLĄDOWY

**DETERMINANTS OF MENTAL HEALTH DISORDERS IN THE COMMUNITY
DURING THE COVID-19 PANDEMIC: A LITERATURE REVIEW**

**DETERMINANTY ZABURZEŃ ZDROWIA PSYCHICZNEGO W SPOŁECZNOŚCI
PODCZAS PANDEMII COVID-19: PRZEGLĄD LITERATURY**

Yoyok Beki Prasetyo^{1(A,B,C,D,E,F)}, **Cindy Velinda**^{2(A,B,D,F)}, **Yulis Setiya Dewi**^{3(A,E,F)}

¹Department of Community Nursing, University of Muhammadiyah Malang, Indonesia

²Student of Nursing, Faculty of Health Sciences, University of Muhammadiyah Malang, Indonesia

³Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

Prasetyo YB, Velinda C, Dewi YS. Determinants of mental health disorders in the community during the COVID-19 pandemic: a literature review. Health Prob Civil. <https://doi.org/10.5114/hpc.2024.134772>

Tables: 3

Figures: 1

References: 37

Submitted: 2023 Oct 17

Accepted: 2024 Jan 23

Address for correspondence / Adres korespondencyjny: Yoyok Bektı Prasetyo, Department Community Nursing, University of Muhammadiyah Malang, Jl. Bendungan Sutami No. 188-A, 65145 Malang, East Java, Indonesia, e-mail: yoyok@umm.ac.id, phone: +62 0341 551149
ORCID: Yoyok Bektı Prasetyo <https://orcid.org/0000-0001-8801-7760>, Yulis Setiya Dewi <https://orcid.org/0000-0003-4407-0433>

Copyright: © John Paul II University in Biała Podlaska, Yoyok Bektı Prasetyo, Cindy Velinda, Yulis Setiya Dewi. This is an Open Access journal, all articles are distributed under the terms of the Creative Commons Attribution-NonCommercialShareAlike 4.0 International (CC BY-NC-SA 4.0) License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited and states its license.

Summary

This study aimed to identify the determinant factors to mental health disorders in the community during the COVID-19 pandemic. The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guide was used in this study. The article search was conducted through PubMed, Proquest, Cambridge University Press, and SpringerLink by using the keywords “Determinant Factors” and “Mental Disorder” or “Mental Health” and COVID-19” or “Corona Virus Diseases 2019”. The articles were selected from the studies published from 2019 to 2021. According to the research results, younger people are three times more likely to experience mental disorders than older people. Women tend to experience mental disorders twice as much as men. People who are afraid of being infected are three times more likely to experience mental health problems than people who are not afraid of being infected. People with chronic diseases tend to experience health problems twice as much as people who do not have chronic diseases (OR 1.30-1.84). The results of this literature review can help nurses

determine and control factors that affect mental health in the community during the COVID-19 pandemic and to achieve good outcomes in order to prevent more serious conditions due to COVID-19.

Keywords: sociodemographic factors, COVID-19, determinants, mental disorder, mental health

Streszczenie

Niniejsze badanie miało na celu identyfikację czynników determinujących, które powodują zaburzenia zdrowia psychicznego w społeczności podczas pandemii COVID-19. Wytyczne *Preferred Reporting Items for Systematic Reviews and Meta-analyses* (PRISMA) zostały wykorzystane w badaniu. Przeprowadzono wyszukiwanie literatury w bazach PubMed, Proquest, Cambridge University Press i SpringerLink, używając słów kluczowych „Determinant Factors” i „Mental Disorder” lub „Mental Health” oraz „COVID-19” lub „Corona Virus Diseases 2019”. Do przeglądu wybrano artykuły opublikowane w latach 2019-2021. Wyniki badań wskazują, że osoby młodsze są trzykrotnie bardziej narażone na zaburzenia psychiczne niż osoby starsze. Kobiety mają tendencję do doświadczania zaburzeń psychicznych dwa razy częściej niż mężczyźni. Osoby obawiające się zarażenia są trzykrotnie bardziej narażone na problemy ze zdrowiem psychicznym niż osoby, które nie obawiają się zarażenia. Osoby cierpiące na choroby przewlekłe doświadczają problemów zdrowotnych dwa razy częściej niż osoby, które nie cierpią na choroby przewlekłe (OR 1,30-1,84). Wyniki niniejszego przeglądu literatury mogą pomóc personelowi pielęgniarstwu w określeniu i kontrolowaniu czynników wpływających na zdrowie psychiczne w społeczności podczas pandemii COVID-19 oraz w uzyskaniu dobrych wyników i zapobieganiu powikłaniom związanym z COVID-19.

Słowa kluczowe: czynniki socjodemograficzne, COVID-19, determinanty, zaburzenia psychiczne, zdrowie psychiczne

Introduction

COVID-19 is a new type of virus that first appeared in Wuhan, Hubei Province, China, and has spread worldwide [1]. This virus attacks the respiratory tract caused by infection with the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) [2]. The World Health Organization (WHO) has declared the novel coronavirus (COVID-19) to be a global pandemic [3]. The impact of COVID-19 is not only on the physical aspect but also the mental aspect of health (such as stress, depression, and anxiety). During the COVID-19 pandemic, people have been more prone to stress, depression, and anxiety.

A large proportion of people in various countries reported that their mental health was adversely affected during the first wave of the pandemic [4]. The prevalence of distress was reported to have increased during the second wave in autumn and winter 2020, which coincided with the re-tightening of restrictions to contain the pandemic. The third pandemic wave in early 2021 (February to May) was less severe than the first and second waves in terms of hospitalizations and deaths [5].

The prevalence of stress, depression, and anxiety as a result of COVID-19 is 29.6%, 31.9%, and 31.9%, respectively [6]. In studies carried out in China in the early stages of the COVID-19 outbreak, the prevalence of severe psychological disorders was 53.8%, moderate-severe anxiety symptoms were reported by 28.8% of the population, moderate-severe depression symptoms were 16.5%, and moderate-severe stress was 8.1% [7]. In Arab studies, 23.6% of the general public reported psychological effects, 28.3% reported depressive symptoms, 24% reported anxiety, and 22.3% reported stress [8]. In studies conducted in Indonesia, as many as 63% of the people experienced anxiety, and as many as 66% reported

depression [9]. Early on in the pandemic, research from China, the US, and Europe demonstrated that mental health deteriorated. For example, the UK had a more significant increase in psychological discomfort than was experienced in prior upward trends, with the prevalence rising from 18.9% in 2018-2019 to 27.3% in April 2020, one month into lockdown [5].

There is still little research to identify the determinants of mental health disorders due to COVID-19 in Indonesia. Research conducted by Ilpaj et al. [10] indicated that the mental health impacts of COVID-19 included fear and anxiety, changes in sleeping and eating patterns, feelings of depression and difficulty concentrating, boredom and stress, drug and alcohol abuse, and the emergence of psychotic disorders. Other research shows that COVID-19 causes moderate to severe stress, anxiety, and depression [11].

The impact caused during the COVID-19 pandemic has led to various problems such as economic instability and unemployment. This can bring about an increase in the incidence of stress and anxiety. Identifying the determinants of mental health disorders during the pandemic is very important. These determining factors include age, gender, level of education, experience related to the SARS virus, and fear of infection [12]. Individuals can maintain mental health by identifying the determinants of mental health disorders. Efforts can be made by overcoming anxiety, stress, and depression, always thinking positively, sorting and choosing information related to the COVID-19 pandemic, and fostering good relationships with family and others.

Aim of the work

This study aims to identify the determinant factors that have been causing mental health disorders in the community during the COVID-19 pandemic.

ONLINE FIRST

Methods

Search strategy

The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guide was used in this study. The research was conducted for the articles by utilizing the PubMed, Proquest, Cambridge University Press, and SpringerLink by using the keywords “Determinant Factors”, “Mental Disorder”, “Mental Health”, “COVID-19” and “Corona Virus Diseases 2019” (Table 1).

Table 1. Search list

Number of articles found in each database	Keywords
PubMed = 92 Proquest = 443 Cambridge University = 324 SpringerLink = 213	Determinant factor AND Mental disorders or mental health AND COVID-19 or Corona Virus Diseases 2019

Study selection

Selections were chosen from research articles published since 2020. The articles focused on people affected by COVID-19 and by mental health problems, including people that remain at risk of mental health problems and people that have suffered from them. Excluded were articles that discuss the physical aspects of the impact of COVID-19 and articles written in languages other than English (Figure 1).




Figure 1. Study flow chart with PRISMA

Data extraction

Information from each study was categorized according to author/year, design/sample/country, primary findings, and study quality.

Literature review results

Study characteristics

The determinant factors that have affected people's mental health during the COVID-19 pandemic can be divided into two aspects, namely, sociodemographic and non-sociodemographic factors. There are 14 studies that state that sociodemographic determinants consist of age, gender, place of residence, education level, and income/economics: Bonsaksen et al. [13]; Guo et al. [14]; Kantor et al. [15]; Makhshvili et al. [16]; Mani et al. [17]; Mazza et al. [18]; Newby et al. [19]; Smith et al. [20]; Stanton et al. [21]; Venugopal et al. [22]; Verma et al. [23]; Cuiyan et al. [24]; Wong et al. [25]; and Zoghby et al. [26]. Ten studies describe non-sociodemographic determinants, namely: fear of infection, history of chronic diseases, working from home, family conflict, and lack of family support (Guo et al. [14], Mazza et al. [18], Newby et al. [19], Stanton et al. [21], Venugopal et al. [22], Wong et al. [25], Zoghby et al. [26], Choi et al. [27], Liu et al. [28], and Winkler et al. [29]) (Table 2).

Table 2. Characteristics of the study based on the determinants

Determinants	Studies
Sociodemographic factors:	Bonsaksen et al. [13]
Age	Guo et al. [14]
Gender	Kantor et al. [15]
Place of residence	Makhshvili et al. [16]
Education level	Mani et al. [17]
Income/economics	Mazza et al. [18]
	Newby et al. [19]
	Smith et al. [20]
	Stanton et al. [21]
	Venugopal et al. [22]
	Verma et al. [23]
	Cuiyan et al. [24]
	Wong et al. [25]

	Zoghby et al. [26]
Non-sociodemographic factors: Fear of infection History of chronic diseases Work from home Family conflict Lack of family support	Guo et al. [14]
	Mazza et al. [18]
	Newby et al. [19]
	Stanton et al. [21]
	Venugopal et al. [22]
	Wong et al. [25]
	Zoghby et al. [26]
	Choi et al. [27]
	Liu et al. [28]
	Winkler et al. [29]

Sociodemographic factors that influence mental health

Age, gender, place of residence, education level, and income / economic factors have affected people's mental health during the COVID-19 pandemic. It is argued that age is a determining factor in experiencing mental health problems during the COVID-19 pandemic (OR=-0.03-2.85) as younger people are three times more likely to experience mental disorders than older people [13,14,17,18,23,25]. Gender is another determinant of mental health problems during the COVID-19 pandemic (OR=0.60-1.70), with women being twice as likely to experience mental disorders as men [13,17,18,23,25]. A place of residence is also one of the determinants of mental health problems during the COVID-19 pandemic, on the grounds that people who live in urban areas are more likely to experience mental health problems than people who live in rural areas [15]. Another determining factor is an education level. The studies show that people with higher education are more likely to suffer mental health problems during the COVID-19 pandemic compared to people with lower education level (OR=0.50-1.26) [15,17,18,23,25]. Finally, people earning higher income are three times more likely to experience mental health problems during the COVID-19 pandemic than people with lower income (OR=0.70-2.95) [13-15,17,18,23] (Table 3).

Table 3. Determinant factors that affect mental health

Author, year	Design, sample, country	Main findings	Study quality*
Bonsaksen et al. 2020 [13]	Cross-sectional study 4,527 respondents Norway	Age (OR: 0.85, 95% CI: 0.80-0.90), gender (OR: 1.69, 95% CI: 1.32-2.16), social support (OR: 0.42, 95% CI: 0.43-0.52), economic concern (OR: 2.95, 95% CI: 2.51-3.57)	7/8 (87.5%)
Choi et al. 2020 [27]	Cross-sectional study 500 respondents Hong Kong	Fear of infection (OR: 2.20, 95% CI: 1.63-2.97), not having a mask (OR: 1.96, 95% CI: 1.50-2.56), unable to work from home (OR: 1.41, 95% CI: 1.13-1.77)	7/8 (87.5%)
Guo et al. 2020 [14]	Cross-sectional study 2,331 respondents China	Age (OR: 0.97, 95% CI: 0.778-1.197), income change (OR: 1.53, 95% CI: 1.23-1.84), chronic disease (OR: 1.84, 95% CI: 1.35-2.50), family conflict (OR: 1.98, 95% CI: 1.66-2.37)	7/8 (87.5%)
Kantor et al. 2020 [15]	Cross-sectional study 1,000 respondents United States	Gender (OR: 0.67, 95% CI: 0.51-0.89), place of residence (OR: 1.32, 95% CI: 0.98-1.79), religion (OR: 0.85, 95% CI: 0.63-1.14), occupation (OR: 1.61, 95% CI: 1.05-2.45), education (OR: 0.88, 95% CI: 0.24-3.17),	6/8 (75%)
Liu et al. 2020 [28]	Cross-sectional study 898 respondents United States	Fear of infection (OR: 2.87, 95% CI: 1.67-4.94), family support (OR: 0.46, 95% CI: 0.32-0.66)	7/8 (87.5%)
Makhashvili et al. 2020 [16]	Cross-sectional study 2,088 respondents Georgia	Age ($p<0.001$), economics ($p<0.001$), occupation ($p<0.001$)	7/8 (87.5%)
Mani et al, 2020 [17]	Cross-sectional study 922 respondents Iran	Age (OR: 2.88, 95% CI: 1.81-4.59), gender (OR: 1.74 95% CI: 1.31-2.31), marital status (OR: 1.35, 95% CI: 0.55-3.31), education (OR: 1.26, 95% CI: 0.75-2.12), economics (OR: 0.88, 95% CI: 0.61-1.28)	6/8 (75%)
Mazza et al, 2020 [18]	Cross-sectional study 2,812 respondents Italy	Age (OR: 0.99, 95% CI: 0.02-0.005), education (OR: 0.78, 95% CI: 0.355-0.123), gender (OR: 0.63, 95% CI: 1.893-0.445), occupation (OR: 0.719, 95% CI: 0.648-0.011), marital status (OR: 1.087, 95% CI: 0.837-1.003), family support (OR: 0.908, 95% CI: 0.405-0.210), work from home (OR: 1.050, 95% CI: 0.214- 0.313), chronic disease (OR: 1.473, 95% CI: 0.198 0.576)	7/8 (87.5%)
Newby et al. 2020 [19]	Cross-sectional study 2,812 respondents UK	Age ($p=0.00$), gender ($p=0.00$), education ($p=0.08$), occupation ($p=0.030$), chronic disease ($p=0.08$), fear of infection ($p=0.03$)	6/8 (75%)
Smith et al. 2020 [20]	Cross-sectional study 932 respondents	Age (OR: 0.83, 95% CI: 0.46-1.49), gender (OR: 2.21, 95% CI: 1.54-3.21), marital	6/8 (75%)

Author, year	Design, sample, country	Main findings	Study quality*
	UK	status (OR: 1.10, 95% CI: 0.76-1.60), family support (OR: 0.77, 95% CI: 0.51-1.15), economics (OR: 0.69, 95% CI: 0.40-1.18)	
Stanton et al. 2020 [21]	Cross-sectional study 1,491 respondents Australia	Age ($p \leq 0.0010$), Gender ($p=0.189$) marital status ($p \leq 0.001$), education ($p=0.002$), income ($p=0.047$), chronic disease ($p=0.001$)	7/8 (87.5%)
Venugopal et al. 2020 [22]	Cross-sectional study 453 respondents India	Age ($p=0.001$), gender ($p=0.055$), place of residence ($p=0.000$), marital status ($p=0.000$), education ($p=0.000$), occupation ($p=0.000$), family support ($p=0.000$), work from home ($p=0.039$)	7/8 (87.5%)
Verma et al. 2020 [23]	Cross-sectional study 354 respondents India	Age (OR: 0.746, 95% CI: 0.317-1.757), gender (OR: 1.26, 95% CI: 0.767-2.076), marital status (OR: 1.002, 95% CI: 0.452-2.224), income (OR: 1.1, 95% CI: 0.556-2.161), occupation (OR: 1.914, 95% CI: 0.072-3.418), education (OR: 1.16, 95% CI: 0.472-2.840)	7/8 (87.5%)
Cuiyan et al. 2020 [24]	Cross-sectional study 1,210 respondents China	Age ($p=0.006$), gender ($p=0.004$), marital status ($p=0.003$), family support ($p \leq 0.00$), occupation ($p=0.004$), education ($p=0.004$)	7/8 (87.5%)
Winkler et al. 2020 [29]	Cross-sectional study 3,021 respondents Czech Republic	Fear of infection (OR: 1.66, 95% CI: 1.38-1.99), economics (OR: 1.44, 95% CI: 1.21-1.71)	6/8 (75%)
Wong et al. 2020 [25]	Cohort study 263 respondents Hong Kong	Age (OR: -0.03, 95% CI: -0.08-0.02), gender (OR: 0.99, 95% CI: 0.30-1.68), education (OR: 0.50, 95% CI: -0.11-1.12), marital status (OR: 0.26, 95% CI: -0.38-0.90), family support (OR: 0.70, 95% CI: -0.16-1.56), chronic disease (OR: 0.38, 95% CI: -0.27-1.03)	9/11 (82%)
Zoghby et al. 2020 [26]	Cross-sectional study 510 respondents (age >18 years) Egypt	Age ($p=0.036$), gender ($p=0.003$), place of residence ($p=0.001$), maritime status ($p=0.740$), education ($p=0.699$), working in the medical field ($p=0.052$), and chronic disease ($p=0.011$)	7/8 (87.5%)

Note: * Joanna Briggs Institute (JBI) critical appraisal.

Non-sociodemographic factors that influence mental health

The studies show that people who are afraid of being infected are three times more likely to experience mental health problems than people who are not afraid of being infected (OR=1.66-2.87) [27-29]. People with chronic diseases tend to experience health problems twice as often as people who do not have chronic diseases (OR=1.3-1.84) [14,18,25]. Also someone who works from home is more prone to experience mental health problems compared to someone who works outside of home during the COVID-19 pandemic. Likewise, family conflicts are determinant to mental health problems in society during the COVID-19 pandemic. People with less family support tend to experience mental health problems more often during the COVID-19 pandemic (OR=0.04-0.90) [13,18,20,28,29].

Discussion of the review results

Sociodemographic factors that influence mental health

The research results show that young people were more likely to experience mental health disorders than older people. This is evidenced by previous research that shows that young adults (18-30 years old) are very vulnerable to experiencing stress because, at that age, they tend to get a lot of information from social media related to COVID-19 [30]. It has been proved that because of the fear that arises, any news or information related to COVID-19, whether verified or not, that is relayed through social media and television will increase panic and fear in the community [27].

The studies also demonstrate that women are more likely to experience mental health disorders than men. Women tend to be more prone to stress, anxiety, and depression, which can

also increase post-traumatic symptoms [31]. Also mothers who suffer miscarriages or experience partner violence are at high risk for developing mental health problems. Moreover, during the COVID-19 pandemic, infected pregnant women were giving birth without their husbands/partners accompanying them, and the babies were immediately separated from the mothers to avoid the COVID-19 infection. This has also brought about profound mental health disorders in mothers both in short-term and long-term [31].

The results of the research indicate that people living in urban areas are more likely to experience mental health problems during the COVID-19 pandemic than people from rural areas. Life in an urban area with a dense population affords less opportunity for social interaction between individuals and can cause an individual to experience loneliness, anxiety, and depression [32]. A crowded environment, noise, and lack of greenery in urban areas can also increase mental health disorders.

The studies [15,17-19,21-26] show that people with higher education experience mental health disorders more than people with lower education. Those with a higher education tended to experience more stress during the COVID-19 pandemic because they had higher self-awareness of their health condition. Not only that, a higher level of education can also be linked to the greater curiosity about health information during the COVID-19 pandemic, which has negatively affected some people's mental health.

The results of this study [13,14,16,17,20,21] reveal that people who earn higher income are more likely to experience mental health problems during the COVID-19 pandemic compared to people with lower income. A good economic situation makes it easy to have access to information and to healthcare services. This in turn may lead to mental health problems caused by being exposed to frightening information such as death incidents that occur every day.

Non-sociodemographic factors that influence mental health

The results of the studies show that people who are afraid of being infected tend to experience mental health problems more than people who are not afraid. Feeling afraid and worried was to be expected during the early days of the COVID-19 pandemic. The research indicate that people's mental health has begun to deteriorate since the beginning of the COVID-19 pandemic which was caused by the fear of being infected with COVID-19 [27] and worrying about infecting their close ones [5,12]. A person who is more worried and afraid of contracting COVID-19 is more likely to have poor mental health [27], as worrying about COVID-19 and feeling depressed can deteriorate a person's mental health [28]. During the pandemic, people were afraid that they or their family members would fall ill but they did not know yet fully the dangerous effects of the pandemic. In addition, discrimination and stigma related to infectious diseases can make people afraid of being infected, which in turn affects their mental health. Makhshvili et al. stated that people's lack of trust in the media regarding COVID-19 information could also cause their mental health to deteriorate [16].

The studies [18,19,21,25,26] show that someone who was suffering from a chronic illness during the COVID-19 pandemic was more likely to experience mental health problems in comparison to someone who was healthy. Chronic diseases such as cardiovascular, neurological, and respiratory diseases were usually found in people who experienced mental health disorders during the pandemic. People who suffer from chronic illnesses and think of themselves as having poor health are more susceptible to other diseases, especially COVID-19, resulting in stress on their mental health.

The research results also indicate that people who work from home experience mental health problems more than people who do not work from home. People who stay at home continuously and feel lonely can experience increased anxiety and depression [15]. People

experience stress, depression, anxiety, and post-traumatic stress disorder (PTSD) because of being lonely [28]. There has been an increase in poor mental health during the COVID-19 pandemic among respondents living alone [14]. There is an increase in PTSD symptoms due to several factors, including the lack of social support [13].

Family conflict is a determining factor in mental health [33]. During the COVID-19 pandemic, many family conflicts occurred due to termination of employment relationships [34,35]. Families whose members lost their jobs, suffered a loss of income and found it difficult to meet their family's needs. This is what makes families experience financial distress and become vulnerable to conflict within the family [36]. Positive coping, having good social strengths, and family support can offer protection from stress or mental disorders during COVID-19 [37].

Conclusions

The determinants of mental health disorders found in the community during the COVID-19 pandemic include age, gender, place of residence, income/economics, chronic diseases, family conflict, and family support. It is hoped that the results of this literature review can help nurses determine and control factors that affect mental health in the community during the COVID-19 pandemic in order to achieve good outcomes and to prevent more serious conditions due to COVID-19. In addition, the results of this study can also be a source of literature and information for academic and clinical nursing, as mental health is important for facing not only COVID-19 but also future health threats. Of course, the results of this research depend on the database analyzed, so expanding the database will provide a more comprehensive view. The benefit of this research after the COVID-19 pandemic is the importance of preparing community resilience to face disasters in the future. Community resilience under challenging

situations is greatly influenced by factors such as skills in dealing with conflict, the importance of support, and togetherness. The implications of this research can be aimed at regional governments by providing health promotion budgets that focus on community resilience in facing disasters. Meanwhile, nurses and other healthcare professionals (such as midwives, physiotherapists, pharmacists and doctors) are making health promotion plans to improve the community's mental health. Further research can be developed using experimental methods on how conflict management or social support affects mental health.

Disclosures and acknowledgements

The authors declare no conflicts of interest with respect to the research, authorship, and/or publication of this article. This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. Artificial intelligence (AI) was not used in the creation of the manuscript.

References:

1. Mikolai J, Dorey P, Keenan K, Kulu H. Spatial patterns of COVID-19 and non-COVID-19 mortality across waves of infection in England, Wales, and Scotland. *Soc Sci Med.* 2023; 338: 116330. <https://doi.org/10.1016/j.socscimed.2023.116330>
2. Borczuk AC. Pathogenesis of pulmonary long COVID-19. *Mod Pathol.* 2024; 37(2): 100378. <https://doi.org/10.1016/j.modpat.2023.100378>
3. Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. *Acta Biomed.* 2020; 91(1): 157-160.

4. Romay-Barja M, Pascual-Carrasco M, De Tena-Dávila MJ, Falcón M, Rodriguez-Blazquez C, Forjaz MJ, et al. How patients with COVID-19 managed the disease at home during the first wave in Spain: a cross-sectional study. *BMJ Open*. 2021; 11: e048702. <https://doi.org/10.1136/bmjopen-2021-048702>
5. Schrepft S, Pullen N, Baysson H, Wisniak A, Zaballa M-E, Pennacchio F, et al. Prevalence and predictors of psychological distress before, during, and after a COVID-19 pandemic wave in Switzerland, 2021. *J Psychiatr Res*. 2023; 158: 192-201. <https://doi.org/10.1016/j.jpsychires.2022.12.042>
6. Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: A systematic review and meta-analysis. *Global Health*. 2020; 16(1): 1-11. <https://doi.org/10.1186/s12992-020-00589-w>
7. Wang C, Pan R, Wan X, Tan Y, Xu L, Cyrus SH, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *QJM*. 2020; 113(5): 311-2. <https://doi.org/10.3390/ijerph17051729>
8. Alkhamees AA, Alrashed SA, Alzunaydi AA, Almohimeed AS, Aljohani MS. The psychological impact of COVID-19 pandemic on the general population of Saudi Arabia. *Compr Psychiatry*. 2020; 102: 152192. <https://doi.org/10.1016/j.comppsy.2020.152192>
9. Ridlo IA. [The COVID-19 pandemic and mental health policy challenges in Indonesia]. *Insa J Psikol dan Kesehat Ment*. 2020; 5(2): 162 (in Indonesian). <https://doi.org/10.20473/jpkm.V5I22020.162-171>

10. Ilpaj SM, Nurwati N. Analysis of the influence of the death rate due to COVID-19 on the mental health of people in Indonesia. *Focus J Pekerj Sos.* 2020; 3(1): 16. <https://doi.org/10.24198/focus.v3i1.28123>
11. Afonso P. [The impact of the COVID-19 pandemic on mental health]. *Acta Med Port.* 2020; 33(5): 356 (in Portugese). <https://doi.org/10.20344/amp.13877>
12. Agodi A, Maugeri A, Favara G, Magnano San Lio R, Puglisi M, Sinatra D, et al. Gender differences in comorbidities of patients with COVID-19: An Italian local register-based analysis. *Heliyon.* 2023; 9(7): e18109. <https://doi.org/10.1016/j.heliyon.2023.e18109>
13. Bonsaksen T, Heir T, Schou-Bredal I, Ekeberg Ø, Skogstad L, Grimholt TK. Post-traumatic stress disorder and associated factors during the early stage of the COVID-19 pandemic in Norway. *Int J Environ Res Public Health.* 2020; 17(24): 9210. <https://doi.org/10.3390/ijerph17249210>
14. Guo Y, Cheng C, Zeng Y, Li Y, Zhu M, Yang W, et al. Mental health disorders and associated risk factors in quarantined adults during the COVID-19 outbreak in China: cross-sectional study. *J Med Internet Res.* 2020; 22(8): e20328. <https://doi.org/10.2196/20328>
15. Kantor BN, Kantor J. Mental health outcomes and associations during the COVID-19 pandemic: a cross-sectional population-based study in the United States. *Front Psychiatry.* 2020; 11: 569083. <https://doi.org/10.3389/fpsy.2020.569083>
16. Makhashvili N, Javakhishvili JD, Sturua L, Pilauri K, Fuhr DC, Roberts B. The influence of concern about COVID-19 on mental health in the Republic of Georgia: a cross-sectional study. *Global Health.* 2020; 16(1): 111. <https://doi.org/10.1186/s12992-020-00641-9>

17. Mani A, Estedlal AR, Kamali M, Ghaemi SZ, Zarei L, Shokrpour N, et al. Mental health status during COVID-19 pandemic in Fars Province, Iran: timely measures. *BMC Public Health*. 2020; 20(1): 1-11. <https://doi.org/10.1186/s12889-020-09928-3>
18. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A nationwide survey of psychological distress among Italian people during the covid-19 pandemic: Immediate psychological responses and associated factors. *International Journal of Environmental Research and Public Health*. 2020; 17(3165): 1-14. <https://doi.org/10.1186/s12889-020-09928-3>
19. Newby JM, O'Moore K, Tang S, Christensen H, Faasse K. Acute mental health responses during the COVID-19 pandemic in Australia. *PLoS One*. 2020; 15(7): e0236562. <https://doi.org/10.1371/journal.pone.0236562>
20. Smith L, Jacob L, Yakkundi A, McDermott D, Armstrong NC, Barnett Y, et al. Correlates of symptoms of anxiety and depression and mental wellbeing associated with COVID-19: a cross-sectional study of UK-based respondents. *Psychiatry Res*. 2020; 291: 113138.
21. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, tobacco and alcohol use in Australian adults. *Int J Environ Res Public Health*. 2020; 17(11): 4065. <https://doi.org/10.3390/ijerph17114065>
22. Venugopal VC, Mohan A, Chennabasappa LK. Status of mental health and its associated factors among the general populace of India during COVID-19 pandemic. *Asia-Pacific Psychiatry*. 2022; 14(2): e12412. <https://doi.org/10.1111/appy.12412>
23. Verma S, Mishra A. Depression, anxiety, and stress and socio-demographic correlates among general Indian public during COVID-19. *Int J Soc Psychiatry*. 2020; 66(8): 756-62. <https://doi.org/10.1177/0020764020934508>

24. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health*. 2020; 17(5): 1729. <https://doi.org/10.3390/ijerph17051729>
25. Wong SYS, Zhang D, Sit RWS, Yip BHK, Chung RY, Wong CKM, et al. Impact of COVID-19 on loneliness, mental health, and health service utilisation: a prospective cohort study of older adults with multimorbidity in primary care. *Br J Gen Pract*. 2020; 70(700): e817-24. <https://doi.org/10.3399/bjgp20X713021>
26. El-Zoghby SM, Soltan EM, Salama HM. Impact of the COVID-19 pandemic on mental health and social support among adult Egyptians. *J Community Health*. 2020; 45(4): 689-695. <https://doi.org/10.1007/s10900-020-00853-5>
27. Choi EPH, Hui BPH, Wan EYF. Depression and anxiety in Hong Kong during COVID-19. *Int J Environ Res Public Health*. 2020; 17(10): 3740. <https://doi.org/10.3390/ijerph17103740>
28. Liu CH, Zhang E, Wong GTF, Hyun S, Hahm HC. Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: clinical implications for U.S. young adult mental health. *Psychiatry Res*. 2020; 290:113172. <https://doi.org/10.1016/j.psychres.2020.113172>
29. Winkler P, Formanek T, Mlada K, Kagstrom A, Mohrova Z, Mohr P, et al. Increase in prevalence of current mental disorders in the context of COVID-19: analysis of repeated nationwide cross-sectional surveys. *Epidemiol Psychiatr Sci*. 2020; 29: e173. <https://doi.org/10.1017/S2045796020000888>
30. Qiu J, Shen B, Zhao M, Wang Z, Xie B, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy

- recommendations. *Gen Psychiatry*. 2020; 33(2): e100213.
<https://doi.org/10.1136/gpsych-2020-100213>
31. Thibaut F, ELNahas G. Women's Mental Health and Lessons Learnt from the COVID-19 Pandemic. *Psychiatr Clin North Am*. 2023; 46(3): 415-426.
<https://doi.org/10.1016/j.psc.2023.04.001>
32. Addai G, Amponsah O, Dinye RD. Social interactions in periodic urban markets and their contributions to sustainable livelihoods: Evidence from Ghana. *Reg Sustain*. 2023; 4(4): 369-377. <https://doi.org/10.1016/j.regsus.2023.10.002>
33. Bilodeau J, Quesnel-Vallée A, Beaugard N, Brault MC. Gender, work-family conflict and depressive symptoms during the COVID-19 pandemic among Quebec graduate students. *Prev Med Reports*. 2021; 24: 101568.
<https://doi.org/10.1016/j.pmedr.2021.101568>
34. Holingue C, Kalb LG, Riehm KE, Bennett D, Kapteyn A, Veldhuis CB, et al. Mental distress in the United States at the beginning of the COVID-19 pandemic. *Am J Public Health*. 2020; 110(11): 1628-1634. <https://doi.org/10.2105/AJPH.2020.305857>
35. Ashcroft R, Donnelly C, Dancy M, Gill S, Lam S, Kourgiantakis T, et al. Primary care teams' experiences of delivering mental health care during the COVID-19 pandemic: a qualitative study. *BMC Fam Pract*. 2021; 1-12. <https://doi.org/10.1186/s12875-021-01496-8>
36. Daks JS, Peltz JS, Rogge RD. Psychological flexibility and inflexibility as sources of resiliency and risk during a pandemic: Modeling the cascade of COVID-19 stress on family systems with a contextual behavioral science lens. *J Context Behav Sci*. 2020; 18: 16-27. <https://doi.org/10.1016/j.jcbs.2020.08.003>
37. Cai Z, Cui Q, Liu Z, Li J, Gong X, Liu J, et al. Nurses endured high risks of psychological problems under the epidemic of COVID-19 in a longitudinal study in

Wuhan China. J Psychiatr Res. 2020; 131:132-137.

<https://doi.org/10.1016/j.jpsychires.2020.09.007>

ONLINE FIRST