

CASE STUDY

STUDIUM PRZYPADKU

**ACTIVITY LEVEL AND QUALITY OF LIFE AMONG PATIENTS UNDERGOING  
KNEE REPLACEMENT SURGERY**

**POZIOM AKTYWNOŚCI I JAKOŚĆ ŻYCIA PACJENTÓW PODDANYCH  
ENDOPROTEZOPLASTYCE STAWU KOLANOWEGO**

**Huda Alfatafta<sup>1(A,B)</sup>, Mahmoud Alfatafta<sup>2(A,B)</sup>, David Onchonga<sup>1(B,C)</sup>, Haitham Khatatbeh<sup>1(C,F)</sup>,  
Faten Amer<sup>1(E,F)</sup>, Peter Than<sup>1(C,D,F)</sup>, Balint Molics<sup>1(C,D,E,F)</sup>, Boncz Imre<sup>1(B,C,E,G)</sup>, Pongrac Acs<sup>1(F,G)</sup>**

<sup>1</sup>Faculty of Health Sciences, University of Pecs, Pecs, Hungary

<sup>2</sup>Orthotics and Prosthetics Department, Faculty of Rehabilitation Sciences, University of Jordan, Amman, Jordan

Alfatafta H, Alfatafta M, Onchonga D, Khatatbeh H, Amer F, Than P, et al. Activity level and quality of life among patients undergoing knee replacement surgery. Health Prob Civil. <https://doi.org/10.5114/hpc.2024.137425>

Tables: 3

Figures: 0

References: 30

Submitted: 2024 Jan 22

Accepted: 2024 March 25

**Address for correspondence / Adres korespondencyjny:** Huda Alfatafta, Faculty of Health Sciences, University of Pécs, Vörösmarty str. 4, H-7621 Pécs, Hungary, e-mail: [huda\\_alfatafta@hotmail.com](mailto:huda_alfatafta@hotmail.com), phone: +36-72/501-500/61682

ORCID: Huda Alfatafta <https://orcid.org/0000-0002-2063-329X>, Mahmoud Alfatafta <https://orcid.org/0000-0003-2511-1658>, David Onchonga <https://orcid.org/0000-0003-4461-797X>, Haitham Khatatbeh <https://orcid.org/0000-0003-2019-6621>, Faten Amer <https://orcid.org/0000-0002-5214-3797>, Peter Than <https://orcid.org/0000-0002-9388-7557>, Boncz Imre <https://orcid.org/0000-0003-3699-6236>, Pongrac Acs <https://orcid.org/0000-0002-4999-7345>.

Copyright: © John Paul II University in Białá Podlaska, Huda Alfatafta, Mahmoud Alfatafta, David Onchonga, Haitham Khatatbeh, Faten Amer, Peter Than, Balint Molics, Boncz Imre, Pongrac Acs. This is an Open Access journal, all articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited and states its license.

### Summary

Knee osteoarthritis (OA) is a widespread disease in Hungary. For severe cases, knee replacement surgery is the last-stage intervention to reduce pain and improve the quality of life. Nevertheless, there is no information about the activity level of Hungarians with severe knee OA. This study evaluated the improvement in the quality of life pre-post knee replacement surgery among Hungarian patients. This study included eight participants ( $70.8 \pm 4.5$  years and  $30.0 \pm 3.0$  kg/m<sup>2</sup>) with severe knee OA undergoing total unilateral knee replacement surgery. The activity level was evaluated with ActivPAL and Short Form-36 one month before surgery and one year after surgery.

One year after surgery, the sedentary time and standing time had significantly improved among the patients by 12% ( $p=0.033$ ) and 65.7% ( $p=0.030$ ), respectively. The number of steps improved by 17.1% ( $p=0.11$ ) based on ActivPAL. Based on the Short Form-36, the overall score, pain, and physical function had improved after one year by 50.0% ( $p=0.050$ ), 23.9% ( $p=0.32$ ), and 5.1% ( $p=0.58$ ), respectively. This is the first study in Hungary that used both objective and subjective tools. The outcomes found that the activity level and pain improved after surgery among Hungarian patients with severe knee OA. However, the sedentary time after the surgery should be reduced to reach the maximum benefits of surgery.

**Keywords:** knee osteoarthritis, knee replacement, knee, physical activity, quality of life

### Streszczenie

Choroba zwyrodnieniowa stawu kolanowego jest szeroko rozpowszechniona na Węgrzech. W ciężkich przypadkach endoprotezoplastyka stawu kolanowego jest ostatnim etapem interwencji mającej na celu złagodzenie bólu i poprawę jakości życia. Brak jednak informacji na temat poziomu aktywności Węgrów z ciężką chorobą zwyrodnieniową stawów kolanowych. W niniejszych badaniach oceniono poprawę jakości życia po endoprotezoplastyce stawu kolanowego wśród węgierskich pacjentów. W badaniach wzięło udział ośmiu uczestników ( $70,8\pm 4,5$  lat i  $30,0\pm 3,0$  kg/m<sup>2</sup>) z ciężką chorobą zwyrodnieniową stawu kolanowego poddanych operacji całkowitej jednostronnej endoprotezoplastyki stawu kolanowego. Poziom aktywności został oceniony za pomocą ActivPAL i Short Form-36 na miesiąc przed operacją i rok po operacji. Po roku od operacji znacznej poprawie uległ czas siedzenia i czas stania pacjentów, odpowiednio o 12% ( $p=0,033$ ) i 65,7% ( $p=0,030$ ). Liczba kroków wzrosła o 17,1% ( $p=0,11$ ) w oparciu o ActivPAL. Na podstawie formularza Short Form-36 ogólny wynik, poziom bólu i sprawności

fizycznej po roku uległy poprawie odpowiednio o 50,0% ( $p=0,050$ ), 23,9% ( $p=0,32$ ) i 5,1% ( $p=0,58$ ). Są to pierwsze na Węgrzech badania, w których wykorzystano zarówno obiektywne, jak i subiektywne narzędzia. Wyniki wykazały, że poziom aktywności i bólu u węgierskich pacjentów z ciężką chorobą zwyrodnieniową stawu kolanowego poprawił się po operacji. Aby jednak osiągnąć maksymalne korzyści z endoprotezoplastyki stawu kolanowego, należałoby skrócić czas siedzenia po operacji.

**Słowa kluczowe:** choroba zwyrodnieniowa stawu kolanowego, endoproteza stawu kolanowego, kolano, aktywność fizyczna, jakość życia

## Introduction

Knee osteoarthritis (OA) is one of the most common diseases among the elderly population globally, as it is associated with age and obesity [1-3]. In Hungary, the prevalence of knee OA was 16.54% in 2011. In a group aged 20-67 years, a total of 2.9% had severe OA [3]. The core symptoms associated with knee OA are pain, low physical activity (PA) levels, and disabilities [1-3].

Conservative treatments, such as pharmacological treatment, orthotics, physiotherapy sessions, manual exercises, acupuncture, and other possible interventions, could be effective for mild-moderate knee OA [4-8]. Total knee replacement (TKR) surgery is the gold-standard treatment to deal with severe knee OA when the pain is no longer relieved by conservative treatments [9,10]. Thus, the expected goals of TKR surgery are reducing pain and improving the quality of life (QOL) [9-11].

## Case description

In Hungary, pain and PA after TKR were evaluated only once using four forms of questionnaires: the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), a short form questionnaire (SF-36), the Knee Society Score (KSS), and a self-made questionnaire [12]. A conference paper reported the QOL before the surgery and six weeks after the surgery and found that the QOL had significantly improved after six weeks from the surgery [12]. However, unfortunately, the full article of this conference paper is not available, some information about the patients' characteristics, numbers, and methodology are missing, and six weeks are not enough to evaluate the QOL improvement. Furthermore, the questionnaires used to assess the PA of patients with knee OA are associated with recall bias and have low-moderate reliability and validity [13,14]. Hence, it was necessary to evaluate the PA of Hungarian patients with severe knee OA using high validity and reliability objective tools, as no databases are available in this field in Hungary. To our knowledge, no study has explored the PA level of Hungarian patients before and after surgery using objective tools or has explored the PA level of Hungarian patients before and after surgery in terms of the number of steps, sedentary time, standing time, and stepping time using a high validity and reliability accelerometer such as ActivPAL. This information is important to understand the PA level and physical behavior of Hungarian patients with severe knee OA before and after surgery using objective tools.

Therefore, the aims of this case-series study were (a) to understand the activity level of Hungarian patients with severe knee OA and (b) to evaluate the activity level of Hungarian patients one year after TKR surgery. The measured variables were evaluated with both objective and subjective monitoring tools.

### *Study design*

This study is a case series study that evaluated the QOL and activity level one month before surgery and one year after surgery among eight Hungarian patients (four females, four males) with an average age of  $70.8 \pm 4.5$  years old and  $30.0 \pm 3.0$  kg/m<sup>2</sup> (Table 1).

**Table 1.** Demographic data of the participants.

<b>Domains</b>	<b>Total (n=8)</b>
<b>Age (y)</b>	70.8±4.5
<b>Gender</b>	4 males, 4 females
<b>BMI (kg/m<sup>2</sup>)</b>	30.7±4.3
<b>Smoking</b>	0
<b>Heart problems</b>	3 (2 males, 1 female)
<b>Diabetes</b>	2 (1 male, 1 female)
<b>Hypertension</b>	6 (2 males, 4 females)

### *Subjects*

The inclusion criteria were the presence of a clinical and radiological diagnosis of OA that required knee replacement surgery. The confirmation of a radiological diagnosis was performed by the same orthopedic surgeon at the orthopedic clinic, at the University of Pecs, to ensure that participants needed knee replacement surgery. Participants were excluded if they had had hip and ankle injuries in the previous five years or if they had co-morbidities or medical conditions that affected PA, such as congestive heart failure or cognitive impairment. Ethical approval was obtained from the University of Pecs (no. 8343/2020 of July 2020) and consent was initially

collected from the participants. All methods were carried out according to relevant guidelines and regulations.

#### *PA measurement tools*

The activity level was evaluated with ActivPAL and a short form (SF-36) questionnaire. ActivPAL (PAL Technologies, Glasgow, UK) is a uniaxial accelerometer (20 g) that calculates the time spent in sedentary, standing, upright, and stepping states and the number of steps per day for up to 14 days [15-19]. ActivPAL is recommended to be used with the elderly more than other monitors, such as ActiGraph, because it has a higher validity and reliability to detect body movement at different speeds than ActiGraph [19,20]. Before use, the monitor was charged and activated with ActivPAL3™ (version 8.11.9.100). Moreover, the Hungarian version of the short form (SF-36) was used to assess QOL. This form is composed of 36 items about eight domains: physical functioning, physical role, bodily pain, general health, vitality, social functioning, emotional role, and mental health. Each question is scaled between zero (poor health status) and 100 (no problem at all) [21-23].

#### *Intervention and TKR surgery*

All of the participants underwent the operation with the medial parapatellar approach that was performed by the orthopedic surgeon. Various prosthetic knee types were used, such as S and N Genesis II, Johnson and Johnson PFC sigma, and Zimmer Nexgen. No compliances were reported after the surgery. The patients stayed five to six days at the hospital after the surgery.

Physiotherapy sessions were standardized according to hospital protocols to minimize confounding factors for both inpatient and outpatient periods (for three months only).

#### *Data collection procedure*

One month before the surgery, all participants were briefed about the study first and signed the consent form. Then, each participant was asked to complete a Hungarian language SF-36 form one month before the surgery. At that session, the participants were instructed to wear the ActivPAL for seven days most of the time (removed during shower time or water activities). The monitors were set to start measuring the PA from the second day of that visit for seven days. The instructions were given verbally and written to ensure that they understood the instructions. The ActivPAL was attached at the mid-thigh with self-adhesive tape under the clothes. The participants were asked to return the monitor on the day of the surgery.

One year after the surgery, the participants were asked to visit the clinic for follow-up, fill out the SF-36, and use the ActivPAL for one week. The same previously mentioned instructions were given again to each patient. Then, the participants were asked to return the monitor after one week of usage by post.

#### *Data analysis*

For data from the SF-36 questionnaire, the average ( $\pm$ SD) of physical functioning, physical role, pain, general health, and the overall QOL score were calculated for each participant at all assessment time points, as these aspects are the most relevant aspects for the study's goal. The data



from the ActivPAL were extracted from the monitor using the software. Then, the downloaded files were imported into Excel. The average ( $\pm$ SD) sedentary time, standing time, stepping time, and the number of steps were calculated for each participant at all assessment time points during the testing period (6-7 days). A valid day is defined as 10 hours of continuous activity with less than three hours of interruptions. All the data were analyzed by SPSS (SPSS Inc., Chicago, IL, USA) using a paired sample test. Significant results were considered if the significance (2-tailed) value was less than 0.05 [24]. Only eight patients were included in the paired sample t-tests. The data were normally distributed based on the Kolmogorov-Smirnov test [25].

### Results

Ten participants initially took part in this study. After one year, one participant was not available for the final study, and the ActivPAL data of one participant were not included in the pre-surgery data because his data were not valid. Thus, eight participants completed the SF-36 form and had valid ActivPAL data for this study (Tables 2 and 3).

**Table 2.** The average ( $\pm$  SD) of the activity level one month before and one year after the TKR surgery based on ActivPal

Domains	Pre-surgery	One year post-surgery	Significant (p-value)	95% Confidence interval for differences (IC)	
				Low	High
<b>ActivPAL data (pre-surgery n=8, post-surgery n=8)</b>					
<b>Time of sedentary (hour)</b>	18.3 $\pm$ 1.9	16.1 $\pm$ 3.1	0.033	0.24	4.1
<b>Time of standing (hour)</b>	3.5 $\pm$ 1.4	5.8 $\pm$ 2.7	0.030	-4.22	-0.11
<b>Time of stepping (hour)</b>	1.5 $\pm$ 0.6	2.2 $\pm$ 0.9	0.046	-1.2	-0.01

<b>Number of steps (number)</b>	6270±2754	7344±3331	0.11	-2507.04	358.0
---------------------------------	-----------	-----------	------	----------	-------

Notes: The number of paired samples is eight.

**Table 3.** The average ( $\pm$  SD) activity level one month before and one year after the TKR surgery is based on the SF-36

Domains	Pre-surgery	One year post-surgery	Significant (p-value)	95% Confidence interval for differences (IC)	
				Low	High
<b>SF-36 score (pre-surgery n=8, post-surgery n=8)</b>					
<b>Overall score</b>	50.0±26.7	75.0±23.1	0.050	-49.9	-0.19
<b>Physical functioning</b>	58.1±20.8	61.2±21.5	0.582	-15.5	9.6
<b>Role limitations due to physical health</b>	46.8±41.0	43.7±7.4	0.89	-50.9	57.2
<b>Bodily pain</b>	41.8±23.5	51.8±25.1	0.32	-32.3	12.7
<b>General health</b>	52.5±11.6	56.2±20.6	0.704	-26.1	18.6

Notes: The number of paired samples is eight.

#### *ActivPAL results*

The PA level was enhanced after one year of knee replacement surgery based on an objective monitor (Table 2). Before the surgery, patients spent  $3.5 \pm 1.4$  hours per day and  $5.8 \pm 2.2$  hours per day standing after one year of the surgery (65.7%,  $p=0.030$ ). The stepping time also improved by 46.6% ( $p=0.046$ ) one year after surgery. Furthermore, the number of steps increased from  $6270 \pm 2754$  to  $7344 \pm 3331$  but insignificantly (17.1%,  $p=0.11$ ). One year after the surgery, patients spent 2.2 hours per day (12%,  $p=0.033$ ) less on sedentary activity than before the surgery.

### *SF-36 results*

The total QOL score of the patients was  $50.0 \pm 26.7$  before the surgery and improved to  $75.0 \pm 23.1$  one year after the surgery ( $p=0.05$ ) (Table 3). The bodily pain based on this questionnaire increased from  $41.8 \pm 23.5$  to  $51.8 \pm 25.1$  (23.9%,  $p=0.32$ ). Furthermore, role limitations due to physical health increased by 6.6% ( $p=0.89$ ). In addition, physical functioning and general health one year after knee replacement surgery improved by 5.1% ( $p=0.58$ ) and 7.0% ( $p=0.70$ ), respectively.

### **Case analysis**

This study measured the activity level before and one year after TKR surgery among Hungarian patients. To our knowledge, this study is the first study in Hungary that used a highly valid objective accelerometer to understand the PA level of Hungarian patients with severe knee OA and their PA improvement one year after TKR. This study found that Hungarian patients with severe knee OA spend most of their daily time in a sedentary manner, with an average of  $18.3 \pm 1.9$  hours per day, and that they only spend almost  $1.5 \pm 0.6$  hours per day walking, with an average of  $6270 \pm 2754$  steps per day. After one year of TKR, Hungarian patients have better PA levels and less pain (less pain or a better QOL in general) based on both objective and subjective tools.

Hence, the activity level increased between 65.7%-12% based on the ActivPAL and between 50%-5.1% after the surgery. However, the sedentary time decreased from  $18.3 \pm 1.9$  to  $16.1 \pm 3.1$  hours per day, and these changes are still not enough to allow Hungarian patients to meet the general PA guidelines [26].

Few studies have evaluated the PA using the same accelerometer with a one-year follow-up [27,28]. Bin Sheeha et al., 2020 [27] evaluated the PA of 33 patients before surgery and one year after. This study found that the stepping time and the number of steps improved significantly by 38.77% and 45.6%, respectively [27]. Moreover, Lützner et al. [28] evaluated the PA level of 97 patients before knee surgery and one year after using ActivPAL. This study found that the number of steps improved by 20.3% (from  $5278 \pm 2999$  to  $6473 \pm 3654$  steps/day), and no changes were seen in the sedentary and stepping time one year after the surgery. However, this result could be due to applying the ActivPAL over the tibia, which is a less reliable position, and the monitor was used for just four days [28].

Furthermore, this study found that the PA of Hungarian patients was improved based on the SF-36 questionnaire. The total score of the SF-36 questionnaire significantly improved by 50% ( $p=0.050$ ) one year after surgery. Similarly, among Greek elderly women, the total SF-36 score significantly improved from  $29.33 \pm 11.3$  before surgery to  $62.35 \pm 2.7$  six months after surgery [29]. It seems that Hungarian patients have higher PA levels than Greek patients before and after surgery. This finding could be due to the very severe pain the Greek patients had before surgery, as 85% of them had continuous severe pain before the surgery [29]. Other studies used different types of questionnaires. For instance, Granat et al. [27] found that PA based on the Oxford knee score (OKS) significantly increased by 142% ( $p=0.00$ ) one year after surgery. Additionally, the OKS score improved six months after surgery from 12 points to 42 points [30].

In summary, the activity level and pain among Hungarian patients with severe knee OA improved after one year of TKR. However, the changes were not enough to meet the general PA guidelines. Long follow-ups and staying active after surgery are still necessary to obtain better outcomes. These findings could be important for therapists who care for Hungarian patients with

severe knee OA to understand their physical captivity limitations before and after surgery. Additionally, they should be helped by focusing on how to enhance the outcomes of the surgery to reach the maximum activity level improvement.

## **Conclusions**

Based on both subjective and objective assessment tools, Hungarian patients had better PA levels and QOL after TKR. However, the sedentary time among Hungarian patients after surgery is still high and might reflect a reduced efficiency of the surgery. In addition, this study emphasized including objective assessment tools, as they are more sensitive and reliable than subjective assessment tools. Thus, it is important to include objective assessment tools in the evaluation to reduce recall bias and represent more information about the PA level. Hence, long-term follow-up and rehabilitation sessions could be required to reduce the sedentary time and increase the efficiency of the surgery. More research should be done in Hungary to understand the effect of TKR surgery with more participants from different parts of the country.

## **Disclosures and acknowledgments**

The authors declare no conflicts of interest concerning the research, authorship, and/or publication of this article. The research was funded by project no. TKP-2021-EGA-10. Artificial intelligence (AI) was not used in the creation of the manuscript.

**References:**

1. Cross M, Lapsley H, Barcenilla A, Parker D, Coolican M, March L. Patient expectations of hip and knee joint replacement surgery and postoperative health status. *Patient*. 2009; 2(1): 51-60. <https://doi.org/10.2165/01312067-200902010-00006>
2. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Annals of the Rheumatic Diseases*. 2014; 73(7): 1323-1330. <https://doi.org/10.1136/annrheumdis-2013-204763>
3. Horváth G, Koroknai G, Ács B, Than P, Bellyei Á, Illés T. Prevalence of radiographic primary hip and knee osteoarthritis in a representative Central European population. *International Orthopaedics*. 2011; 35(7): 971-975. <https://doi.org/10.1007/s00264-010-1069-6>
4. Alfatafta HH, Hutchins S, Liu A, Jones R. Effect of a knee-ankle-foot orthosis on knee kinematics and kinetics in an individual with varus knee alignment. *JPO: Journal of Prosthetics and Orthotics*. 2016; 28(4): 186-190. <https://doi.org/10.1097/JPO.0000000000000111>
5. Alfatafta H, Onchonga D, Alfatafta M, Zhang I, Boncz I, Lohner S, et al. Effect of using knee valgus brace on pain and activity level over different time intervals among patients with medial knee OA: systematic review. *BMC Musculoskeletal Disorders*. 2021; 22(1): 1-9. <https://doi.org/10.1186/s12891-021-04513-0>

6. Crawford DC, Miller LE, Block JE. Conservative management of symptomatic knee osteoarthritis: a flawed strategy?. *Orthopedic Reviews*. 2013; 5(1). <https://doi.org/10.4081/or.2013.e2>
7. Sánchez Romero EA, Fernández-Carnero J, Calvo-Lobo C, Ochoa Sáez V, Burgos Caballero V, Pecos-Martín D. Is a combination of exercise and dry needling effective for knee OA?. *Pain Medicine*. 2020; 21(2): 349-363. <https://doi.org/10.1093/pm/pnz036>
8. Sánchez-Romero EA, González-Zamorano Y, Arribas-Romano A, Martínez-Pozas O, Fernández Espinar E, Pedersini P, et al. Efficacy of manual therapy on facilitatory nociception and endogenous pain modulation in older adults with knee osteoarthritis: A case series. *Applied Sciences*. 2021; 11(4): 1895. <https://doi.org/10.3390/app11041895>
9. Malviya A, Lingard EA, Weir DJ, Deehan DJ. Predicting range of movement after knee replacement: the importance of posterior condylar offset and tibial slope. *Knee Surg Sports Traumatol Arthrosc*. 2009; 17(5): 491-498. <https://doi.org/10.1007/s00167-008-0712-x>
10. Ferket BS, Feldman Z, Zhou J, Oei EH, Bierma-Zeinstra SMA, Mazumdar M. Impact of total knee replacement practice: cost-effectiveness analysis of data from the Osteoarthritis Initiative. *BMJ*. 2017; 356: j1131. <https://doi.org/10.1136/bmj.j1131>
11. Tambascia RA, Antunes Vasconcelos R, Mello W, Portes Teixeira P, Bevilaqua Grossi D. Pre-operative functional parameters of patients undergoing total knee arthroplasty. *Physiother Res Int*. 2016; 21(2): 77-83. <https://doi.org/10.1002/pri.1622>
12. Busa M, Gunther T, Mazzag K, Vizsy M, Boncz I, Molics B. PMS6 quality of life of patients underwent implantation of knee endoprosthesis. *Value in Health*. 2019; 22: S694-S695. <https://doi.org/10.1016/j.jval.2019.09.1550>

13. Collins NJ, Roos EM. Patient-reported outcomes for total hip and knee arthroplasty: commonly used instruments and attributes of a “good” measure. *Clinics in Geriatric Medicine*. 2012; 28(3): 367-394. <https://doi.org/10.1016/j.cger.2012.05.007>
14. Collins NJ, Prinsen CAC, Christensen R, Bartels EM, Terwee CB, Roos EM. Knee Injury and Osteoarthritis Outcome Score (KOOS): systematic review and meta-analysis of measurement properties. *Osteoarthritis Cartilage*. 2016; 24(8): 1317-29. <https://doi.org/10.1016/j.joca.2016.03.010>
15. Edwardson CL, Winkler EAH, Bodicoat DH, Yates T, Davies MJ, Dunstan DW, et al. Considerations when using the activPAL monitor in field-based research with adult populations. *Journal of Sport and Health Science*. 2017; 6(2): 162-178. <https://doi.org/10.1016/j.jshs.2016.02.002>
16. Lyden K, Keadle SK, Staudenmayer J, Freedson PS. The activPAL™ accurately classifies activity intensity categories in healthy adults. *Medicine and Science in Sports and Exercise*. 2017; 49(5): 1022. <https://doi.org/10.1249/MSS.0000000000001177>
17. Dahlgren G, Carlsson D, Moorhead A, Häger-Ross C, McDonough SM. Test–retest reliability of step counts with the ActivPAL™ device in common daily activities. *Gait & Posture*. 2010; 32(3): 386-390. <https://doi.org/10.1016/j.gaitpost.2010.06.022>
18. Taraldsen K, Askim T, Sletvold O, Einarsen EK, Gruner Bjåstad K, Indredavik B, et al. Evaluation of a body-worn sensor system to measure physical activity in older people with impaired function. *Physical Therapy*. 2011; 91(2): 277-285. <https://doi.org/10.2522/ptj.20100159>



19. Ryan CG, Grant PM, Tigbe WW, Granat MH. The validity and reliability of a novel activity monitor as a measure of walking. *British Journal of Sports Medicine*. 2006; 40(9): 779-784. <https://doi.org/10.1136/bjism.2006.027276>
20. Kim Y, Barry VW, Kang M. Validation of the ActiGraph GT3X and activPAL accelerometers for the assessment of sedentary behavior. *Measurement in Physical Education and Exercise Science*. 2015; 19(3): 125-137. <https://doi.org/10.1080/1091367X.2015.1054390>
21. Ware Jr JE, Gandek B. Overview of the SF-36 health survey and the international quality of life assessment (IQOLA) project. *Journal of Clinical Epidemiology*. 1998; 51(11): 903-912. [https://doi.org/10.1016/S0895-4356\(98\)00081-X](https://doi.org/10.1016/S0895-4356(98)00081-X)
22. Kalantar-Zadeh K, Kopple JD, Block G, Humphreys MH. Association among SF36 quality of life measures and nutrition, hospitalization, and mortality in hemodialysis. *Journal of the American Society of Nephrology*. 2001; 12(12): 2797-2806. <https://doi.org/10.1681/ASN.V12122797>
23. Czimbalmos Á, Nagy Z, Varga Z, Husztik P. [Patient satisfaction survey with SF-36 questionnaire, determination of normal values in Hungary]. *Népegészségügy*. 1999; 80(1): 4-19 (in Hungarian).
24. Ross A, Willson VL. Paired samples T-test, in basic and advanced statistical tests. In: Ross A, Willson VL., editors. *Basic and advanced statistical tests*. Rotterdam: SensePublishers. 2017. p. 17-19. [https://doi.org/10.1007/978-94-6351-086-8\\_4](https://doi.org/10.1007/978-94-6351-086-8_4)
25. Lilliefors HW. On the Kolmogorov-Smirnov test for normality with mean and variance unknown. *Journal of the American Statistical Association*. 1967; 62(318): 399-402. <https://doi.org/10.1080/01621459.1967.10482916>

26. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007; 116(9): 1081.  
<https://doi.org/10.1161/CIRCULATIONAHA.107.185649>
27. Sheeha BB, Granat M, Williams A, Johnson DS, Jones R. Does free-living physical activity improve one-year following total knee arthroplasty in patients with osteoarthritis: a prospective study. *Osteoarthritis and Cartilage Open*. 2020; 2(3): 100065.  
<https://doi.org/10.1016/j.ocarto.2020.100065>
28. Lützner C, Kirschner S, Lützner J. Patient activity after TKA depends on patient-specific parameters. *Clinical Orthopaedics and Related Research*. 2014; 472(12): 3933-3940.  
<https://doi.org/10.1007/s11999-014-3813-5>
29. Tsonga T, Kapetanakis S, Papadopoulos C, Papathanasiou J, Mourgiaris N, Georgiou N, et al. Evaluation of improvement in quality of life and physical activity after total knee arthroplasty in Greek elderly women. *The Open Orthopedics Journal*. 2011; 5: 343.  
<https://doi.org/10.2174/1874325001105010343>
30. Frimpong E, et al. Improvements in objectively measured activity behaviors do not correlate with improvements in patient-reported outcome measures following total knee arthroplasty. *The Journal of Arthroplasty*. 2020; 35(3): 712-719.  
<https://doi.org/10.1016/j.arth.2019.10.016>